

GREAT TRINITY FOREST

Hardwood Silviculture

Volume 28

Table of Contents

Section	Page #
Cover Crops Help Tree Seedlings Beat Weed Competition	1
A Comparison of Large-Scale Reforestation Techniques Commonly Used on Abandoned Fields in the Lower Mississippi Alluvial Valley	8
Restoring Bottomland Hardwood Forests: A Comparison of Four Techniques	15
An Afforestation System for Restoring Bottomland Hardwood Forests: Biomass Accumulation of Nuttall Oak Seedlings Interplanted Beneath Eastern Cottonwood	21
Early Response of Interplanted Nuttall Oak to Release from an Eastern Cottonwood Overstory	30
The Role of Large Container Seedlings in Afforesting Oaks in Bottomlands	35
A Comparison of Oak Regeneration Conditions Following Midstory Injection and Partial Overstory Removal in a Tombigbee River Terrace	42
Herbicide Release of 4 Year old, Naturally Regenerated Bottomland Oaks-10 Years Results	46
Oak Regeneration: Serious Problems Practical Recommendations	51
Seedling Care and Reforestation Standards	384
Literature Cited	401

Great Trinity Forest Management Plan

HARDWOOD SILVICULTURE

Cover Crops Help Tree Seedlings Beat Weed Competition

(Produced by the LandOwner Resource Centre and the Ontario Ministry of Natural Resources)



COVER CROPS HELP TREE SEEDLINGS BEAT WEED COMPETITION

Planting cover crops, such as rye grass or clover, can be an important step in turning fragile or marginal lands into productive forests. By slowing the spread of weeds, cover crops help seedlings compete for sunlight, nutrients and water, with reduced need for herbicides or other weed control.

Some cover crops also enrich and stabilize the soil. They are particularly useful in preventing slopes from eroding which, in turn, reduces the amount of silt carried into waterways. A

THE BENEFITS OF COVER CROPS


REDUCED HERBICIDE USE


Hardwood seedlings grow best when they have an ample supply of sunlight, nutrients and moisture. When other plants compete for the same resources, these seedlings rarely survive long enough to outgrow the competition.

Cover crops keep competing plants at bay, with reduced need for herbicides or other weed control. Because they grow low to the ground, cover crops give seedlings access to sunlight, while discouraging competing weeds from taking root. While it's true cover crops also compete with seedlings, they are not as damaging as weeds.

Field trials conducted by the Ministry of Natural Resources have shown that cover crops can cut herbicide use by as much as 50 per cent and the cost of chemical weed control by up to \$25 a hectare. As well, rodent damage to young seedlings was reduced to less than five percent of the trees, while tree survival rates were as high as 99 per cent. tractor, seeder and mower are needed to grow cover crops in large areas.

This Extension Note explains how to choose, plant and grow cover crops in order to improve the growth of hardwood seedlings.





WHEN TO PLANT COVER CROPS

Cover crops can be planted either one year before tree seedlings or during the same year.

When cover crops are planted the year before, soil erosion and water runoff are reduced. To effectively slow the growth of competing weeds, sow white clover early so that it becomes established before the trees are planted. If you are using machines to plant the trees, however, there are some disadvantages to early sowing. Tire traction can be reduced. Because cover crops hold soil moisture, there may also be fewer days in the spring when fine-textured soils are dry enough for planting.

If you plant cover crops during the same year as seedlings, using a machine to plant trees is easier. With this option, however, cover crops may not grow enough in the first year to prevent weeds from germinating. The use of a second, faster growing cover crop, such as winter wheat, timothy grass or barley, along with your primary cover crop can help solve this problem.

PREVENTING SOIL EROSION

The root systems of cover crops help prevent erosion by securing the soil. They are particularly useful on slopes where rain can carry soil into waterways and degrade water quality and fish habitat.

Cover crops can also improve the condition of the soil. Depending on the species, they can help loosen compacted soil and increase soil organic matter, moisture and fertility.

FOOD FOR WILDLIFE


Some cover crops, such as white clover and rye grass, are excellent winter foods for birds, small mammals and deer.

SELECTING A SITE

Cover crops thrive on slopes and flat areas, but they don't do well on low sites that may be prone to flooding.

CHOOSING COVER CROP SPECIES

When choosing the best cover crop for your tree plantation, it's important to know whether or not the cover crop produces its own seed or if new seed must be purchased and sown each year. Also, you should be aware that some cover crops can increase pest problems or attract rodents that might feed on the seedlings. One agricultural pest, for example, thrives on millet. The Ministry of Natural Resources has tested two cover crops traditionally used in agriculture — white clover and rye grass.



A COMPARISON OF THE BENEFITS OF TWO COMMON COVER CROP SPECIES

	WHITE CLOVER	RYE GRASS
Appearance	 There are several types of white clover, including wild white clover and Dutch white clover Ladino clover is not recommended for tree plantations because it grows to a height of 34 centimetres 	Grows rapidly in dense clumps
Weed Suppression	 Although it can take up to a year to become established, white clover suppresses weed germination 	 Becomes established quickly Its dense cover, large root mass and ability to absorb nitrogen, suppresses weeds, specially late annual weeds and quack grass
Life Span	Most varieties live five yearsSelf-seeding	 Annual rye grass dies in winter Perennial rye grass survives winter Self-seeding
Competition and Control	 Can spread into the vegetation-free areas around tree seedlings When necessary, growth can be controlled with herbicides 	 Can spread into the vegetation-free areas around tree seedlings Not easily controlled by herbicides
Soil Building	Increases soil fertility by producing nitrogenCan reduce frost heaving on fine-textured soil	 Increases soil organic matter Can reduce frost heaving on fine-textured soil
Wildlife	Excellent winter food for upland game birds, rabbits and deer	Excellent winter food for deer
Varieties and Amount Required	 Dutch white clover requires five kilograms of seed a hectare Wild white clover varieties, such as Grasslands Huia or New Zealand, require five kilograms of seed a hectare combined with five kilograms of the creeping red fescue variety Jasper 	The many varieties require eight kilograms of seed a hectare
Cost	About \$23 a hectare	About \$15 a hectare
Sowing	 Sow in early spring just as frost comes out of ground Areas that have adequate moisture can be seeded no later than mid-August Seedings in September or October rarely survive the winter When sowing, add the recommended nitrogen producing bacteria to the hopper 	 Sow in early spring for best results Areas that have adequate moisture can be seeded in summer Can be seeded from mid-August to mid-September when there is less competition from germinating weeds

	WHITE CLOVER	RYE GRASS
Disadvantages	 Can increase species of worms that feed on seedling roots if area around trees is not kept free of vegetation Can provide habitat for mice and other rodents that feed on seedlings 	Can provide habitat for mice and other rodents that feed on seedlings
Maintenance	 To reduce rodent habitat, mow in early September For best self-seeding, mow in summer when seeds are ready for dispersal Allow the plants to reach 10 centimetres in height before the first mowing 	 To reduce rodent habitat, mow in early September For best self-seeding, mow in summer when seeds are ready for dispersal
Site Conditions	 Low tolerance to drought Does not germinate well in wet areas Ideal soil pH is 5.1 to 7.5 	 Low tolerance to drought and heat Ideal soil pH is 5.5 to 7.5



Figure 1— Green ash seedlings are planted 2.4 metres apart. The cover crop, white clover, is grown in a 1.2 metre strip between the rows of trees. The 1.3 metre strips immediately surrounding the trees should be kept free of weeds by cultivating or applying herbicides. Use of cover crops in this manner reduces the amount of herbicide required in plantations by 50 per cent.

HOW TO PLANT COVER CROPS

Cover crops are planted with the same equipment used to plant other agricultural crops. To establish cover crops between rows of trees, follow these steps:

PREPARING THE SITE

Fields with silt loam to clay loam soils should be ploughed and disced in the fall, if you intend to plant a cover crop the following spring. Sandy loam and loam soils should be ploughed and disced in the spring, just before a cover crop is planted.

On fields where agricultural crops grew during the previous season or where erosion is a problem, cover crops can be sown directly over crop stubble.

SOWING

With a broadcast seeder, sow clover just as the frost is coming out of the ground in the spring. If seed is sown later in the season, ensure that there is enough moisture in the soil for germination. When seeding clover, add the recommended bacterial inoculant (available from seed suppliers) to the seed hopper before seeding. Follow the instructions on the inoculant package. The bacteria help to convert atmospheric nitrogen to forms that plants can use.

Calibrate your seeder. Broadcast the seeds according to the rates specified in Table 1.

PLANTING

If you are planting trees the year after the cover crop is sown, mow the cover crop in the fall. If a lot of weeds appear in the first season, make sure you mow the weeds before they go to seed. Raise the height of the mower so that it cuts all the weeds that are taller than the cover crop.

Plant trees in rows that are far enough apart to allow mowing equipment to cut the cover crop.

PROTECTING PLANTS FROM WEEDS

Weed growth should be controlled in a one-metre area around each seedling for the first three years of a seedling's life. This can be done by cultivating or by using the herbicide simazine.

If weeds are controlled manually, cultivate or hoe each spring and fall, and as required in the summer.

To control weeds with simazine, apply the herbicide immediately after planting. In addition to killing weeds, it prevents white clover and rye grass from germinating in the treated area. Apply simazine again each spring or fall for three years or until tree branches prevent access by an agricultural sprayer.

MAINTAINING COVER CROPS

Because mowing reduces habitat for rodents that feed on the tree seedlings over winter, be sure to mow at least once a year in early September. However, more frequent mowing may be necessary. Mowings that are done when the cover crop is going to seed will help broadcast the seeds. If ground nesting birds, such as bobolinks or field sparrows, are present, do not mow until the young birds have left the nest.



Cover crops should be mowed at least once a year. This reduces habitat for rodents that might feed on tree seedlings.

COVER CROP RESEARCH

The effectiveness of cover crops as tools for re-establishing hardwood forests in southern Ontario is being evaluated by the Ministry of Natural Resources. Field trials are being conducted through the Vegetation Management Alternatives Program, which is developing safe and effective alternatives to herbicide spraying. Field tests have already shown that cover crops reduce the need for herbicide spraying in young plantations.

For more information on cover crop evaluation in southern Ontario contact: Silvia Strobl, Plantation Forester Ontario Ministry of Natural Resources P.O. Box 605 Brockville, Ontario K6V 5Y8 phone (613) 342-8524 fax (613) 342-7544

For information on cover crop varieties and the cost of seeds, contact seed suppliers or your local office of the Ontario Ministry of Agriculture and Food, and Rural Affairs.



The Ministry of Natural Resources is investigating the use of cover crops, including commercially available mixes of grains and clovers, as a means of reducing weed competition with tree seedlings. The grains establish quickly in the first year providing both weed control and food for wildlife. By the second year, the clovers are densely established.

Produced by the LandOwner Resource Centre and the Ontario Ministry of Natural Resources, with financial assistance from Tree Plan Canada.



© 1994, Queen's Printer for Ontario Printed in Ontario, Canada 50515 ISSN 1198-3744 (10 k P.R., 95 01 30)

Cette publication est également disponible en français. Page 7 of 404

Great Trinity Forest Management Plan HARDWOOD SILVICULTURE

A Comparison of Large-Scale Reforestation Techniques Commonly Used on Abandoned Fields in the Lower Mississippi Alluvial Valley

(Presented at the Tenth Biennial Southern Silvicultural Research Conference, Shreveport LA, February 16-18, 1999.)

A COMPARISON OF LARGE-SCALE REFORESTATION TECHNIQUES COMMONLY USED ON ABANDONED FIELDS IN THE LOWER MISSISSIPPI ALLUVIAL VALLEY'

Callie Jo Schweitzer and John A. Stanturf²

Abstract-Reforesting abandoned land in the lower Mississippi alluvial valley has attracted heightened attention. Currently, federal cost share programs, such as the Wetland Reserve Program and the Conservation Reserve Program, are enticing landowners to consider reforesting lands that are marginally productive for agriculture. This study examined four reforestation techniques commonly used in the area. These techniques were applied operationally, following methods used under federal reforestation programs. Four techniques were implemented on **8.1-ha** blocks in a completely randomized design with three replications. The entire study site was prepared by disking following soybean harvest in the fall of 1994. The techniques tested were: (1) direct seeding **Nuttali** oak (*Quercus nuttallii* Palmer) acorns (2) **planting 1-0 bareroot Nuttali** oak seedlings (3) establishing a nurse crop of eastern cottonwood (*Populus deltoides* Marsh.) and then underplanting 1-O **bareroot Nuttali** oak seedlings (4) control (natural succession). All techniques were implemented by April 1995, except the underplanted oaks, which were planted following two growing seasons for the cottonwood (oak underplanted in March 1997). Results of this study are reported for 3 years of survival and growth of the seed germinants and seedlings, for cottonwood survival and growth, and interplanted oak survival and growth following 1 year. Natural invasion onto the site is also discussed. Comparisons are made among the four reforestation techniques, with ideas for incorporating this information into the administration of federal cost share reforestation programs.

INTRODUCTION

1

Large-scale reforestation of former agricultural lands in the lower Mississippi alluvial valley (LMAV) continues to attract interest. Lands that were cleared in the 1960's primarily for soybean production are undergoing a use shift again, this time back to trees. The decision by many landowners to reforest these lands has been aided, in part, by the increased availability of reforestation programs, such as the Wetland Reserve Program (WRP). This program provides the landowner with a one-time easement payment, technical expertise, and cost-share to cover part or all of the reforestation costs. Applying the information amassed for large-scale reforestation of hardwoods on former agricultural lands has been challenging. There are perhaps several reasons for this. Technology transfer outlets, such as written reports and other publications, may not be the most suitable methods of disseminating this information. Additionally, the target audience is diverse, and many of the practitioners are unknown.

Research continues on expanding our current reforestation knowledge base from a small to a larger scale. The basic techniques for introducing desirable hardwood species on former agricultural land have been worked out. However, much of the preliminary work has been done on a much smaller scale. Currently, large-scale reforestation is occurring on thousands of ha in the LMAV. Much of this work has been done under the Wetland Reserve Program, and seedling and acorn survival rates have been low. One objective of this study was to test four commonly used reforestation techniques and to monitor both growth and survival so that operational techniques can be adjusted accordingly.

In many reforestation projects in the LMAV, emphasis has been placed on hard-mast producing species such as the

oaks. Typical reforestation on recently abandoned fields involves the establishment of one to three overstory species, usually oaks. Sowing of acorns is a conventional method for **establishing new** stands of oak even though failures often occur. Researchers have indicated that planting 1-O **bareroot** nursery stock or direct seeding acorns can work well for most bottomland hardwood species planted on a variety of sites in the LMAV (Allen 1990, Baker and **Blackmon** 1973, Johnson 1983, Johnson and Krinard 1985, Krinard and Kennedy 1987).

This project was designed to test one alternative reforestation technique which combines a faster growing species, cottonwood (*Populus deltoids* Marsh.) and a slower growing species, **Nuttall** oak (*Quercus nuttallii* Palmer), and to contrast this technique with more traditional approaches of planting **bareroot** seedlings or direct seeding of acorns of **Nuttall** oak. The control treatment for this study is to do nothing and allow natural field succession to occur.

Compared to other reforestation techniques, one advantage of using the cottonwood nurse crop may be the creation of a more favorable microclimate for oak growth and survival. Obviously, we cannot test this until later in the study. The early growth of cottonwood allows for the rapid establishment of a forest canopy. The advantage of this canopy is that it may lend itself to accelerating natural succession by attracting birds and small mammals that are vectors for dispersal of heavy seed.

The major disadvantage of pure cottonwood plantations to wildlife may be the paucity of hard mast. Although some may also feel that cultivation works against restoration goals, it has been found that wildlife importance values for all wildlife food plants in several cottonwood plantations studied peaked in the fourth, **fifth** and sixth growing seasons

¹ Paper presented at the Tenth Biennial Southern Silvicultural Research Conference, Shreveport, LA, February 16-18, 1999.

² Research Forester, USDA Forest Service, Southern Research Station, Forest Inventory and Analysis Research Work Unit, Asheville, NC 28802; and Research Soil Scientist, USDA Forest Service, Southern Research Station, Center for Bottomland Hardwoods Research, Stoneville, MS 38776, respectively.

(Wesley and others 1981). The planting scheme under study here will provide for hard mast, and **cultivation** cessation after establishment will hopefully serve to promote plant establishment that would benefit wildlife and restoration in the later growing seasons.

In addition to providing wildlife habitat, the high survival rates and rapid growth of cottonwood will enable harvest in 10 years, generating income in a relatively short time period. **Stanturf** and others (1998) conducted a financial analysis for a private landowner of a cottonwood plantation and found that the cottonwood provided a cash flow over the rotation, with a positive internal rate of return. Data collected from the current study will allow for future costs analysis with the additional returns of the oak together with the cottonwood.

OVERVIEW AND METHODS

Four reforestation techniques have been installed on former agricultural land located in Sharkey County, Mississippi. The site is 2.5 km east of the community of Anguilla, approximately 1 km south of State Highway 14, and immediately north of the Delta National Forest. This land was transferred to the **USDI** Fish & Wildlife Service from the Farmers Home Administration in 1993, and is administered by personnel at the Yazoo National Wildlife Complex.

A recent soil survey of the tract conducted by the Natural Resources Conservation **Service(NRCS)** indicated that the soil is of the Sharkey series, a heavy clay with shrink-swell properties, described as a very fine, montmorillonitic, **nonacid,** thermic **Vertic** Haplaquepts (Personal communication. Floyd Wood. 1995. Soil biologist, NRCS, 100 Capitol St., Jackson, MS 39289). The Sharkey series consists of poorly drained, clayey soils formed in fine textured sediment in slack water areas along the Mississippi River. The shrink-swell nature typical of clays with montmorillinitic mineralogies results in 2-10 cm wide cracks up to 1.5 m deep that form under dry conditions, and close upon wetting. The hydrologic and edaphic conditions of our study site typify most of the abandoned agricultural land available for and undergoing reforestation in this region.

The experimental design follows a randomized complete block design with three replicates located in different portions of the tract. Treatment plots are approximately rectangular in shape and 8.1 ha in size. Treatment plots were established in October 1994 (Blocks II and III) and February 1995 (Block I). The assignment of treatments to plots was done using a random numbers table, and is summarized in table **1**.

Agricultural production on the study site ended in fall 1994. The entire study site was double **disked** (including the natural succession treatment) following soybean removal. Acorns were collected in the Delta National Forest, placed in water and non-viable acorns that floated were discarded. Acorns were then stored in ventilated polyethylene bag at 1.7 °C. The Fish & Wildlife Service conducted the planting/sowing using their standard equipment and techniques. In May 1995, acorns were machine sown at 1 **.1** m X 3.7 m spacing, with one acorn placed at each planting spot. Oak seedlings **(1-0** stock) were obtained from Fratesi Nursery, Leland, MS (seed source Delta National Forest). Seedlings were machine planted in March 1995 at 3.7 m X 3.7 m spacing. Table I-Treatment plot assignment in each block

	Block								
Treatment plot	ls.	11	111						
1 2 3 4	PLN NAT NUR sow	sow N A T PLN NUR	NUR s o w NAT PLN						

PLN = Plant bare-root Nuttall oak seedlings (1-O stock),

NAT = natural old field succession, NUR = cottonwood/Nuttall oak intercrop, SOW = direct seeded Nuttall oak acorns.

Four cottonwood clones (ST-88, ST-72, ST-76, and **S7C-1**) were established using plantation establishment procedures practiced by Crown Vantage, Inc. in spring 1995 (table 2). Cottonwood cuttings were hand-planted in pure blocks at 3.7 m X 3.7 m spacing in March 1995. Each cottonwood clone **area (2** ha per clone per block) was split, with one-half of the area (1 ha) receiving weed control by disking in 1995 only, and the other half receiving the same weed control in 1995 and 1996.

In March 1997, the underplanted **Nuttall** oak seedlings were planted underneath the cottonwood, in every other row (3.7 m X 7.4 m spacing). Additional seedlings were also planted in **0.4-ha** blocks in the open fields adjacent to each cottonwood plot. Survival, height, and diameter data were collected in 1997, following one growing season.

Four permanent measurement plots were installed in each treatment plot in fall 1995. For the direct seeded and planted treatments, measurement plots were rectangular, approximately 0.2 ha in size. Sample measurement plots for cottonwood clones were also 0.2 ha and placed, by clone, in the 1995 and the 1995-96 **disked** areas. All cottonwood, planted and direct seeded spots were flagged in each treatment measurement plot. The control areas were sampled by installing 64 circular plots (6.45-m radius). This sample area equals 0.85 of the total ha sampled for each control treatment plot, which is comparable to the total area sampled in the direct seeded and planted measurement plots (0.2 ha X 4 quadrants). Height and diameter data have been collected for all woody stems in all treatments following the third growing season **(1** 997).

Statistical analysis followed a Model I **ANOVA.** The SAS statistical package (SAS Institute, Inc. 1990) was used for data analysis, incorporating Duncan's New Multiple Range test for mean comparisons. All significant differences are reported at a \leq 0.05.

RESULTS

Seedling and Acorn Germinant Plots

The seedling height and diameter growth increased significantly over 3 years (table **3). Seedling** survival was 64 percent after 3 years, averaging 489 trees per ha.

Table P-Schedule of operations for cottonwood/Nuttall treatment

.

Dates	Activity
October 1994	Two-pass site preparation disking Row establishment and liquid nitrogen applied in trenches
March 1995 March 1995	 @ 112 kg N ha⁻¹ Plant cottonwood Spray herbicide in band over dormant cuttings (oxyfuorfen @ 0.26 kg ha⁻¹ + glyphosate
May 1995	① 1.4 kg ha-') One-pass disking, followed 2 weeks later by second pass
June &July 1995	at right angle to first Basal application of oxyfluorofen @ 0.7 kg ha"
August 1995	One-pass disking, followed 2 weeks later by second pass at right angle to first
Summer 1995	Insect control for cottonwood leaf beetles (carbanyl @ 0.92 kg ha-')
June 1996	Insect control for cottonwood leaf beetles (carbanyl @ 0.92 kg ha")
June & July 1996 March 1997	One-pass disking Plant Nuttall oak seedlings at offset position from cottonwood
Winter 2004	Cottonwood pulpwood harvest

Table **3—Seedling** and acorn germinant height, diameter, and survival comparisons among growing seasons

		Year	
Variable	1997 ^a	1996	1995
Seedling height (cm) Seedling diameter (mm) Seedling survival (percent) Germinant height (cm) Germinant diameter (mm) Germinant survival (percent)	27.4a 3.4a	52.9b 6.2b 59.6a 18.4b 2.5b 10.6b	1.3c .lc 62.8a .5c .lc 17.6a

^eDifferent letters within rows indicate significantdifferences among means: average n = 1135 for seedlings: average n = 530 for germinants.

Acorn germinant survival was 16 percent, which was not significantly different from the 1995 or 1996 survival percentages (table 3). The 1995 acorn germinant survival was greater than that of 1996, and 1996 survival was less than 1997 survival. The most likely explanation for these discrepancies was sampling error. Acorn germinants averaged 225 stems per ha after 3 years. Height and diameter of germinants increased consistently over the three growing seasons.

Control Plots

In 1996, in a total of 0.85 ha sampled in each control plot, seven trees were found in Block 1 (five green ash *Fraxinus pennsylvanica* Marsh., two cedar elms *Ulmus* crassifolia **Nutt.)**, five in Block 2 (four green ash, one cedar elm) and five in Block 3 (one green ash, four cedar elms). More volunteer trees were found in 1997 (table 4). In 1997, the average number of trees per ha was **195, 25** and 20 per block. The high number of swamp dogwoods (Comus foemina Mill.) found in Block 1 heavily weighted this average.

Cottonwood Plots

After three growing seasons, the clones displayed differences in growth, with ST-66 > S7C-1 > ST-75 > ST-72 for both height and diameter (table 5). Height and diameters were greater for those cottonwood clones that received 1 year of weed control compared to those that were **disked** in 1995 and 1996 (table 6). The results for survival among clones by weed control regime are given in table 7.

Cottonwood-Red Oak: Underplanted and Open-Grown Oaks

Following one field growing season, the underplanted oaks were significantly taller (48.2 cm for underplanted, 43.9 cm for open grown) and had greater average diameters (6.1 mm for underplanted, 7.7 mm for open grown) compared to those grown in the adjacent open fields. Survival for underplanted oaks averaged 57 percent after one growing season, while open-grown oaks had a 43 percent survival rate.

Table 4-Number and species of volunteer trees found on control plots, total of 0.85 ha sampled for each block; Sharkey Research Site, **1997**

	Block						
Таха		II	III				
Fraxinus Pennsylvania Green ash	11	14	2				
Ulmus crassifolia	16	5	5				
Cedar elm Comus stricta	125	0	6				
Swamp dogwood Diospyros Virginia Persimmon	7	0	0				
Celtis laevigata	8	2	52				
Sugarberry' <i>Gieditsia triacanthos</i> Honeylocust	0	0	1				

Table	5Height,	diameter,	and	total	number	of	sprout	comparisons	
among	cottonwood	clones	for 1	1997 (data				

Clone type	Diameter ^a	Height	Sprouts		
	ст	т	Number		
ST-66 S7C-1 ST-75 ST-72	10.1a 9.0b 8.7c 8.1d	9.49a ^{8.71 b} 7.85c 7.49d	1.2c 1.3b 1.5a 1.4a		

⁸ Different letters within columns indicate significant differences among means; average n = 815.

Table **6—Height,** diameter, total number of sprouts, and survival comparisons of means between **1** year (1995) and 2 years (199586) of weed control, by clone, following three growing seasons (1997 data)

Variable	Weed control	S7C-1*	ST-66	ST-72	ST-75
	Year(s)				
Height(m) Diameter (cm) Sprouts (no.) Survival (percent)	1995 199596 1995 1995-96 1995-96 1995-96 1995 1995-96	6.9a 5.8b 7.1a 5.8b 1.7a 1.7a 97.6a 94.4a	7.6a 6.0b 8.1a 6.2b 1.6a 1.5b 96.4b 95.1a	5.9a 5.5b 6.1a 5.6b 1.5a 1.5a 88.6a 82.2a	6.6a 5.6b 7.1a 5.8b 1.7a 1.6b 93.2a 92.8a

⁸Different letters within columns, by clone, indicate significant differences between means.

3

1948 - 195 MA

Table 7-1997 cottony control regime-l year		among clones by weed (199596) of weed control
	199596 weed	
Clone	control'	control
	••••• P	ercent
S7C-1 ST-66 ST-72 ST-75	97.6a 96.4ab 88.6c 93.2b	94.4ab 95.1a 82.2b 92.8ab

^eDifferent letters within columns indicate significant differences among means,

٦

11

DISCUSSION

After three growing seasons, results from these four reforestation techniques are beginning to shed some light on expected reforestation on this site. It is understandable that the cottonwood growth and survival are superior to that of the oaks. Not only is cottonwood the fastest growing tree species in North America, but also the care and attention given to its establishment are displayed in the results. Seedling survival was approximately as expected for a **large**-scale operation. At an average of **489** trees per ha, seedling establishment would be deemed above adequate under guidelines set by NRCS for federal tree establishment programs, especially WRP. Under WRP, 308 hard mast stems per ha after three growing seasons meets the success criteria. Therefore, the 225 trees per ha for the acorn germinants would fall short of this goal.

Seedling survival, compared to other similarly reforested sites, was average to below average. Allen (1990) evaluated oak plantations established by the **USDI** Fish & Wildlife Service personnel on refuges in west-central Mississippi. Seven out of ten stands Allen assessed had over 489 trees per ha. Krinard and Kennedy (1987) observed survival **rates** of hardwood seedlings planted on Sharkey clay soil from 69 to 97 percent after 2 years, and 57 to 98 percent after 4 years. They found that **Nuttall** oak seedling survival was 85 percent (2 years) and 80 percent (4 years). Wittwer (1991) observed 78 percent survival of planted bottomland oaks after 3 years. Savage and others (1989) reported 370 trees per ha and a 64 percent survival rate for both seedlings and germinants on reforested bottoms in Louisiana.

To date, obtaining adequate stocking of oaks on good to poor sites has been difficult with either natural or artificial regeneration. It has been suggested that the inherent sluggish growth habitat of oak is the principal cause of regeneration failures. This slow growth rate combined with intensive vegetative competition and frequent animal browsing has plagued forest managers in their attempts to regenerate oaks. Based on research trials and our knowledge of direct seeding on public and private lands in recent years, the overall likelihood of regeneration success with oaks in a given year is somewhat less with direct seeding than with planting **(Bullard** and others 1992).

Before acorns germinate, predation from mammals and birds constitutes the major cause of loss (Harmer **1994a**, b; Korstian 1927). A pilot small mammal trapping study (600 trap nights) conducted on a single plot of the direct seeded and the cottonwood treatments in October 1995 indicated that very dense small mammal populations were present on the direct seeded treatment (202 animals captured 216 times) compared to the cottonwood (2 animals captured 2 times). Sigrnodon *hispidus* was the most abundant of five species captured (Willis and others 1996). Therefore, rodent predation of acorns may have contributed to the lower survival rates and stocking in the direct seeded treatments, which had extensive weed cover compared to the bare soil conditions in the cottonwood treatments.

The expected survival of operationally direct seeded acorns has been reported at 35 percent (Johnson and Krinard 1985, Kennedy 1993). Theoretically, a large number of acorns should increase the possibility for some acorns to escape predation and germinate. Others have experienced this same problem and have addressed it by simply increasing the number of seeds sown. For example, Willoughby and others (1996) described results and recommendations from several studies designed to establish new **woodlots** by direct sowing. Their recommendation for acorns was to sown 10,000 per ha, with the aim of establishing 1,000 reasonably even-spaced vigorous trees by year 10. This figure allowed for losses from germination failure, predation, drought, herbicide damage, and weed competition. This planting density was also observed by the authors on State-run forests in Denmark, where the sowing and subsequent germination and growth of oak was so thick that it appeared almost like a shrub row.

The survival and stocking on these reforested sites must be viewed in the context of the landowners' objectives. If having a few widely spaced trees in fields dominated by herbaceous vegetation is the goal, it appears as if the current direct seeding or natural succession methods will work. However, if the objective is to restore these sites back to some level of a functional forested wetland, regardless of the function, more time and effort are needed to implement the techniques. The general knowledge of reforestation on old field sites exists; it now becomes a matter of using that information property to get the desired results. Perhaps the biggest challenge is not working out the individual aspects of large-scale reforestation but putting all those pieces together in one scheme. We can only continue to match species to sites, use proper seed and seedling handling, strive for good quality planting stock, perform site preparation, and perhaps most importantly, have on-site supervision to help make the pieces fit.

Plantations of fast growing tree species in short rotation cycles are going to play a vital role in meeting the rising demand for woody biomass production. Populus has shown promising growth and productivity, can be harvested in short periods, and has combined well, thus far, with the red oak intercropping system. Twedt and Portwood (1997) noted that in addition to production-related benefits, planting earlysuccessional species can (1) promote rapid colonization by migrant birds (2) enhance plant species diversity (3) provide a more rapid financial return to landowners, and (4) enhance the public's perception of reforestation efforts. The ongoing research into physiological response of oaks established beneath cottonwood, as well as the possible changes in the edaphic and microclimate environment under the cottonwoods, will aid in our prognosis of implementing such a system.

ACKNOWLEDGMENTS

Funding for this study came in part from the National **Council** of the Paper Industry for Air and Stream Improvement. The authors wish to acknowledge the input of Jim Shephard, NCASI, Jeff Portwoad, Crown Vantage, Inc., Tim Wilkins, Lamar Dorris and Ronnie Haynes, **USDI** Fish &Wildlife . Service, Dexter Bland, Emile Gardiner, Paul **Hamel**, Davld Hoge, and Todd Parker, USDA Forest Service.

REFERENCES

Allen, J.A. 1990. Establishment of bottomland oak plantations on the Yazoo National Wildlife Refuge Complex. Southern Journal of Applied Forestry. 14(4): 206-210.

Baker, J.B.; **Blackmon**, B.G. 1973. Summer fallowing helps establish cottonwood on old**fields**. Mississippi Agriculture and. Forestry Experiment Station Research Highlights.**36(6)**: 5.

- Bullard, D.; Hodges, J.D.; Johnson, R.L.; Straka, T.J. 1992. Economics of direct seeding and planting for establishing oak stands on old-field sites in the South. Southern Journal of Applied Forestry. 16(1): 34-40.
- Harmer, R. 1994a. Natural regeneration of broadleaved trees in Britain: I. Historical aspects. Forestry. 67: 179-188.
- Harmer, R. 1994b. Natural regeneration of broadleaved trees in Britain: II. Seed production and predation. Forestry. 67: 275-286.
- Johnson, R.L. 1983. Nuttall oak direct seeding still successful after 11 years. Res. Note SO-301. New Orleans: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 3 p.
- Johnson, R.L.; Krinard, R.M. 1965. Oak seeding on an adverse site. Res. Note SO-319. New Orleans: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 p.
- Kennedy, H.E. 1993. Artificial regeneration of bottomland oaks. In: Loftis, David L.; McGee, Charles E.. eds. Oak regeneration: serious problems, practical recommendations: Symposium proceedings; 1992 September 8-10; Knoxville, TN. Gen. Tech. Rep. SE-64. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 3 p.
- Korstlan, C.F. 1927. Factors controlling germination and early survival in oaks. Bull.19. New Haven, CT: Yale University, School of Forestry: 115.
- Krinard, R.M.; Kennedy, H.E. 1987. Fifteen-year growth of six planted hardwood species on Sharkey clay soil. Res. Note SO-336. New Orleans: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 p.

- SAS Institute Inc. 1990. SAS user's guide: statistics. Version 6. 4th ed. Carv. NC: SAS Institute Inc. [Number of pages unknown].
- Savage, L.; Prftchett, D.W.; DePoe, C.E. 1989. Reforestation of a cleared bottomland hardwood area in northeast Louisiana. Restoration and Management Notes. 7(2): 88.
- Stanturf, J.A.; Schweitzer, C.J.; Gardiner, E.S. 1998. Afforestation of marginal agricultural land in the Lower Mississippi River Alluvial Valley. U.S.A. Silva Fennica. 32(2): 281-297.
- Twedt, D.J.; Portwood, J. 1997. Bottomland hardwood reforestation for neotropical migratory birds: are we missing the forest for the trees? Wildlife Society Bulletin. 25(3): 647-652.
- Wesley, D.E.; Perkins, C.J.; Sullivan, AD. 1981. Wildlife in cottonwood plantations. Southern Journal of Applied Forestry. 5(1): 37-42.
- Willis, K.; \Voodson, C.; Hamel, P.H. 1998. Small mammal populations on agricultural land in Mississippi Alluvial Valley restored to forests: first year results. In: Proceedings of the Mississippi chapter of the wildlife society annual meeting; 1996 October 17; Jackson, MS. [Place of publication unknown]: [Publisher unknown]: 11-12.
- Willoughby, I.; Kerr, G.; Jinks, R.; Gosling, P. 1996. Establishing new woodlands by direct sowing. For. Comm. Res. Inf. Note-285. Edinburgh, UK: Forestry Authority Research Division. 7 p.
- Wittwer, R.F. 1991. Direct-seeding of bottomland oaks in Oklahoma. Southern Journal of Applied Forestry **15(1)**: 17-22.

Great Trinity Forest Management Plan

HARDWOOD SILVICULTURE

Restoring Bottomland Hardwood Forests: A Comparison of Four Techniques

(IUFRO Division 1 Conference – Book of Abstracts)

RESTORING BOTTOMLAND HARDWOOD FORESTS: A COMPARISON OF FOUR TECHNIQUES

John A <u>Stanturf</u>^{*1}, Emile S. Gardiner², James P. Shepard³, Callie J. Schweitzer⁴, C. Jeffrey Portwood⁵, and Lamar Dorris⁶

 ¹ USDA Forest Service, Stoneville, MS USA
 US Forest Service, 320 Green Street, Athens, GA 30602 USA; tel. 001-706-559-4316; email <u>istanturf@fs.fed.us</u>
 ² USDA Forest Service, Stoneville, MS USA
 ³ National Council for Air and Stream Improvement, Gainesville, FL
 ⁴ US Forest Service, Normal AL, USA
 ⁵ Temple-Inland Forest, Diboll. TX, USA
 ⁶US Fish and Wildlife Service, Hollandale, MS, USA

Introduction

Large-scale afforestation of former agricultural lands in the Lower Mississippi Alluvial Valley (LMAV) is one of the largest forest restoration efforts in the world and continues to attract interest from landowners, policy makers, scientists, and managers. The decision by many landowners to afforest these lands has been aided in part by the increased availability of public and private incentive programs such as the Wetlands Reserve Program (WRP). The WRP provides a landowner with a one-time easement payment, technical expertise, and reimbursement for part or all of the afforestation costs. Large-scale afforestation is occurring on thousands of hectares in the LMAV (King and Keeland 1999; Stanturf et al. 2000; Schoenholtz et al. 2001).

Early results from the WRP were discouraging (Stanturf et al. 2001a); seedling and acorn survival rates were low, despite much available information on planting and direct seeding techniques (Stanturf et al. 1998). Although the basic techniques for afforesting native hardwood species have been worked out, few studies compared several techniques on the same site. In response to questions from managers, we undertook a study to compare operational techniques for afforesting bottomland hardwoods. In addition to standard approaches of planting bare-root seedlings and direct seeding acorns, we included an interplanting technique using a fast-growing, native species (*Populus deltoides* Bartr. ex Marsh.) with an oak species (*Quercus nuttallii* Palmer) and another treatment of doing nothing and depending upon natural invasion. The specific objectives of the study were to demonstrate and compare four restoration techniques in terms of survival, accretion of vertical structure, and species diversity.

Methodology

The study was located in Sharkey County, MS (N32°58' W90°44'), in the Yazoo River Basin. The land was actively cropped until the study was established. The hydrologic and edaphic conditions of the study site were typical of land available for restoration in LMAV. Soils were mapped as the Sharkey series of very-fine, smectitic, thermic chromic Epiaquerts. Sharkey soils consist of poorly drained clays formed in fine textured sediment in slack water areas in the Mississippi River floodplain. The shrink-swell nature of Vertisols results in 2 to 10 cm wide cracks up to 1.5 m deep that form under dry conditions, and close when saturated.

The experiment was a randomized complete block design with three replicates located in different portions of the tract. Treatment plots were 8.1 ha and approximately rectangular. Treatments were chosen to represent a gradient in restoration intensity, from natural invasion, direct seeding Nuttall oak, planting Nuttall oak, to interplanting eastern cottonwood with Nuttall oak. The natural invasion treatment was the baseline to compare passive versus active restoration. Direct seeding and planting bareroot seedlings are routinely used restoration techniques in the LMAV under federal incentive programs (Stanturf et al. 2000). The interplanting technique combines a fast growing species, eastern cottonwood, as a nurse crop for the slower growing oak. The cottonwood can be harvested in as little as 10 years, providing a financial return to the landowner (Stanturf and Portwood 1999).

Direct seeding and planting treatments were installed in February 1995 and the cottonwood was planted in the interplanting treatment in March 1995. Acorns for the direct seeding treatment were collected from nearby natural stands, placed in water and non-viable acorns that floated were discarded. Acorns were stored in ventilated polyethylene bags at 1.7 °C until sowing. Acorns were machine sown by Fish and Wildlife Service staff in May 1995. Spacing was 1.1 m by 3.7 m, with one acorn placed at each planting spot. Bareroot 1-0 Nuttall oak seedlings were obtained from a commercial nursery. Oak seedlings were machine planted by Fish and Wildlife Service staff in March 1995 at 3.7 m by 3.7 m spacing.

Eastern cottonwood cuttings were hand planted according to procedures used operationally by forest industry (Stanturf et al. 2001b). Four commercially available clones were planted at 3.7 m by 3.7 m spacing. Three clones had been selected from native populations along the Mississippi River (ST66, ST72, ST75) and one was from an east Texas population (S7C1); all four clones are used operationally. Clonal material was provided by Crown Vantage (now Tembek) and grown in their nursery at Fitler, MS. Two growing seasons later (March 1997), Nuttall oak seedlings were interplanted under the cottonwood. Oaks were planted between every other cottonwood row so the cottonwood can be harvested without damage to the oaks. Spacing for these oak seedlings was 3.7 m by 7.4 m.

Four permanent measurement plots were installed in each treatment plot in autumn 1995. Survival, height and diameter growth were measured in the active restoration treatments annually through the fifth growing season. Height and diameter data were collected for all woody stems in the natural invasion treatment annually beginning with the second growing season. The interplanted oak seedlings, planted after the second growing season of the overstory, were measured annually.

Results and Discussion

There were no significant differences in soil chemical properties at any depth between the four restoration treatments. Soil bulk density was not significantly different for the treatments, although there were significant differences between blocks. After five growing seasons, the cottonwoods were the tallest trees and had the greatest density. Height of planted oaks averaged almost 1.4 m and was significantly taller than the direct seeded and interplanted oaks, which did not differ. Even though the direct-seeded oaks had been growing on the site two years longer than the interplanted oaks, they were not significantly taller. The direct-seeded oaks had greater diameter than the interplanted and they accumulated more biomass

The abandoned soybean field was invaded over time by woody species disseminated by wind and by birds. Swamp dogwood (*Cornus stricta*), common persimmon (*Diospyros virginiana*), green ash (*Fraxinus pennsylvanica*), sugarberry (*Celtis laevigata*), American elm (*Ulmus Americana*), hawthorns (Craetaegus species), cedar elm (*Ulmus crassifolia*), honeylocust (*Gleditsia triacanthos*), and deciduous holly (*Ilex decidua*) were the woody species found in the measurement plots.

This project was designed to test one alternative afforestation technique that combines a faster growing species with a slower growing species and to contrast this technique with more traditional approaches of planting bareroot seedlings or direct seeding of acorns. The control treatment for this study is to do nothing and allow natural invasion to occur. The early growth of cottonwood allowed for the rapid establishment of a forest canopy. The advantage of this canopy is that it may lend itself to accelerating natural succession by attracting birds and small mammals that are vectors for dispersal of heavy seed. The major disadvantage of pure cottonwood plantations to wildlife may be the paucity of hard mast. Although some may also feel that the intensive cultivation needed to establish cottonwood works against restoration goals, other studies have found that wildlife importance values for all wildlife food plants in cottonwood plantations studied peaked in the fourth, fifth and sixth growing season. The interplanting scheme under study here will provide for hard mast; once cultivation ceases after establishment, other herbaceous plants will establish.

Literature Cited

King, S.L. and B.D. Keeland. 1999. Restoration Ecology 7: 348-359.

Noss et al. 1995

Schoenholtz, S.H., J.P. James, R.K. Kaminski, B.D. Leopold, and A.W. Ezell. 2001. Wetlands 21: 602-613.

Stanturf, J.A., E.S. Gardiner, P.B. Hamel, M.S. Devall, T.D. Leininger, T.D. and M.L. Warren, Jr. 2000. Journal of Forestry 98: 10-16.

Stanturf, J.A. and C.J. Portwood. 1999. Pages 66-72 *in* J.D. Haywood, editor, Proceedings tenth biennial southern silvicultural research conference. U.S. Department of Agriculture Forest Service, Southern Research Station, Asheville, NC. General Technical Report SRS-30.

Stanturf, J.A., C.J. Schweitzer, and E.S. Gardiner. 1998. Silva Fennica 32:281-297.

Stanturf, J.A., S.H. Schoenholtz, C.J. Schweitzer, and J.P. Shepard. 2001a. *Restoration Ecology* 9:189-200.


Stanturf, J.A., C. van Oosten, M. Coleman, D. Netzer, and C.J. Portwood. 2001b. Pages 153-206 in D. Dickmann, J. Isebrands, and J. Richardson, editors, *Poplar Culture in North America*. National Research Council Press, Ottawa, Canada.

Meeting the challenge: Silvicultural Research in a Changing World

June 14-18, 2004 Hotel Mercure, La Grande Motte Montpellier, France

IUFR

IUFRO Division 1 Conference Program Book of abstracts



IUFRO Division 1 conference – Book of abstracts Edited by John Parrotta, Henri-Félix Maître, Daniel Auclair, Marie-Hélène Lafond CIRAD, UMR AMAP, TA 40/PS2, 34398 Montpellier cedex 5, France June 2004

Great Trinity Forest Management Plan

HARDWOOD SILVICULTURE

An Afforestation System for Restoring Bottomland Hardwood Forests: Biomass Accumulation of Nuttall Oak Seedlings Interplanted Beneath Eastern Cottonwood

(December 2004 Restoration Ecology Vol. 12 No. 4, pp.525-532)

An Afforestation System for Restoring Bottomland Hardwood Forests: Biomass Accumulation of Nuttall Oak Seedlings Interplanted Beneath Eastern Cottonwood

Emile S. Gardiner,^{1,2} John A. Stanturf," and Callie J. Schweitzer⁴

Abstract

Bottomland hardwood forests of the southeastern United States have declined in extent since European settlement. Forest restoration activities over the past decade, however, have driven recent changes in land use through an intensified afforestation effort on former agricultural land. This intense afforestation effort, particularly in the Lower Mississippi Alluvial Valley, has generated a demand for alternative afforestation systems that accommodate various landowner objectives through restoration of sustainable forests. We are currently studying an afforestation system that involves initial establishment of the rapidly growing native species eastern cottonwood (Populus deltoides Bartr. ex Marsh.), followed by enrichment of the plantation understory with Nuttall oak (Quercus nuttallii Palm.). In this article, we examine the growth and biomass accumulation by Nuttall oak seedlings to determine whether this species can be established and whether it will develop beneath the cottonwood overstory. After 3 years of growth beneath cottonwood canopies, Nuttall oak seedlings were similar in height (126 cm), but were 20% smaller in root-collar diameter than seedlings established in open fields. Seedlings established in the

open accumulated more than twice the biomass of seedlings growing beneath a cottonwood canopy. However, the relative distribution of accumulated biomass in seedlings did not differ in the two environments. Ten percent of total seedling biomass was maintained in leaf tissue, 42% was maintained in stem tissue, and 48% was maintained in root tissue on open-grown seedlings and seedlings established in the understory of cottonwood plantations. Though establishment in the more shaded understory environment reduced Nuttall oak growth, seedling function was not limited enough to induce changes in plant morphology. Our results suggest that an afforestation system involving rapid establishment of forest cover with a quick-growing plantation species, followed by understory enrichment with species of later succession, may provide an alternative method of forest restoration on bottomland hardwood sites and perhaps other sites degraded by agriculture throughout temperate regions.

Key words: forest restoration, *Populus deltoides*, *Quercus nuttallii*, regeneration, wetland forest.

Introduction

Forest restoration activities have become prevalent around the globe as marked by an afforestation rate of about 1.6 million hectares per year between 1990 and 2000 (FAO 2001). Of critical importance to forest conservation and sustainability in the temperate zone of North America is the restoration of bottomland hardwood forest ecosystems. Bottomland hardwood forests are deciduous and associated with alluvial soils on the floodplains of rivers and streams that dissect the southern and eastern United

¹USDA Forest Service, Southern Research Station, Center for Bottomland Hardwoods Research, Stoneville, MS 38776, U.S.A.

²Address correspondence to E. S. Gardiner, email egardiner@fs.fed.us

³USDA Forest Service, Southern Research Station, Forestry Sciences

Laboratory, Athens, GA 30602, U.S.A.

⁴USDA Forest Service, Southern Research Station, Bent Creek Experimental Forest, Normal, AL 35762, U.S.A.

© 2004 Society for Ecological Restoration International

States. The largest contiguous area of bottomland hardwood forest originally occurred in the Lower Mississippi Alluvial Valley (LMAV), but forest cover in this region has been reduced to an estimated 26% of the former 10.1 million-ha forest (Stanturf et al. 2000; Gardiner & Oliver, in press). In the last decade, a concentrated forest restoration effort has developed in the LMAV where afforestation activities have reestablished native tree species on more than 193,000 ha of degraded agricultural land marginally suited for crop production (Stanturf et al. 1998; Schoenholtz et al. 2001; Gardiner & Oliver, in press). The ultimate success of this massive forest restoration effort will be determined by the basic implementation of sound afforestation practices.

Interests in carbon sequestration, game habitat enhancement, timber production, water quality protection, and conservation of biological diversity are among the driving forces behind afforestation activities in the LMAV (King & Keeland 1999). Basic silvicultural practices for afforestation of bottomland hardwood forests **were** developed and have been applied since the 1960s (Allen 1990; Newling 1990). Conventional practices, which generally evolved from earlier techniques for establishing single-species plantations, have proven successful for some afforestation needs, but there is a need for alternative afforestation practices that provide additional options for achieving various resource management objectives (Gardiner et al. 2002). Development of afforestation systems that quickly produce complex vertical forest structure will provide alternative pathways to attain various restoration objectives.

Recently, plantations have been used effectively to restore and conserve tropical forests (Parrotta et al. 1997). Some of these systems have demonstrated value for restoring particular ecological processes or ecosystem functions. Several authors, including Parrotta (1995), Lugo (1997). and Powers et al. (1997), provide examples of plantations catalyzing natural forest regeneration and tree species richness on degraded sites. Plantations are often used to facilitate restoration of native forest cover on degraded tropical sites because they provide rapid development of forest structure attractive to wildlife species, and an associated understory microenvironment favorable to germination and establishment of native tree species (Parrotta 1992; Keenan et al. 1997; Lugo 1997). Other workers have extended the application of plantations for forest restoration by establishing rapidly growing species that develop an understory microenvironment favorable to establishment of planted seedlings of late-successional species (Ashton et al. 1997, 199X). Application of similar plantation establishment approaches involving fast-growing canopy species and understory enrichment plantings has not been widely studied or used for forest restoration in temperate regions (Ashton et al. 1997).

We are currently investigating the development and application of an interplanting system for restoration of bottomland hardwood forests on former agricultural fields in the LMAV. This restoration sequence is initiated by establishing the native species eastern cottonwood (Populus deltoides Bartr. ex Marsh.) to rapidly develop forest structure on the site. Following the establishment of the cottonwood plantation, the understory is enriched by planting Nuttall oak (Quercus nuttallii Palm.), a hardmast-producing, disturbance-dependent species of later successional seres. Initial results indicate that this interplanting system has potential as an alternative afforestation practice that can address multiple forest restoration objectives, for example, rapid forest biomass production and quick assembly of bird species richness (Hamel 2003). However, the ability of Nuttall oak to establish and grow in the understory of eastern cottonwood remains an uncertainty. In the current study, we examine biomass accumulation of Nuttall oak seedlings established under two contrasting afforestation techniques (interplanted beneath eastern cottonwood vs. planted in the open). The purpose of this investigation was to gain knowledge on the early

morphological development and biomass distribution of Nuttall oak seedlings that may be incorporated into future afforestation and forest restoration practices.

Methods

Study Site

The study was established in the LMAV on a former agricultural site located in Sharkey County, Mississippi, U.S.A. $(32^{\circ}58' \text{ N}, 90^{\circ}44' \text{ W})$. The site is situated in the humid, subtropical region of the temperate zone where the mean annual precipitation is 1,318 mm, and air temperature averages from 27.8 "C in July to 7.5°C in February (Scott & Carter 1962). Sharkey Clay, a very fine, smectitic, thermic, chromic Epiaquerts, is the predominant soil on the site. Core samples collected across the study site indicated that texture of this alluvial soil ranges from 7 to 12% sand, 22 to 42% silt, and 46 to 71% clay in the surface horizon (O-7.5 cm).

Establishment of Experimental Stands

More than 25 years of agricultural production ended on the study site with the harvest of a soybean (Clycine max [L.] Merrill) crop in the fall of 1994. We initiated this experiment in March 1995 by establishing a replicated eastern cottonwood plantation. Three 8.1-ha cottonwood stands were established according to methods used by Crown Vantage paper company as previously described by Gardiner et al. (2001). Cottonwood cuttings were hand-planted on a 3.7 \times 3.7-m spacing, and each stand replication received mechanical cultivation for competition control during the first two growing seasons. After the second growing season (February 1997), 1-0, bare-root Nuttall oak seedlings were interplanted between every other cottonwood row (3.7 x 7.3-m spacing). Additionally, a Nuttall oak stand was established immediately adjacent to each of the three interplanted cottonwood stands for the purpose of providing open-grown seedlings that would serve as experimental controls for morphology and growth measurements. Thus, the experiment included three replicated stands of Nuttall oak seedlings interplanted beneath eastern cottonwood and three replicated stands of opengrown Nuttall oak seedlings. The three Nuttall oak stands were hand-planted on a 3.7 \mathbf{x} 3.7-m spacing with the same stock material used for interplanting the cottonwood stands. Fifty experimental seedlings were selected in each stand and protected from small mammal herbivory with 0.6-m tall and 0.6-m diameter wire mesh shelters.

Experimental Stand Conditions

Sampling for this investigation began after oak seedlings completed three growing seasons. Table 1 provides a summary of the early stand characteristics that developed under each afforestation treatment. The eastern cottonwood nurse crop developed rapidly on this former

Tablel.	Stand	characteristics	of	open-grown	Nuttall	oak	and	eastern	cottonwood	interplanted	with	Nuttall	oak	established	on	former	agricultural
land in	Sharke	ey County, MS	5, 1	U.S.A.													

	Stand Type		
	Open-Grown Nuttall Oak	Cottonwood Interplanted with Oak	
Eastern cottonwood density (stems/ha) Eastern cottonwood basal area (m²/ha)		716 8.3 272 1,480	
Nuttall oak density (stems/ha)	405 4,230		
Herbaceous competitior (kg/ha) Light availability (μ mol m ⁻² s ⁻¹)*	1,047	32.5	

The open-grown Nuttall oak stands were 3 years old, eastern cottonwood stands WETE 5 years old, and interplanted Nuttall oaks were 3 years old.

*Light availability data arc the mean photosynthetic photon flux density from a cloud-free day (10 July 1998) in the middle of the second growing season for Nuttall oak seedlings

agricultural site providing a forest structure that appeared to reach maximum leaf area index 3 years after establishment. The cottonwood canopy reduced light availability to interplanted oak seedlings (approximately 30% of light available in the open), but the amount of herbaceous competition in the understory of cottonwood plantations was less than 35% of that in the open (Table 1). It should be noted that growth and productivity of eastern cottonwood is very site specific. Bottomland soils that are excessively flooded, for example, are not appropriate for establishment of eastern cottonwood.

Measurements and Statistical Procedures

After the third growing season for Nuttall oak, six randomly chosen oak seedlings were destructively sampled from each treatment combination (36 total). Height (cm) and root-collar diameter (mm) were measured, then each sample seedling was excavated from the soil and separated into leaf, stem, and root tissues. The number of leaves on each seedling was counted, then total leaf area was measured with an area meter (LI-COR, Lincoln, NE, U.S.A.). Average blade area was determined by dividing the total leaf area of the seedling by the number of leaves. Roots were washed to remove adhering soil particles then seedling biomass components were oven-dried at 50°C until completely desiccated. Oven-dried tissues were weighed, and proportional biomass accumulation by tissue type was computed according to the following formulae: leaf weight ratio (LWR) = leaf weight/total seedling weight; stem weight ratio (SWR) = stem weight/total seedling weight; and root weight ratio (RWR) = root weight/total seedling weight. Analyses of variance (ANOVA) according to a randomized block design (three levels of block, two levels of treatment) were conducted to test for a treatment effect on 12 response variables. Response variables tested included seedling height, rootcollar diameter, leaf weight, stem weight, root weight, total seedling weight, number of leaves, blade area, total leaf area, LWR, SWR, and RWR. Individual tests were conducted on each response variable at an alpha level of 0.05.

Results

Stem Growth

Three years after establishment, Nuttall oak height was not influenced by the presence of the eastern cottonwood canopy, as seedlings averaged approximately 126cm tall regardless of afforestation treatment (Table 2). Mean heights observed on sampled seedlings represent positive growth of more than two times the height of seedlings at planting. Seedling root-collar diameter also showed positive growth during the 3-year establishment period (Table 2), but the magnitude of this response was influenced by the afforestation treatment. Nuttall oak seedlings interplanted in the understory of cottonwood stands were 20% smaller in root-collar diameter compared to those established in the open (Table 2).

Biomass Accumulation

After 3 years of establishment on former agricultural land, accumulation of Nuttall oak seedling biomass was greatest in stems and roots (Fig. 1). However, the amount of biomass accumulated by seedlings differed according to the afforestation treatments. Seedlings established in the understory of eastern cottonwood accumulated less biomass than open-grown seedlings (Fig. 1). This reduction was observed for all plant components such that interplanted seedlings accumulated 54% less leaf mass (p = 0.0096), 59% less stem mass (p = 0.0242), and 50%

Table 2. Initial and third-year height and root-collar diameter ofNuttall oak seedlings planted on a former agricultural field in SharkeyCounty, MS, U.S.A.

Variable	Open-Grown	Interplanted	p-Value
Initial			
Height (cm)	50.4 ± 2.8	53.6 ± 2.6	0.4117
Root-collar diameter (mm)	8.1 ± 0.5	7.4 ± 0.5	0.3707
Year 3			
Height (cm)	128.4 ± 11.9	123.9 ± 14.1	0.7285
Root-collar diameter (mm)	26.7 ± 2.2	21.4 ± 1.3	0.0135

Values are expressed as mean \pm standard error.




Figure 1. Biomass accumulation of Nuttall oak seedlings established in the open and in the understory of an eastern cottonwood stand in Sharkey County, MS, U.S.A. The seedlings were established 3 years prior to excavation. Light bars represent open-grown seedlings, dark bars represent underplanted seedlings, and error bars are \pm standard error of the mean.

(p = 0.0113) less root mass relative to seedlings established in the open. Accordingly, Nuttalloak seedlings established in the open developed a total mass more than two times the mass of seedlings established beneath eastern cottonwood (p = 0.0148) (Fig. 1).

Biomass Distribution in Leaf Tissue

The comparatively greater amount of leaf biomass observed on open-grown Nuttall oak seedlings is partially attributed to a larger number of leaves (250% more) relative to interplanted seedlings (Table 3). Leaf blades of open-grown seedlings, however, were generally smaller in area (52%) than those of the interplanted treatment (Table3). Despite the greater number of leaves observed on open-grown seedlings, we were unable to detect a treatment effect on total leaf area (Table3). Thus, the photosynthetic surface area of interplanted seedlings did not differ from that maintained by open-grown seedlings.

Biomass Distribution in Seedlings

Three years after establishment, Nuttall oak seedlings accumulated the smallest proportion of fixed carbon in leaf tissue (10%) (Fig. 2). Stem tissue comprised 42%

Table 3. The number of leaves, mean blade area, and mean total leaf area observed on 3-year-old Nuttall oak seedlings planted on a former agricultural field in Sharkey County, MS. U.S.A.

Variable	Open-Grown	Interplanted	p-Value
Number of leaves Blade area (cm') Leaf area (cm')	$\begin{array}{c} 345 \pm 83 \\ 10.4 \ \pm 0.88 \\ 3,259 \pm 734 \end{array}$	$\begin{array}{c} 98 \pm 16 \\ 21.5 \pm 1.98 \\ 2,157 \pm 402 \end{array}$	$\begin{array}{c} 0.0023 \\ < 0.0001 \\ 0.0953 \end{array}$

Values are expressed as mean \pm standard error.




Figure 2. Proportional biomass accumulation of Nuttall oak seedlings established in the open and in the understory of an eastern cottonwood stand in Sharkey County, MS, U.S.A. The seedlings were established 3 years prior to excavation. LWR \equiv leaf weight ratio. SWR = stem weight ratio, RWR \equiv root weight ratio. Light bars represent open-grown seedlings, dark bars represent underplanted seedlings, and error bars are \pm standard error of the mean.

whereas root tissue comprised about 48% of total seedling biomass (Fig.2). Irrespective of the large differences observed for biomass accumulation, the proportional distribution of biomass within seedlings was not altered by stand environment (Fig.2). Interplanted Nuttall oak seedlings accumulated leaf (p = 0.9736), stem (p = 0.0847), and root (p -0.1042) biomass in equal proportions to open-grown seedlings.

Discussion

Seedlings planted on former agricultural fields in the LMAV are prone to encounter numerous stress agents including vigorous herbaceous weed competition, mammal and insect herbivory, and drought and flooding (Allen et al. 2001). Small seedlings of poor vigor are generally more vulnerable to some of these environmental stresses than are vigorous large seedlings and saplings (Allen et al. 2001). For example, large seedlings and saplings have an advantage over smaller seedlings in the event of shallow flooding because the crown of the taller seedling is at a lower inundation risk. Thus, it is critical for establishment success that seedlings planted on former agricultural fields in the LMAV exhibit vigorous early growth. Irrespective of establishment treatment, Nuttall oak seedlings examined in this study averaged nearly 25 cm of annual height growth during the 3-year establishment period. This growth rate was within the expected range for bare-root, bottomland oak seedlings established on former agricultural fields without the use of competition control (Kennedy 1981, 1993). The presence of the eastern cottonwood canopy did not reduce average seedling height 3 years after establishment. In contrast, root-collar diameter was larger on seedlings established and grown in the open.

Though root-collar growth was not maximized when seedlings were established in the understory of eastern cottonwood, the root-collar growth we observed on interplanted seedlings appeared as good or better than other published observations for open-grown Nuttall oak planted on similar soils (Kennedy 1981; Patterson & Adams 2003). However, it is not known whether these interplanted seedlings with relatively smaller stem diameters will be prone to wind damage when released.

Our research may be the first to examine oak seedling growth and development in the understory of another broadleaf species established on former agricultural land. Studies examining oak seedling height and diameter growth in relation to establishment beneath partial overstories of natural broadleaf species stands have demonstrated mixed results. Teclaw and Isebrands (1993), who worked in natural stands in Wisconsin, demonstrated that northern red oak (Quercus rubra L.) seedlings may exhibit better height growth under a partial canopy than in the open, particularly where open sites are subject to vigorous competition and late spring frosts. Truax et al. (2000), who compared development of northern red oak and bur Michx.) growing in open fields oak and beneath natural aspen (*Populus tremuloides* Michx.) stands in Canada, observed contrasting results for these two species. They observed a better height and diameter growth increment for northern red oak established beneath aspen, whereas bur oak exhibited a better height and diameter growth increment when established in the open (Truax et al. 2000). Results from our study were most closely in line with those presented by Gcmmcl et al. (1996), who studied the development of pedunculate oak (Quercus robur L.) under different canopy densities of natural stands in southern Sweden. In their study, seedlings established beneath the partial canopies grew to the same height as those established in the open, but rootcollar diameter was less than that of open-grown seedlings. It is obvious that the range of responses documented in the literature precludes generalizations on how oak seedling height and diameter will respond to an understory environment as responses appear specific to species, forest environment, and silvicultural system.

Biomass accumulation by Nuttall oak, particularly whole plant biomass for seedlings growing on former agricultural land in the LMAV, has not been well documented (Schlaegel & Willson 1983). In the current study, Nuttall oak seedlings established under two different afforestation treatments showed positive biomass accumulation over the 3-year study period, but biomass accumulation rate was determined by the environment in which seedlings were established. Our findings indicate that Nuttall oak seedlings established in the understory of eastern cottonwood plantations will maintain a positive accumulation of leaf, stem, and root biomass, but the accumulation rate will be less than for seedlings established in **the** open. This observation is in line with the decrease in root-collar diameter we observed on interplanted seedlings. This finding conflicts with observations on oak seedling biomass accumulation beneath artificial shade cloth. Biomass accumulation by cherrybark oak (Quercus pagoda Raf.) and pedunculate oak seedlings established under moderate levels of light availability can be as high or greater than for seedlings established under full sunlight (Ziegenhagen & Kausch 1995; Gardiner & Hodges 1998), because the artificial shade is thought to reduce seedling water stress. However, our finding is consistent with that of Gemmel et al. (1996), who reported greater biomass accumulation by pedunculate oak seedlings established in the open versus beneath partial canopies in southern Sweden. Moderating light availability with the use of partial overstories in natural stands or sparse canopies of plantation species does not appear to benefit oak seedling biomass accumulation to the same extent observed when seedlings are cultured under artificial shade.

Nuttall oak seedlings established in the understory of eastern cottonwood stands produced fewer leaves than those seedlings grown in the open, but they produced leaves with a blade area twice the size of those on opengrown seedlings. Because of these characteristics, seedlings from both afforestation treatments showed similar leaf areas. Oak seedlings studied by other workers exhibited a wide range in leaf number, blade size, and total leaf area responses along light gradients. Pedunculate oak, black oak (Quercus velutina Lam.), and banj oak (Quercus leucotrichophora A. Camus) showed a higher number of leaves in high light environments than in low light environments (Gottschalk 1994; Thadani & Ashton 1995; Welander & Ottosson 1998). Some oak species increase blade size under low irradiance, whereas others do not exhibit this characteristic (Callaway 1992). Valley oak (Quercus lohata Nee) and blue oak (Quercus douglasii H. & A.) did not alter total leaf area when grown under a range of light availability, but coast live oak (Quercus agrifolia Nee) seedlings did respond to increased light availability by decreasing blade area, which led to a decrease in total leaf area (Callaway 1992). Most species of oak will probably decrease leaf area under low light availability, but this response is typically not observed until extremely low levels of light are experienced (Gottschalk 1994; Ziegenhagen & Kausch 1995). The light levels observed in this study (approximately 30% of light available in the open) along with the ability of seedlings to maintain appreciable leaf area in the understory of eastern cottonwood are consistent with our premise that the understory environment in eastern cottonwood plantations appears suitable for Nuttall oak seedling establishment and growth.

Though establishment beneath the cottonwood canopy reduced total biomass accumulation by Nuttall oak seedlings, it is interesting that the relative distribution of accumulated biomass in seedlings was similar for both treatments. For the open and the understory environment, leaf tissue comprised 10%, stem tissue comprised 42%, and root tissue comprised 48% of total seedling biomass. These results illustrate that Nuttall oak seedlings accumulated near equal proportions of biomass above- and belowground 3 years after establishment on a former agricultural field.

Several studies conducted on various oak species indicate that oak seedlings acclimate morphologically to environmental stresses through proportional biomass accumulation rates (Kolb et al. 1990; Thadani & Ashton 1995; Ashton & Larson 1996; Gardiner & Hodges 1998). For example. Kolb and Steiner (1990) demonstrated that northern red oak seedlings acclimated to belowground competition by increasing the proportion of biomass accumulated in roots. Welander and Ottosson (1998) reported a decrease in the root-to-shoot ratio of pedunculate oak seedlings with decreasing light availability. This response favors biomass accumulation in aboveground components to improve light gathering in low light environments (Welander & Ottosson 1998). The fact that seedlings established beneath eastern cottonwood accumulated less biomass than those established in the open indicates that light, soil moisture, or another factor were not sufficiently available in the understory to maximize Nuttall oak seedling growth. Though growth was not maximized, proportional accumulation of biomass indicates that resource availability in the eastern cottonwood understory did not limit seedling function substantially enough to alter Nuttall oak seedling morphology.

Stanturf et al. (2001) argued that afforestation is a necessary first step toward ecological restoration of bottomland hardwood ecosystems on former agricultural land. Conventional afforestation practices have generally targeted the establishment of mast-producing overstory species such as bottomland oaks (Allen 1997: Schoenholtz et al. 2001). Bottomland oaks have been favored because of their high value as a component of wildlife habitat and their limited dispersal; it is also thought that light-seeded species will establish naturally (Allen 1997; King & Keeland 1999). A primary criticism of conventional afforestation practices employed on bottomland sites centers on their effectiveness at initiating restoration of ecological functions of bottomland hardwood ecosystems. For example, Allen (1997), who studied woody plant invasion on 10 afforestation sites in the LMAV, expressed concern that conventional afforestation practices were leading to stands with low woody species diversity relative to natural bottomland hardwood stands. He observed a limited amount of natural invasion and inferior growth by light-seeded species in oak plantations, especially ii stands were not immediately adjacent to a seed source (Allen 1997). Likewise, Ouchley et al. (2000) examined the composition of a historic bottomland hardwood forest, concluding that current silvicultural and afforestation practices foster a higher component of oak species than what was characteristic of historic bottomland hardwood forests. Allen (1997) and Ouchley et al. (2000) provide arguments that support establishment of more complex plantations with a higher component of light-seeded species to facilitate restoration of tree species diversity

on bottomland hardwood afforestation sites. Their arguments and observations by others illustrate a clear need to identify and implement afforestation practices that will provide pathways toward catalyzing development of biodiversity components in bottomland ecosystems.

The eastern cottonwood-Nuttall oak interplanting system examined in this manuscript is a unique alternative for establishing bottomland hardwood forests because it quickly develops a forest cover, which may facilitate restoration of some ecological functions more rapidly than conventional afforestation. For example, results from this study indicate the understory microenvironment of eastern cottonwood stands is sufficient for establishment of the shade-intolerant Nuttall oak; hence it is likely that other bottomland hardwood species can invade the understory. Indeed, Hodges (1997) reported that shadeintolerant species such as American sycamore (Platanus occidentalis L.), sweet pecan (Curvu illinoensis [Wang.] K. Koch), green ash (Fraxinus pennsylvanica Marshall), and sweetgum (Liquidambar styraciflua L.) are often present in the understory of natural eastern cottonwood stands. Our own observations indicate that given a seed source, numerous bottomland hardwood species readily become established in the understory of eastern cottonwood plantations. Thus, use of a plantation system such as this may serve to catalyze invasion by other bottomland hardwood species, thereby increasing species richness on the site as is the case for similar plantation systems studied in the tropics (Parrotta 1992; Lugo 1997; Powers et al. 1997).

Hamel (2003), who studied winter bird use on afforestation sites, described another example of how the eastern cottonwood-Nuttall oak interplanting system can provide a swift trajectory toward developing attributes of a natural ecosystem. He observed a more rapid assemblage of forest canopy-dwelling birds on sites receiving the eastern cottonwood–Nuttall oak interpianting system relative to sites receiving afforestation with oak alone (Hamel 2003). Hamel (2003) attributed this rapid assemblage of forest canopy birds to the quick development of forest cover by eastern cottonwood. Similarly, Twedt and Portwood (1997) found a it-fold increase in the number of bird species that established breeding territories in 5- to 7-year-old eastern cottonwood stands versus 4- to 6-yearold oak plantings. As our research efforts continue to focus on this eastern cottonwood-Nuttall oak afforestation system, we are encouraged that additional benefits to restoration of bottomland hardwood ecosystem functions will be revealed. Future work with this afforestation system will concentrate on the developmental trajectory of various biodiversity components in bottomland hardwood ecosystems, including understory flora, soil arthropods, small mammals, and insect assemblages.

Management implications

Several management implications of practical significance to forest restoration practitioners working in the LMAV

and other temperate regions may be drawn from this work. First, we documented substantial height growth and positive biomass accumulation for the 3-year period Nuttall oak seedlings were established in the understory of eastern cottonwood stands. We are encouraged that these interplanted seedlings are rapidly developing into large saplings that should respond quickly to release and are at a lower risk of sustaining damage from herbivory or flooding. We also noted similar morphologies between seedlings established in the open and beneath eastern cottonwood, indicating that resources, that is, light, water, and nutrients, required for growth by interplanted seedlings were sufficiently available to maintain seedling function. These findings complement earlier work on leaflevel photosynthetic characteristics of Nuttall oak seedlings established beneath eastern cottonwood (Gardiner et al. 2001) and support the application of this interplanting system as an alternative method of establishing Nuttall oak as a component in restored bottomland hardwood stands on former agricultural land in the LMAV.

Second, we present favorable results from an experimental afforestation system that begins with rapid development of forest cover on former agricultural land through the establishment of a fast-growing, native plantation species (eastern cottonwood). After the establishment of forest cover, the plantation understory is enriched with a slower-growing, disturbance-dependent species (Nuttall oak) that is generally considered mid-successional. The promising application of this interplanting system in the LMAV provides a forest restoration model that can be tried with other species in other temperate regions. Advantages of this afforestation model include, but arc not limited to, a speedy transition from cleared land to forest structure and associated forest functions (Hamel 2003), the potential to realize an early financial return on the restoration investment (Stanturf & Portwood 1999), and the opportunity to establish complex forest structure through enrichment of the understory with slowergrowing, desirable species.

Finally, the magnitude of the afforestation effort currently concentrated in the LMAV presents a considerable opportunity for forest restoration, particularly for bottomland hardwood ecosystems. Because of biological and social factors inherent to the LMAV, for example, site heterogeneity, tree species diversity, ownership characteristics, landowner objectives, and restoration project scale, there is a strong need to provide alternative mechanisms for achieving restoration of sustainable forest ecosystems. We demonstrate that research aimed at understanding basic plant processes can aid in the development of biologically feasible afforestation systems that will provide the foundation for advancing sustainable forest restoration practices.

Acknowledgments

We thank several people and organizations that collaborated on the successful establishment and maintenance of this experiment. Primary contributors to this research included Jim Shepard with the National Council of the Paper Industry for Air and Stream Improvement, Inc., Tim Wilkins, Lamar Dorris, and Ronnie Haynes with the U.S. Fish and Wildlife Service, and Jeff Portwood (formerly with Crown Vantage). Mathew Stroupe, Ralf Zellin, and Ron Fisher assisted with seedling excavations. Several others, including Pat Weber and Jackie Henne-Kerr, have been a source of encouragement and constructive dialogue regarding forest restoration in the LMAV. We also thank our colleagues who provided helpful reviews of earlier drafts of this manuscript.

LITERATURE CITED

- Allen, J. A. 1990. Establishment of bottomland oak plantations on the Yazoo National Wildlife Refuge Complex. Southern Journal of Applied Forestry 14:206–210.
- Allen, J. A. 1997. Reforestation of bottomland hardwoods and the issue of woody species diversity. Restoration Ecology 5:125–134.
- Allen, J. A., B. D. Keeland, J. A. Stanturf, A. F. Clewell, and H. E. Kennedy, Jr. 2001. A guide to bottomland hardwood restoration. USGS, Biological Resources Division Information and Technology Report USGS/BRD/ITR-2000-0011; USDA, Forest Service, Southern Research Station, General Technical Report SRS-40. U.S. Geological Survey, Reston, Virginia.
- Ashton, P. M. S., S. Gamage, I. A. U.N. Gunatilleke, and C. V. S. Gunatilleke. 1997. Restoration of a Sri Lankan rainforest: using Caribbean pine *Pinus caribaea* as a nurse for establishing late-successional tree species. Journal of Applied Ecology 34:915–925.
- Ashton, P. M. S., S. Gamage, I. A. U. N. Gunatilleke, and C. V. S. Gunatilleke. 1998. Using Caribbean pine to establish a mixed plantalion: testing effects of pine canopy removal on plantings of rain forest tree species. Forest Ecology and Management 106:21 1-222.
- Ashton, M. S., and B. C. Larson. 1996. Germination and seedling growth of *Quercus* (section *Erythrobalanus*) across openings in a mixeddeciduous forest of southern New England. USA. Forest Ecology and Management 80:81–94.
- Callaway, R. M. 1992. Morphological and physiological responses of three California oak species to shade. International Journal of Plant Science **153**:434–441.
- FAG (Food and Agricultural Organization of the United Nations). 2001. Global forest resources assessment 2000—main report. Food and Agriculture Organization of the United Nations, FAO Forestry Paper 140. FAG, Rome, Italy.
- Gardiner, E. S., and J. D. Hodges. 1998. Growth and biomass distribution of cherrybark oak (*Quercus pagoda* Raf.) seedlings as influenced by light availability. Forest Ecology and Management 108:127–134.
- Gardiner, E. S., and J. M. Oliver. Restoration of bottomland hardwood forests in the Lower Mississippi Alluvial Valley, USA. In J. A. Stanturf, and P. Madsen, editors. Restoration of boreal and temperate forests. CRC Press, Boca Raton, Florida. (in press).
- Gardiner, E. S., D. R. Russell. M. Oliver, and L. C. Dorris, Jr. 2002. Bottomland hardwood afforestation: state of the art. Pages 75–86 in M. M. Holland, M. L. Warren. and J. A. Stanturf. editors. Proceedings of a conference on sustainability of wetlands and water resources; how well can riverine wetlands continue to support society into the 21st century? USDA Forest Service, Southern Research Station, General Technical Report SRS-SO. IJSDA Forest Service, Asheville, North Carolina.
- Gardincr, E. S., C. J. Schweitzer, and J. A. Stanturf. 2001. Photosynthesis of Nuttall oak (*Quercus nuttallii* Palm.) seedlings interplanted

beneath an eastern cottonwood (*Populus deltoides* Bartr. ex Marsh.) nurse crop. Forest Ecology and Management **149**:283–294.

- Gemmel, P., U. Nilsson, and T. Welander. 1996. Development of oak and beech seedlings planted under varying shelterwood densities and with different site preparation methods in southern Sweden. New Forests 12:141–161.
- Gottschalk, K. W. 1994. Shade, leaf growth and crown development of *Quercus rubra*, *Quercus velutina*, *Prunus serotina* and *Acer rubrum* seedlings. Tree Physiology 14:735–749.
- Hamel, P. B. 2003. Winter bird community differences among methods of bottomland hardwood forest restoration: results after seven growing seasons. Forestry 76:189–197.
- Hodges, J. D. 1997. Development and ecology of bottomland hardwood sites. Forest Ecology and Management 90:117–125.
- Keenan, R., D. Lamb, O. Woldring, T. Irvine, and R. Jensen. 1997. Restoration of plant biodiversity beneath tropical tree plantations in Northern Australia. Forest Ecology and Management 99: 117–131.
- Kennedy, H. E., Jr. 1981. Foliar nutrient concentrations and hardwood growth influenced by cultural treatments. Plant and Soil 63:307–316.
- Kcnncdy, H. E., Jr. 1993. Artificial regeneration of bottomland oaks. Pages 241-249 in D. L. Loftis, and C. E. McGee, editors. Oak regeneration: serious problems, practical recommendations. USDA Forest Service, Southeastern Forest Experiment Station, General Technical Report SE-X4. USDA Forest Service, Asheville, North Carolina.
- King, S. L., and B. D. Keeland. 1999. Evaluation of reforestation in the Lower Mississippi River Alluvial Valley. Restoration Ecology 7: 34X-359.
- Kolb, T. E., and K. C. Steiner. 1990. Growth and Biomass partitioning of northern red oak and yellow-poplar seedlings: effects of shading and grass root competition. Forest Science 36:34–44.
- Kolb, T. E., K. C. Steiner, L. H. McCormick, and T. W. Bowersox. 1990. Growth response of northern red-oak and yellow-poplar seedlings to light. soil moisture and nutrients in relation to ecological strategy. Forest Ecology and Management 38:65-7X.
- Lugo, A. E. 1997. The apparent paradox of reestablishing species richness on degraded lands with tree monocultures. Forest Ecology and Management 99:9–19.
- Newling, C. J. 1990. Restoration of the bottomland hardwood forest in the Lower Mississippi Valley. Restoration and Management Notes 8: 23-2X.
- Ouchley, K., R. B. Hamilton, W. C. Barrow, Jr., and K. Ouchley. 2000. Historic and present-day forest conditions: implications for bottomland hardwood forest restoration. Ecological Restoration 18:21–25.
- Parrotta, J. A. 1992. The role of plantation forests in rehabilitating degraded tropical ecosystems. Agriculture, Ecosystems and Environment 41:115–133.
- Parrotta, J. A. 1995. Influence of overstory composition on understory colonization by native species in plantations on a degraded tropical site. Journal of Vegetation Science 6:627–636.

- Parrotta, J. A., J. W. Turnbull, and N. Jones. 1997. Catalyzing native forest regeneration on degraded tropical lands. Forest Ecology and Management 99:1–7.
- Patterson. W. B., and J. C. Adams. 2003. Soil, hydroperiod and bedding effects on restoring bottomland hardwoods on flood-prone agricultural lands in North Louisiana, USA. Forestry 76:181–188.
- Powers, J. S., J. P. Haggar, and R. F. Fisher. 1997. The effect of overstory composition on understory woody regeneration and species richness in 7-year-old plantations in Costa Rica. Forest Ecology and Management 99:43–54.
- Schlaegel, B. E., and R. B. Willson. 1983. Nuttall oak volume and weight tables. USDA, Forest Service, Research Paper SO-1X6. USDA Forest Service, New Orleans, Louisiana.
- Schoenholtz, S. H., J. P. James, R. M. Kaminski, B. D. Leopold. and A. W. Ezell. 2001. Afforestation of bottomland hardwoods in the Lower Mississippi Alluvial Valley: status and trends. Wetlands 21:602-613.
- Scott, F. T., and R. C. Carter. 1962. Soil survey of Sharkey County. Mississippi. United States Department of Agriculture. Soil Conservation Service, Washington, D.C.
- Stanturf, J. A., E. S. Gardiner, P. B. Harnel, M. S. Devall, T. D. Leininger, and M. E. Warren, Jr. 2000. Restoring bottomland hardwood ecosystems in the Lower Mississippi Alluvial Valley. Journal of Forestry 98:10–16.
- Stanturf, J. A., and C. J. Portwood. 1999. Economics of afforestation with eastern cottonwood (*Populus deltoides*) on agricultural land in the Lower Mississippi Alluvial Valley. Pages 66-72 in J. D. Haywood, editor. Proceedings of the Tenth Biennial Southern Silvicultural Research Conference. USDA, Forest Service, General Technical Report SRS-30. USDA Forest Service, Asheville, North Carolina.
- Stanturf, J. A., S. H. Schoenholtz, C. J. Schweitzer, and J. P. Shepard. 2001. Achieving restoration success: myths in bottomland hardwood forests. Restoration Ecology 9:189–200.
- Stanturf, J. A., C. J. Schweitzer, and E. S. Gardiner. 1998. Afforestation of marginal agricultural land in the Lower Mississippi River Alluvial Valley, U.S.A. Silva Fennica 32:281–297.
- Teclaw, R. M., and J. G. Isebrands. 1993. An artificial regeneration system for establishing northern red oak on dry-mesic sites in the Lake States, USA. Annals des Sciences Forestières 50:543–552.
- Thadani, R., and P. M. S. Ashton. 1995. Regeneration of banj oak (Quercus leucotrichophora A. Camus) in the central Himalaya. Forest Ecology and Management 78:217–224.
- Truax, B., F. Lambert, and D. Gagnon. 2000. Herbicide-free plantations of oaks and ashes along a gradient of open to forested mesic environments. Forest Ecology and Management 137:155–169.
- Twedt. D. J., and J. Portwood. 1997. Bottomland hardwood reforestation for neotropical migratory birds: are we missing the forest for the trees? Wildlife Society Bulletin **25:**647–652.
- Welander, N. T., and B. Ottosson. 1998. The influence of shading on growth and morphology in seedlings of *Quercus robur* L. and *Fagus sylvatica* L. Forest Ecology and Management 107:117–126.
- Ziegenhagen, B., and W. Kausch. 1995. Productivity of young shaded oaks (*Quercus robur* L. <u>as</u> corresponding to shoot morphology and leaf anatomy. Forest Ecology and Management **72:**97–108.

Great Trinity Forest Management Plan

HARDWOOD SILVICULTURE

Early Response of Interplanted Nuttall Oak to Release from an Eastern Cottonwood Overstory

(Connor, Kristina F., ed. 2006. Proceedings of the 13th Biennial southern silvicultural research conference. Gen. Tech. Rep. SRS – 92. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 640 p.)

EARLY RESPONSE OF INTERPLANTED NUTTALL OAK TO RELEASE FROM AN EASTERN COTTONWOOD OVERSTORY

Emile S. Gardiner¹

Abstract—Forest restoration activity in the Lower Mississippi Alluvial Valley has generated a demand for alternative afforestation practices that can accommodate diverse landowner objectives. An alternative afforestation practice now being studied involves rapid establishment of a forest canopy of eastern cottonwood (*Populus deltoides* Bartr. ex Marsh.), followed by interplanting with seedlings of slower-growing bottomland hardwood species. An experiment conducted in Mississippi shows that Nuttall oak (*Quercus nuttallii* Palm.) seedlings can develop into vigorous sapling-sized reproduction when interplanted in the understory of eastern cottonwood, and that the cottonwood overstory can be harvested without adversely affecting the interplanted reproduction. Harvesting damage was minimal, and excellent survival, vigorous sprouting, and adequate growth of Nuttall oak reproduction interplanted beneath eastern cottonwood indicate that Nuttall oak reproduction can be established under and released from an eastern cottonwood overstory.

INTRODUCTION

Over the past 15 years, a substantial area of cropland in the Lower Mississippi Alluvial Valley (LMAV) has been converted to hardwood forest plantations (Gardiner and Oliver 2005). Due to the extensiveness of afforestation in this region, plantation establishment practices and costs are often minimized, and this conventional approach frequently contributes to poor stocking or slow growth of stands. It has been found that conventional afforestation practices may not be the best means of accomplishing certain landowner objectives (Stanturf and others 2000, Stanturf and others 2001). Recent interest in establishing plantations for objectives such as biodiversity, carbon sequestration, economic sustainability, and water quality has led managers to seek alternative afforestation practices that promote development of complex plantations (Gardiner and others 2002).

In one alternative afforestation system now being tested in the LMAV, rows of bottomland hardwood species of later successional seres, such as Nuttall oak (Quercus nuttallii Palm.), are interplanted between previously established rows of eastern cottonwood (Populus deltoides Bartr. ex Marsh.) (Gardiner and others 2001, Gardiner and others 2004). In this system, the cottonwood stand serves as a nurse for the other hardwood reproduction developing in the understory. At the end of the 10- to 15-year cottonwood rotation, the cottonwood is harvested and the hardwood reproduction that has been established in the understory is released. This alternative has several ecological and economic advantages over conventional afforestation practices in the LMAV. These advantages include more rapid carbon sequestration (Stanturf and others 2003), quicker development of forest bird assemblages (Hamel 2003), earlier economic return on the afforestation investment (Stanturf and Portwood 1999), and favorable stand conditions for regeneration of other bottomland tree and shrub species (Gardiner and others 2004). Thus, this interplanting system can provide for multiple objectives while catalyzing bottomland hardwood ecosystem processes.

Though substantial benefits of this interplanting system have been documented, it has not been demonstrated that the

cottonwood biomass can be harvested without significant damage to the developing reproduction in the understory. Also, it has not been shown that the interplanted Nuttall oak reproduction will respond satisfactorily once the eastern cottonwood overstory is harvested. In this study, a mature plantation of eastern cottonwood interplanted with Nuttall oak was harvested experimentally to address these questions. More specifically, the objectives of this research were to (a) quantify the extent of harvesting damage to interplanted Nuttall oak reproduction when the eastern cottonwood overwood was removed; and (b) quantify sprouting frequency, survival, and growth of interplanted Nuttall oak reproduction following release from the eastern cottonwood overstory.

METHODS

The study was conducted in a 10-year-old eastern cottonwood plantation adjacent to Steele Bayou on the Fitler Plantation, Issaquena County, MS. Soil on the site is mapped to the Commerce series (fine-silty, mixed, superactive, nonacid, thermic Fluvaquentic Endoaquepts) (Wynn and others 1959), and this series is generally well suited for culture of eastern cottonwood. Operational practices described in Gardiner and others (2001) were employed to establish the cottonwood stand on a conventional spacing of 12 by 12 feet in the winter of 1991-92. In the winter of 1993-94, rows of 1-0 bareroot Nuttall oak seedlings were planted between rows of cottonwoods. The oak rows were 24 feet apart, and the oaks were 12 feet apart within rows. Oak seedlings were hand planted and did not receive competition control or other cultural treatments after planting.

Six plots were delineated. Each measured approximately 180 by 240 feet and encompassed 15 rows of eastern cottonwood and 7 rows of Nuttall oak, each row being 240 feet long. We inventoried all Nuttall oak reproduction in each plot, assigning a unique number to each stem. Forty Nuttall oak stems in each plot were randomly selected to serve as sample plants (240 total plants). We measured the height (feet), root-collar diameter (inches), and diameter at breast height (inches) of each sample plant.

¹ Research Forester, USDA Forest Service, Southern Research Station, Center for Bottomland Hardwoods Research, Stoneville, MS 38776.

Citation for proceedings: Connor, Kristina F., ed. 2006. Proceedings of the 13th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS–92. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 640 p.

In November 2001, the plots were harvested with a John Deere 843H feller buncher. There were two harvesting treatments, and these were assigned to plots randomly. Three plots received each harvesting treatment. In one treatment, all cottonwood stems were felled and removed from the site as whole trees with a grapple skidder. In the other treatment, all cottonwood stems were felled, all Nuttall oak stems were severed by the feller buncher, and a grapple skidder was used to skid all harvested cottonwood from the site. The first treatment was designed both to show how much damage a typical harvesting operation might do to Nuttall oak reproduction in a stand of this kind and to determine the growth response to release by the interplanted Nuttall oak reproduction. The second treatment was designed to provide insight into the amount of damage to, and stool viability and growth response of, interplanted Nuttall oak reproduction that was coppiced.

Immediately after the harvest, we assessed physical damage sustained by sample plants. We noted if reproduction had bark removed, a broken shoot, or was uprooted during harvesting. Following three consecutive growing seasons after the harvest, we measured stem height and the number of sprouts on all surviving Nuttall oak sample plants. Analysis of variance for a completely random design was conducted on plot means of stem damage and stem height for each growing season. Tests were conducted at an $\alpha = 0.05$ level.

RESULTS AND DISCUSSION

Logging Damage to Reproduction

Logging damage to Nuttall oak reproduction averaged 6 percent across the study site. Most of the observed logging damage was recorded as leaning or bent stems, and this damage appeared to be caused as cottonwood trees were felled rather than by maneuvering of logging equipment. No uprooting of plants was observed. Previous experience of harvesting operations in other cottonwood stands suggested that the felling and skidding equipment would require about 16 feet to maneuver between trees. The 24-foot spacing between Nuttall oak rows in this study made it possible to maneuver harvesting machinery without significantly damaging reproduction interplanted in the understory.

Damage to and Sprouting of Nuttall Oak Stumps

In plots that were harvested completely, Nuttall oak stems were clipped about 10 inches above groundline by the feller buncher. Typical logging damage was not observed in these plots because shoots that might have been injured by cottonwood felling were cut during the harvest. However, cutting of the small-diameter Nuttall oak stems with the feller head caused splintering of about 64 percent of the Nuttall oak stumps. Also, nearly 12.5 percent of the stems were not completely severed from the stump but were partially severed and broken over by the feller buncher. The Nuttall oak reproduction remained viable and sprouted readily. After the first growing season, there was an average of 4.3 ± 0.7 sprouts per stump. Mortality within clumps progressed slowly, and stumps supported 3.8 ± 0.4 sprouts 3 years after harvest. At that time, 99 percent of all stumps had live sprouts.

Natural or anthropogenic disturbances in bottomland hardwood stands often kill or remove the shoot of advance oak reproduction. Vigorous oak saplings generally coppice readily following disturbance if the root system and root collar remain intact (Hodges and Gardiner 1993). Because established oak saplings sprout readily, they typically maintain a high value in regeneration of natural bottomland stands (Belli and others 1999). Our findings are consistent with literature on oak sprouting in natural stands in that Nuttall oak artificially established in the understory of eastern cottonwood coppiced readily and survived well following overstory harvest. Our findings on Nuttall oak coppicing and the lack of logging damage to stumps when only the cottonwood overstory was removed may support interplanting of reproduction in a more dense configuration if desired, e.g., a 12 by 12 foot spacing.

Height Development of Reproduction

Prior to the harvest, Nuttall oak reproduction averaged 11.6 \pm 1.1 feet in height, and mean height did not differ by harvest treatment (*P* = 0.5304) (fig. 1). Saplings released by the cotton-wood harvest had not shown an above-ground response by the end of the first growing season when they had an average height of 11.7 \pm 1.3 feet. Nuttall oak stump sprouts responded vigorously, growing to almost half of their preharvest height (5.7 \pm 0.4 feet). By the end of the third growing season, reproduction that was not coppiced maintained a 25 percent height advantage over coppiced plants (*P* = 0.036). However, coppice was growing vigorously, averaged 13.0 \pm 0.7-feet tall, and had surpassed its preharvest height (fig. 1).

Height distributions of the released Nuttall oak illustrate the development of each reproduction type into taller height classes (fig. 2). Fifty-five percent of saplings that were not coppiced were in a height class taller than 10 feet prior to harvest. Three years following release, 89 percent had advanced into height classes taller than 10 feet. Sixty-five percent of saplings that were coppiced were in a height class taller than 10 feet prior to harvest, and 87 percent had grown into these taller height classes 3 years after treatment (fig. 2).

Slow above-ground response to release has often been cited as a factor that makes bottomland oak reproduction less competitive than reproduction of more intolerant species following stand disturbance (Hodges and Gardiner 1993, Lockhart and others 2000). Several physiological and morphological factors




Figure 1—Mean height of coppiced and intact Nuttall oak reproduction interplanted beneath eastern cottonwood. This reproduction was released from the cottonwood overstory following year zero.




Figure 2—Height distribution of intact (below) and coppiced (above) Nuttall oak reproduction that was interplanted beneath eastern cottonwood. This reproduction was released from the cottonwood overstory following year zero.

characteristic of oak reproduction may contribute to this sluggish response (Hodges and Gardiner 1993), but some authors have suggested the use of shoot clipping or coppicing to rejuvenate above-ground growth of oak reproduction, particularly where this reproduction exhibits poor apical vigor (Deen and others 1993, Gardiner 1998, Janzen and Hodges 1987, Lockhart and others 2000). In this study, established Nuttall oak saplings responded slowly to release from the eastern cottonwood overstory, showing no appreciable height increment the first growing season after harvest. These saplings, however, eventually acclimated to the disturbance and made substantial height growth by the third growing season after harvest. Height growth of Nuttall oak stump sprouts responded rapidly to release, but this growth had not compensated for the reduction in stem length 3 years after clipping. In contrast, Stanturf and Kennedy (1996) demonstrated that the 5-year height of top-pruned cherrybark oak (Q. pagoda Raf.) seedlings did not differ from the average height of seedlings that

were not top-pruned. Nevertheless, oak reproduction established in the understory of eastern cottonwood will respond to removal of the canopy on afforestation sites and should attain upper canopy positions in the subsequent stand.

SUMMARY

A valid concern when designing and establishing interplanted plantations for forest restoration is providing for harvest of the nurse crop and effective release of the understory reproduction. We found that an eastern cottonwood overstory can be harvested with minimal damage to interplanted Nuttall oak reproduction if spacing between and within rows is as it was in this experiment. Mortality of Nuttall oak reproduction was minimal through three growing seasons whether stems remained intact or were severed by felling equipment. Additionally, splintering of the Nuttall oak stumps observed in this study did not reduce coppicing. Our results may support interplanting of reproduction in a more dense configuration if desired, e.g., a 12 by 12 foot spacing. Height development of released Nuttall oak saplings was slow initially, but these saplings eventually acclimated to the open environment and were growing well by year three. Nuttall oak coppice developed rapidly over three growing seasons but still lagged behind saplings that were not coppiced. The lack of harvesting damage and the excellent survival, vigorous sprouting, and adequate growth of Nuttall oak reproduction interplanted beneath eastern cottonwood indicate that reproduction of this species can be interplanted and established under an eastern cottonwood overstory and then released successfully. These results support establishment of more complex plantations to address multiple management objectives when restoring forests on former agricultural land.

ACKNOWLEDGMENTS

Jackie Henne-Kerr provided invaluable assistance in designing and establishing this experiment and in coordinating and implementing the cottonwood harvest on the Fitler Plantation. Encouragement to conduct this study was provided by Pat Weber of Creative Forestry and Jeff Portwood of Temple-Inland Forest Products Corporation. Bryce Burke, Beth Corbin, Sam Franklin, and Ben Ware provided invaluable assistance in the field.

LITERATURE CITED

- Belli, K.L.; Hart, C.P.; Hodges, J.D.; Stanturf, J.A. 1999. Assessment of the regeneration potential of red oaks and ash on minor bottoms of Mississippi. Southern Journal of Applied Forestry. 23: 133-138.
- Deen, R.T.; Hodges, J.D.; Janzen, G.C. 1993. The influence of midstory-understory control treatments on regeneration of red oak species following clearcutting. In: Brissette, J.C., ed. Proceedings of the seventh biennial southern silvicultural research conference. Gen. Tech. Rep. SO-93. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 117-121.
- Gardiner, E.S. 1998. Water relations and morphology of Nuttall oak seedling sprouts. In: Waldrop, T.A., ed. Proceedings of the ninth biennial southern silvicultural research conference. Gen. Tech. Rep. SRS-20. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 284-288.
- Gardiner, E.S.; Oliver, J.M. 2005. Restoration of bottomland hardwood forests in the Lower Mississippi Alluvial Valley, U.S.A. In: Stanturf, J.A.; Madsen, P., eds. Restoration of boreal and temperate forests. Boca Raton, FL: CRC Press: 235-251.

- Gardiner, E.S.; Russell, D.R.; Oliver, M.; Dorris, L.C., Jr. 2002. Bottomland hardwood afforestation: state of the art. In: Holland, M.M.; Warren, M.L.; Stanturf, J.A., eds. Proceedings of a conference on sustainability of wetlands and water resources: how well can riverine wetlands continue to support society into the 21st Century? Gen. Tech. Rep. SRS-50. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 75-86.
- Gardiner, E.S.; Schweitzer, C.J.; Stanturf, J.A. 2001. Photosynthesis of Nuttall oak (*Quercus nuttallii* Palm.) seedlings interplanted beneath an eastern cottonwood (*Populus deltoides* Bartr. ex Marsh.) nurse crop. Forest Ecology and Management. 149: 283-294.
- Gardiner, E.S.; Stanturf, J.A.; Schweitzer, C.J. 2004. An afforestation system for restoring bottomland hardwood forests: biomass accumulation of Nuttall oak seedlings interplanted beneath eastern cottonwood. Restoration Ecology. 12: 525-532.
- Hamel, P.B. 2003. Winter bird community differences among methods of bottomland hardwood forest restoration: results after seven growing seasons. Forestry. 76: 189-197.
- Hodges, J.D.; Gardiner, E.S. 1993. Ecology and physiology of oak regeneration. In: Loftis, D.; McGee, C.E., ed. Oak regeneration: serious problems, practical recommendations. Gen. Tech. Rep. SE-84. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 54-65.
- Janzen, G.C.; Hodges, J.D. 1987. Development of advanced oak regeneration as influenced by removal of midstory and understory vegetation. In: Phillips, D.R., comp. Proceedings of the fourth biennial southern silvicultural research conference. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 455-461.

- Lockhart, B.R.; Hodges, J.D.; Gardiner, E.S. 2000. Response of advance cherrybark oak reproduction to midstory removal and shoot clipping. Southern Journal of Applied Forestry. 24: 45-50.
- Stanturf, J.A.; Gardiner, E.S.; Hamel, P.B. [and others]. 2000. Restoring bottomland hardwood ecosystems in the Lower Mississippi Alluvial Valley. Journal of Forestry. 98: 10-16.
- Stanturf, J.A.; Gardiner, E.S.; Schoenholtz, S. 2003. Interplanting for bioenergy and riparian restoration in the southern USA. In: Nicholas, I.D., comp. Proceedings of the conference on short rotation crops for bioenergy. [Place of publication unknown]: International Energy Agency, Bioenergy Task. 30: 241-249.
- Stanturf, J.A; Kennedy, H.E., Jr. 1996. Survival and growth of planted and direct-seeded cherrybark oak in South Carolina. Southern Journal of Applied Forestry. 20: 194-196.
- Stanturf, J.A.; Portwood, C.J. 1999. Economics of afforestation with eastern cottonwood (*Populus deltoides*) on agricultural land in the Lower Mississippi Alluvial Valley. In: Haywood, J.D., ed. Proceedings of the 10th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS-30. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 66-72.
- Stanturf, J.A.; Schoenholtz, S.H.; Schweitzer, C.J.; Shepard, J.P. 2001. Achieving restoration success: myths in bottomland hardwood forests. Restoration Ecology. 9: 189-200.
- Wynn, A.H., Jr.; Alcorn, D.E.; Parker, R.L. 1959. Soil survey of Issaquena County Mississippi. Series 1959, no. 2. Washington, DC: U.S. Department of Agriculture, Soil Conservation Service. 42 p.

Great Trinity Forest Management Plan

HARDWOOD SILVICULTURE

The Role of Large Container Seedlings in Afforesting Oaks in Bottomlands

(Connor, Kristina F., ed. 2006. Proceedings of the 13th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS-92. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 640 p.)

THE ROLE OF LARGE CONTAINER SEEDLINGS IN AFFORESTING OAKS IN BOTTOMLANDS

Daniel C. Dey, John M. Kabrick, Michael Gold¹

Abstract—We planted large container (RPM[®]) and 1-0 bareroot seedlings of pin oak (*Quercus palustris* Muenchh.) and swamp white oak (*Q. bicolor* Willd.) in crop fields in the Missouri River floodplain. We also evaluated the benefits of soil mounding and a grass (*Agrostis gigantea* Roth) cover crop. RPM[®] oak seedlings had significantly greater survival and basal diameter increment after 3 years than bareroot seedlings. RPM[®] trees lost significantly more height during the first 3 years than bareroot seedlings. Cover grew substantially greater in the natural vegetation than the redtop grass fields. Oak seedlings in redtop grass cover grew substantially more in diameter and height than oaks competing with natural vegetation. Soil mounding had no significant effect on oak survival or growth. Swamp white oak RPM[®] seedlings produced acorns annually the first 4 years. Planting large container seedlings in redtop grass improved early oak regeneration success and rapidly restored acorn production.

INTRODUCTION

Public land managers and private land owners have a strong interest in regenerating native oak species (*Quercus* sp.) on what are largely agricultural floodplains. Bottomland oak species are highly valued for timber products and wildlife habitat. They are also of conservation concern because of the substantial decline in oaks from historic levels and the difficulty in regenerating them on highly productive floodplains.

The Great Flood of 1993 inundated floodplains throughout much of the summer ruining bottomland farms and causing extensive mortality of oaks in forests along the Missouri and Mississippi Rivers. Since 1993, many abandoned bottomland crop fields have naturally regenerated to forests dominated by pioneering species such as cottonwood (Populus deltoides Bartr. ex Marsh.), silver maple (Acer saccharinum L.), sycamore (Platanus occidentalis L.), and willow (Salix nigra Marsh.). These species are abundant in remnant floodplain forests. thus ensuring a local seed supply. They are prolific annual seed producers, and their seed are easily dispersed by wind and water. Seed germination is favored on mineral soils in open environments, typical of conditions following abandonment of bottomland crop fields. Their seedlings exhibit rapid juvenile growth, which makes them highly competitive on the productive bottomland soils.

Former small- to moderate-sized bottomland crop fields have developed into well-stocked sapling stands dominated by these pioneer species in the years since 1993. Cottonwood, willow, sycamore, silver maple, and other early successional tree species are native to bottomlands and are considered desirable reproduction. A widespread pattern in forest succession on former bottomland crop fields is the lack of oaks and other nut-producing trees. For example, Shear and others (1996) found a lack of hard mast species in 50-year-old forests that naturally regenerated on bottomland crop fields in southwestern Kentucky. Thus, artificial regeneration of oaks is needed to increase the likelihood that oaks are present in future forests. Traditional methods of planting bareroot oak seedlings or direct seeding acorns in bottomlands have not always been successful. For example, in a survey of 4-year-old Wetland Reserve Program plantings in the Mississippi River floodplain, Schweitzer and Stanturf (1997) found that only 9 percent of the total reforested land in 13 Mississippi counties met the Natural Resources Conservation Service requirement for at least 125 hard mast stems per acre in 3-year-old stands.

Oak regeneration failures in bottomland crop fields are largely a result of the low competitiveness of small oak seedlings on sites that are capable of producing tremendous herbaceous biomass in one summer. Small oak seedlings also are less competitive than the pioneer tree species that invade abandoned crop fields. In addition, oak plantings are not often maintained by controlling competing vegetation, which makes successful oak regeneration less likely. Oak species are moderately tolerant to intolerant of shade and unable to persist in the heavy shade of competing vegetation.

Oak regeneration success can be improved, in part, by planting large seedlings, particularly those having well-developed root systems (Gardiner and others 2002, Johnson and others 2002, Kormanik and others 1998, Schultz and Thompson 1997, Stanturf and others 1998). Nursery managers can produce hardwood bareroot seedlings with large root systems that have five or more large lateral roots by undercutting the taproot, growing seedlings for a second year. Air pruning the roots of seedlings grown in open-bottomed containers is another way to promote the growth of large lateral roots and dense fibrous root systems (Dey and others 2004).

In the Midwest, several private nurseries have begun commercial production of large (e.g., 3- to 5-gallon) container seedlings that are being used in the afforestation of bottomland crop fields on public and private lands. The Forrest Keeling Nursery of Elsberry, MO, has developed a nursery cultural technique, the Root Production Method (RPM®), to produce high-quality hardwood seedlings that have large basal diameter and height and a substantial fibrous root system (Dey

¹ Research Foresters, USDA Forest Service, North Central Research Station, Columbia, MO 65211; and Associate Director, University of Missouri, Center for Agroforestry, Columbia, MO 65211, respectively.

Citation for proceedings: Connor, Kristina F., ed. 2006. Proceedings of the 13th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS–92. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 640 p.

and others 2004). For the past 5 years or so, these large container seedlings have been planted in various floodplain situations throughout the Midwest. In 1999, we began a study to evaluate the performance of these large container seedlings in the afforestation of bottomland crop fields. This paper presents early survival and growth performance of large container seedlings in regenerating pin oak and swamp white oak in bottomland crop fields along the lower Missouri River. We discuss the role of large container seedlings in regenerating oaks in floodplains.

METHODS

Dey and others (2003) provided a detailed explanation of the experimental design and study establishment. In summary, a study was established to evaluate methods for regenerating pin oak and swamp white oak on former agricultural crop fields in the Missouri River floodplain in the fall of 1999. The study fields had been in crop production for years before this study. Soils at the study sites were mapped as Sarpy Fine Sand (mixed, mesic, Typic Udipsamments), Haynie Silt Loam (coarsesilty, mixed, superactive, calcarious, mesic Mollic Udifluvents), and Leta Silty Clay (clayey over loamy, smectitic, calcareous, mesic Aeric Fluvaquents). Two floodplain properties were used in this study, Smoky Waters and Plowboy Bend Conservation Areas, both managed by the Missouri Department of Conservation. The Smoky Waters site, which lies at the confluence of the Missouri and Osage Rivers, is not protected by levees and is subject to flooding. The site at Plowboy Bend, located along the Missouri River, is protected by a levee.

Both 1-0 bareroot and 3- and 5-gallon RPM[®] seedlings were planted to evaluate the effect of seedling size and nursery stock type on the survival and growth of pin oak and swamp white oak seedlings. Seedlings were planted in soil mounds created with a rice plow or in unmounded soil and with either a cover crop of redtop grass or with natural vegetation that normally colonizes abandoned bottomland crop fields.

Dey and others (2004) have described the RPM[®] cultural technique in some detail. Using oak as an example, the process generally involves (1) germinating seed on soil media in shallow, open-mesh, bottomless trays, (2) transplanting seedlings into 4-inch square bottomless band containers at the end of the first shoot flush, and (3) a final transplanting of seedlings into 3- or 5-gallon pots whether they are grown in the nursery for 1 or 2 years, respectively.

Initial total height and basal stem diameter 1 inch above the ground were measured on all seedlings after planting and again at the end of each growing season. Seedling survival was determined each year, and animal damage to seedlings was recorded. Animal damage consisted primarily of shoot clipping and girdling of main stems by eastern cottontail rabbits (*Sylvilagus floridanus* Allen) and white-tailed deer (*Odocoileus virginianus* Boddaert).

Analysis of variance (ANOVA) was used to determine significant (P < 0.05) growth rate differences among species and stock type combinations between fields with and without redtop grass and between mounded and non-mounded planting units. In this ANOVA, site was a random effect and cover crop treatment, mounding treatment, and species and stock type combinations were fixed effects. The significance of treatment effects on tree growth and survival was determined using their respective site interactions as error terms. For significant effects, orthogonal contrasts were used to compare mean differences. Logistic regression was used to develop models that predict survival based on seedling characteristics and management treatments.

RESULTS AND DISCUSSION

Comparison of Initial Planting Stock

Average basal diameters of 5-gallon RPM[®] seedlings were slightly larger than 3-gallon seedlings for both pin oak and swamp white oak, and diameters of both RPM[®] stock types were substantially larger than the bareroot seedlings (fig. 1). Pin oak and swamp white oak RPM[®] seedlings averaged 0.6 to 0.8 inches in basal diameter compared to 0.3 inches for bareroot seedlings. Similar trends were seen in average height of seedlings among the different stock types. Heights of RPM[®] seedlings averaged 4.9 feet or larger compared to bareroot seedlings, which averaged 2.3 feet or less. Pin oaks were generally slightly larger than swamp white oaks within a given stock type.




Figure 1—Average basal diameter (1-inch above the root collar) and shoot length of pin oak and swamp white oak RPM[®] and bareroot seedlings typical of those used in this study.

Table 1—Root system differences between 1-0 bareroot
(BR) and RPM [®] pin oak and swamp white oak seedlings
planted in this study

Stock type	Root volume	Root dry weight
	$cm^3 H_2O$	g
Pin oak BR	26	17
Pin oak RPM [®] 3 gallon	236	117
Pin oak RPM [®] 5 gallon	223	118
Swamp white oak BR	33	20
Swamp white oak RPM [®] 3 gallon	141	64
Swamp white RPM® 5 gallon	252	138

The root systems of RPM[®] seedlings had substantially more volume and dry weight than bareroot seedlings, regardless of species (table 1). The root systems of 3- and 5-gallon pin oak RPM[®] seedlings were similar in volume and dry weight. For all RPM[®] seedlings, root volume and dry weight were greatest for 5-gallon swamp white oak RPM[®] seedlings and smallest for 3-gallon swamp white oak. However, the smallest RPM[®] root systems were substantially larger than those of bareroot seedlings.

The RPM[®] process builds a large root mass with numerous sizeable lateral roots concentrated just below the root collar in a short period of time (e.g., 210 days to produce a plant-able 3-gallon pin oak seedling). Johnson and others (2002), Kormanik and others (1998), and Schultz and Thompson (1997) have emphasized that the key to successful oak regeneration, both natural and artificial, is the formation of a large root system. Any silvicultural systems or nursery practices that create favorable conditions for the growth of oak root systems enhance oak's competitiveness.

The perceived advantages of container grown RPM[®] seedlings include (1) rapid early growth that is driven by a large, intact root system, which minimizes transplant shock and facilitates early root growth in the establishment year; and (2) seedlings tall enough (e.g., \geq 5 feet) to minimize deer herbivory on the terminal shoot and to improve chances that the live crown will remain above growing-season floods. Survival of seedlings is greatly enhanced if their live crowns avoid inundation during growing-season floods. Floods that cause defoliation of hardwood seedlings force trees to expend their remaining energy forming a second crown, which places them at a greater disadvantage with other competing vegetation on these highly productive sites.

Natural Disturbances

Rabbit herbivory—Every winter after the first year, cottontail rabbits have girdled and shoot clipped oak seedlings and oak sprouts. The amount and severity of rabbit damage to planted oaks varied greatly between the cover crop treatments (i.e., redtop grass versus natural vegetation fields).

In the natural vegetation fields, the composition and structure of winter cover provided by forbs promoted higher rabbit densities (3 rabbits per acre) than in the redtop grass fields (1.0 rabbit per acre) (Dugger and others 2004). In the winter, the dead tops of forbs and clumps of Johnsongrass [*Sorghum*

halepense (L.) Pers.] remained somewhat erect, providing cover that was as high as 3 feet. However, redtop grass matted down to 0.6 feet and provided little hiding cover for rabbits from predators. Thus, rabbits were able to move freely across the natural vegetation fields causing damage to nearly all of the seedlings each winter. Rabbits clipped the shoots of all bareroot seedlings and severely girdled (more than half of the circumference of the stem) 90 percent or more of the RPM® seedlings in the natural vegetation fields by the end of the second winter. In comparison, damage to oak seedlings by rabbits was much less extensive and severe in the redtop grass fields. These differences in winter habitat between the cover crop treatments affected rabbit densities and movements, which in turn contributed to the significant differences in oak seedling survival, growth, and acorn production between the cover crop treatments.

Flooding—In 2 of the past 4 years (2001, 2002), the study site at Smoky Waters was flooded for up to 3 weeks in June. Duration of flooding was variable across the redtop grass and natural vegetation fields due to differences in elevation. Depth of flooding was also variable, with maximum depths reaching 4 to 5 feet, enough to completely inundate bareroot seedlings and oak sprout clumps formed by rabbit-girdled RPM® trees.

Survival

Survival of oak RPM[®] seedlings remained high (> 94 percent) during the first 3 years (fig. 2), while survival of bareroot seedlings continued to decline for both swamp white oak and pin oak. There was no significant difference in survival between 3- and 5-gallon RPM[®] seedlings. After 3 years in the field, survival of swamp white oak bareroot seedlings was significantly higher (P < 0.01) than pin oak bareroot seedlings. Swamp white oak bareroot survival averaged 76 percent, while survival for pin oak bareroot seedlings was 54 percent. There was no significant difference (P = 0.24) between 3- and 5-gallon RPM[®] seedling survival, nor between swamp white oak and pin oak RPM[®] seedling survival (P = 0.87). Soil mounding and cover crop treatments did not significantly affect oak seedling survival.

Basal Diameter Growth

Basal diameter increment after 3 years was significantly greater (P < 0.01) for RPM[®] seedlings than bareroot stock, regardless of species (fig. 2). There was no significant difference in basal diameter increment between 3- and 5-gallon RPM® seedlings. The average basal diameter of all RPM® oak seedlings increased 0.3 inches in the first 3 years, whereas bareroot seedlings increased only 0.1 inches. There was no significant difference (P = 0.34) in basal diameter increment between the 3- and the 5-gallon RPM® seedlings. The basal diameter of pin oak 5-gallon RPM® seedlings increased the most during the first 3 years, averaging 0.4 inches of new growth. Basal diameter increment was least (0.03 inches in 3 years) for pin oak bareroot seedlings. The above analysis includes rabbit-damaged and undamaged trees. By removing the rabbit-damaged trees, average basal diameter increment was 0.6 inches for RPM® seedlings and 0.1 inches for bareroot trees.

Although soil mounds functioned as anticipated by improving drainage and aeration, they did not significantly affect diameter growth. Kabrick and others (2005) have reported more




Figure 2—Survival, basal diameter and height of swamp white oak and pin oak RPM[®] and bareroot seedlings during the first 3 years in this study.

detailed effects of soil mounding on soil properties and oak regeneration based on the results from this study. Basal diameter increment of all trees combined was substantially larger in redtop grass fields (0.6 inches) than in natural vegetation fields (0.1 inches); however, no significant differences can as yet be reported (P = 0.08). For undamaged trees, average basal diameter increment was 0.6 inches for RPM[®] seedlings in redtop grass fields and 0.2 inches in natural vegetation fields, while the basal diameter of bareroot seedlings increased by 0.1 inches in redtop grass but decreased by 0.1 inches in natural vegetation fields.

Height Growth

Average height increment after 3 years was negative for most species and nursery stock types, because cottontail rabbits caused extensive damage by girdling the stems of RPM[®] seedlings or by clipping the shoots of bareroot seedlings at ground-level (fig. 2). Severely girdled or clipped trees that survived produced multiple-stemmed sprout clumps, but new shoot growth was insufficient to recover to original seedling heights, especially for RPM[®] seedlings.

Three-year height increment was significantly less (P < 0.01) for RPM® than bareroot seedlings. There was no significant difference in height increment between 3- and 5-gallon RPM® seedlings. For bareroot seedlings that had been shoot-clipped by rabbits, annual sprout growth came close to, or slightly exceeded, the initial height, resulting in small negative or positive increments in height. In contrast, net height increment was much lower in initially tall RPM® trees, because rabbit girdling, which occurred in the lower 1.0 foot of the stem, caused shoot dieback to near ground-level. In addition, trees were often repeatedly damaged by rabbits each winter. Three-year height increment averaged -1.6 feet for the RPM® seedlings. Despite rabbit browsing, RPM® trees remained taller than bareroot seedlings 3 years after planting. Undamaged trees in the redtop grass fields had slightly positive average height growth (0.3 feet for bareroot and RPM® seedlings), but net growth was negative in natural vegetation fields, averaging -1.7 feet for RPM® and -0.4 feet for bareroot seedlings.

Height growth of undamaged RPM[®] seedlings may be low because these trees were planted on a 30 by 30 foot spacing; and widely spaced, open-grown trees often experience reductions in height growth, especially trees with weak epinastic control such as the oaks (Oliver and Larson 1996). Also, height growth of oak reproduction is slow at first, because seedlings characteristically allocate photosynthates to root growth often at the expense of shoot growth (Johnson and others 2002). Height growth of RPM[®] and bareroot seedlings may also be limited by low levels of foliar nitrogen, which averaged 2.05 percent at Smoky Waters and 1.71 percent at Plowboy Bend Conservation Areas (Kabrick and others 2005).

An analysis of all trees by cover crop treatment showed that 3-year height increment was significantly higher (P = 0.02) for oak seedlings growing in the redtop grass fields than those trees competing with natural vegetation. There may be less light competition in the redtop grass fields during the growing season than in the natural vegetation fields. Redtop grass typically grows to a height of 1.5 to 2.0 feet, whereas herbaceous ground cover in the natural vegetation fields grew to over 6.6 feet in height, overtopping many of the oak seedlings.

Also, rabbit densities were less and fewer trees were damaged in the redtop grass than in the natural vegetation fields. There was no significant difference in height growth among trees on mounded and unmounded soils.

Acorn Production

Swamp white oak RPM[®] seedlings that were 18 to 24 months old at time of planting produced acorns in each of the first 4 years following outplanting (table 2). Acorn production occurred in a small proportion (3.5 percent) of the 2,522 swamp white oak RPM[®] seedlings their first year in the field. Most of the production (60 percent) occurred in oaks from 5-gallon containers, but larger 3-gallon container RPM[®] trees also produced acorns. Individual trees were able to produce as many as 125 acorns. The probability of a RPM[®] swamp white oak seedling producing at least one sound acorn in the first year after planting was significantly related to initial basal diameter and height of the seedling (Grossman and others 2003).

The number of trees bearing acorns dropped in year 2 because rabbit herbivory and shoot dieback took RPM® trees in the natural vegetation fields out of production. Production in years 3 and 4 came almost exclusively from RPM® trees growing in the redtop grass fields. A single pin oak RPM® seedling produced acorns for the first time in the fourth year. Consistent, early production of acorns is surprising considering that open-grown oaks do not begin producing seed until they are 20 to 30 years old (Burns and Honkala 1990). In contrast, no bareroot oak seedlings have produced acorns after four growing seasons. Early fruiting and nut production in RPM® trees is a substantial benefit for wildlife, and it restores the ability for oaks to regenerate naturally on former bottomland crop fields where lack of local seed supply is a factor limiting oak regeneration.

Role of Large Container Seedlings in Regenerating Hard Mast Species

In the afforestation of bottomland crop fields, managers need to consider the contributions that can be expected from natural regeneration and the need for artificial regeneration. In extremely large fields, artificial regeneration may be needed even for pioneering species such as cottonwood, silver maple, sycamore, and willow (Allen 1990, Twedt and Wilson 2002). Relying on natural regeneration in large fields may result in understocked stands, in part because of the difficulty in timing an appropriate seedbed and adequate competition control with seed dispersal to the site. In smaller fields, pioneering species often are able to regenerate to well-stocked stands

Table 2—Acorn production in swamp white oak RPM[®] seedlings during the first 4 years at Smoky Waters and Plowboy Bend Conservation Areas

Year	No. nut- bearing trees	Average no. acorns per tree
1	86	4.3
2	29	5.2
3	70	6.3
4	152	12.5

because establishment conditions are easier to control through site preparation and seed sources are often adjacent to the area. Regardless of field size, managers are almost always forced into artificially regenerating heavy nut species such as the oaks due to the lack of a natural seed source.

Planting bareroot seedlings and direct seeding are the most common methods for regenerating oaks in agricultural floodplains. Planting container seedlings is less common, especially trees grown in very large (3- to 5-gallon) containers. Of course, the initial cost of these seedlings is much higher than bareroot seedlings. For example, a 3-gallon container oak seedling may cost \$8.00 compared to \$0.50 to \$1.00 for a bareroot seedling. Our study is too young to consider an economic analysis, which should consider the cost of the entire regeneration prescription needed to produce a successful tree, one that is free-to-grow and that will be in a codominant or dominant position in the mature forest.

We have seen that large container seedlings have significantly higher survival and diameter growth than bareroot seedlings after 3 years. It is still too early in our study to determine if a planted oak is successful or not in the context of our definition of success but if trends continue, it will take fewer large container seedlings to get a successful tree than if bareroot seedlings are planted.

The manager's objectives must be considered before identifying artificial regeneration methods for restoring forests to agricultural floodplains. Public managers often want hard mast species as a component of a native bottomland forests. Thus, planting large container seedlings on a 60 by 60 foot spacing (12 trees per acre) may be an affordable way to increase the probability that hard mast species will be a component of the overstory and capable of producing good mast crops periodically. Interplanting with bareroot seedlings or direct seeding may be used to supplement hard mast regeneration and diversify the forest. Natural regeneration of pioneer species may fill in if an adequate seed source is available and site preparation is properly done.

In managing the greater floodplain property, which may be as large as 3,000 acres or more, it may be desirable to have hard mast production somewhere on the property but not necessarily on every acre. In this case, managers may choose to plant large container seedlings on tighter spacings, e.g., 30 by 30 feet (48 trees per acre) or 20 by 20 feet (109 trees per acre) and manage them more intensively on small parcels, e.g., 1 to 5 acres or so, located throughout the floodplain.

Commonly on abandoned bottomland crop fields, young (< 10 years), dense sapling stands of cottonwood, willow and sycamore have developed. In these stands, managers may consider planting large container oak seedlings in group openings. Openings can be created by combinations of mechanical and chemical treatments. Or, large container seedlings may be underplanted in thinned cottonwood sapling stands.

The early nut-producing capacity demonstrated by large container oak seedlings is of particular interest to owners of properties managed as waterfowl hunt clubs or public greentree reservoirs. In many cases, acorn production on these properties is limited by the low abundance of oak trees, by the advanced age of oak trees that are declining in their acorn production capacity, or because many of the mature oak trees were lost in a catastrophic flood such as the Great Flood of 1993. Although the initial cost of large container stock is relatively high compared to bareroot seedlings, acorn production is so desirable and such a part of waterfowl management that greentree reservoir managers and duck club owners are willing to pay for restoring acorn production.

There is no single silver bullet for regenerating oaks and other hardmast species in agricultural floodplains. More often managers are designing afforestation projects using a diversity of species and by combining natural regeneration with a variety of artificial regeneration techniques including direct seeding, planting bareroot and container seedlings, and using cuttings as a means of vegetatively reproducing a species. This integrated approach to regeneration is more likely to produce a diverse forest that is more resilient to flooding and other perturbations than are less-diverse monoculture forests and plantations.

Plant succession proceeds rapidly once bottomland crop fields are abandoned. Hence, there is a narrow window of opportunity for establishing trees in high enough stocking that they dominate the growing space before other competing vegetation establishes its dominance. Using natural and artificial regeneration in a planned manner improves the chances of obtaining well-stocked forests. Planting large container stock may be one way of ensuring that the forest has a component of oak.

ACKNOWLEDGMENTS

This work was funded, in part, through the University of Missouri Center for Agroforestry under cooperative agreements 58-6227-1-004 with the ARS and C R 826704-01-2 with the US EPA. The results presented are the sole responsibility of the P.I. and/or MU and may not represent the policies or positions of the ARS or EPA.

LITERATURE CITED

- Allen, J.A. 1990. Establishment of bottomland oak plantations on the Yazoo National Wildlife Refuge Complex. Southern Journal of Applied Forestry. 14: 206-210.
- Burns, R.M.; Honkala, B.H., eds. 1990. Silvics of North America, vol. 2. Hardwoods. Agric. Handb. 654. Washington, DC: U.S. Department of Agriculture, Forest Service. 877 p.
- Dey, D.C.; Kabrick, J.M.; Gold, M.A. 2003. Tree establishment in floodplain agroforestry practices. In: Proceedings of the eighth North American agroforestry conference. [CD-ROM]. Corvallis, OR: Association for Temperate Agroforestry: 102-115.
- Dey, D.C.; Lovelace, W.; Kabrick, J.M.; Gold, M.A. 2004. Production and early field performance of RPM[®] seedlings in Missouri floodplains. In: Michler, C.H.; Pijut, P.M.; Van Sambeek, J.W. [and others], eds. Proceedings of the sixth walnut council research symposium. Gen. Tech. Rep. NC-243. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Station: 59-65.

- Dugger, S.; Dey, D.C.; Millspaugh, J.J. 2004. Vegetation cover effects mammal herbivory on planted oaks and success of reforesting Missouri River bottomland fields. In: Connor, K.F., ed. Proceedings of the 12th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS-71. Asheville, NC: U.S. Dept of Agriculture, Forest Service, Southern Research Station: 3-6.
- Gardiner, E.S.; Russell, D.R.; Oliver, M.; Dorris, L.C., Jr. 2002. Bottomland hardwood afforestation: state of the art. In: Holland, M.M.; Warren, J.L.; Stanturf, J.A., eds. Proceedings of a conference on sustainability of wetlands and water resources. Gen. Tech. Rep. SRS-50. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 75-86.
- Grossman, B.C.; Gold, M.A.; Dey, D.C. 2003. Restoration of hard mast species for wildlife in Missouri using precocious flowering oak in the Missouri River floodplain, USA. Agroforestry Systems. 59: 3-10.
- Johnson, P.S.; Shifley, S.R.; Rogers, R. 2002. The ecology and silviculture of oaks. Wallingford, Oxon, UK: CABI Publishing CAB International. 503 p.
- Kabrick, J.M.; Dey, D.C.; Van Sambeek, J.W. [and others]. 2005. Soil properties and growth of swamp white oak and pin oak on bedded soils in the lower Missouri River floodplain. Forest Ecology and Management. 204: 315-327.
- Kormanik, P.P.; Sung, S.S.; Kass, D.J.; Schlarbaum, S. 1998. Effect of seedling size and first-order-lateral roots on early development of northern red oak on mesic sites. In: Waldrop, T.A., ed. Proceedings of the ninth biennial southern silvicultural research conference. Gen. Tech. Rep. SRS-20. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 247-252.
- Oliver, C.D.; Larson, B.C. 1996. Forest stand dynamics. New York, NY: John Wiley & Sons, Inc. 520 p.
- Schultz, R.C.; Thompson, J.R. 1997. Effect of density control and undercutting on root morphology of 1-0 bareroot hardwood seedlings: five-year field performance of root-graded stock in the central USA. New Forests. 13: 301-314.
- Schweitzer, C.J.; Stanturf, J.A. 1997. From okra to oak: reforestation of abandoned agricultural fields in the Lower Mississippi Alluvial Valley. In: Meyer, D.A., ed. Proceedings of the 25th annual hardwood symposium. Memphis, TN: National Hardwood Lumber Association: 131-138.
- Shear, T.H.; Lent, T.J.; Fraver, S. 1996. Comparison of restored and mature bottomland hardwood forests of southwestern Kentucky. Restoration Ecology. 4(2): 111-123.
- Stanturf, J.A.; Schweitzer, C.J.; Gardiner, E.S. 1998. Afforestation of marginal agricultural land in the Lower Mississippi River Alluvial Valley, U.S.A. Silva Fennica. 32(3): 281-287.
- Twedt, D.J.; Wilson, R.R. 2002. Supplemental planting of early successional tree species during bottomland hardwood afforestation. In: Outcalt, K.W., ed. Proceedings of the 11th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS-48. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 358-364.

Great Trinity Forest Management Plan

HARDWOOD SILVICULTURE

A Comparison of Oak Regeneration Conditions Following Midstory Injection and Partial Overstory Removal in a Tombigbee River Terrace

(Connor, Kristina F., ed.2004. Proceedings of the 12th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS – 71. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 594 p.)

A COMPARISON OF OAK REGENERATION CONDITIONS FOLLOWING MIDSTORY INJECTION AND PARTIAL OVERSTORY REMOVAL IN A TOMBIGBEE RIVER TERRACE

Stephen E. Peairs, Andrew W. Ezell, Keith L. Belli, and John D. Hodges¹

Abstract—Bottomland hardwood stands comprised of a large oak component can be regenerated through natural oak regeneration. To maximize oak regeneration potential, the stand manager must practice silvicultural techniques that allow for necessary sunlight penetration to reach the forest floor. Control of stems occupying the midstory/understory and partial overstory removal are steps that may be required to ensure adequate stocking levels of natural regeneration. While the need for midstory/understory control has been documented, questions often arise regarding the level of intensity of overstory removal. Natural regeneration responses of oak species were observed 1 year after a midstory/understory injection treatment and partial overstory removal. Treatments were conducted on a bottomland hardwood stand near Bellamy, AL. The study area consisted of four 12-acre blocks with three 4-acre treatment units located in each block. The study design was a randomized complete block. Injection treatments using imazapyr were applied during late fall of 2001. There was no negative impact from the injection observed on non-target stems, and seedling establishment generally increased. Partial overstory removal treatments were completed with operational harvests, leaving residual stand basal areas of 30, 50, and 70 square feet. Seedling survival was observed from data collected from 108 0.01-acre sample plots. Results indicate that seedling survival was impacted by harvest. Damage occurring from logging operations was also evaluated on all residual trees.

INTRODUCTION

To establish natural regeneration in a hardwood stand, land managers need an understanding of silvicultural procedures involved. Silvicultural applications such as midstory/ understory control and partial harvests may be required to ensure adequate stocking of oak regeneration to promote the future stand. Control of undesirable species using imazapyr causes mortality to injected stems, enhances regeneration, and does not damage crop stems (Ezell and others 1999). The level of the partial harvest intensity may impact the amount of advance reproduction present following harvesting activity. The intensity of partial harvest affects the amount of sunlight penetration that reaches the forest floor. Gardiner and Hodges (1998) found that oak seedling growth was maximized when sunlight availability was between 27-53 percent. Low understory light levels present in bottomland stands may be the most limiting factor on the establishment and growth of oaks (Hodges and Gardiner 1993). Thus, land managers need to understand how these practices can influence oak regeneration.

The amount of harvesting damage occurring to residual stems following harvesting activity is also an important consideration. Potential losses due to reduction of residual stems' log grade can be assessed if an estimation of damage can be depicted. The intensity of partial harvesting might also impact the amount of initial oak regeneration following harvesting. The objectives for this study are (1) to evaluate the effect of midstory/understory control on initial oak regeneration, (2) determine the impact of harvesting activity on initial oak regeneration, and (3) to evaluate damage resulting from varying harvest intensities.

MATERIALS AND METHODS Study Site Description

The study site is located near Bellamy, AL on property owned by Delaney Development, Inc. and managed by Springdale Land Management. The hardwood stand is situated on a terrace micro-geographical position along the Tombigbee River. Stand composition is a mixed oak bottomland forest with a well developed midstory. Species occupying the overstory include water oak (Quercus nigra L.), swamp chestnut oak (Q. michauxii Nutt.), white oak (Q. alba L.), cherrybark oak (Q. pagoda Raf.), sweetgum (Liquidambar styraciflua L.), and various hickory species (Carva spp.). The understory and midstory tree species primarily consist of American hornbeam (Carpinus caroliniana Walt.), eastern hophornbeam (Ostrya virginiana Mill.), American beech (Fagus grandifolia Ehrh.), and winged elm (Ulmus alata Michx.), with scattered oak species.

STUDY DESIGN

The study was arranged as a randomized complete block design. The research site consisted of four rectangular blocks that were 12.06 acres in size. Three of these blocks were utilized as "treatment areas" while retaining one block as control. A midstory/understory control treatment using imazapyr was applied during the dormant season of 2001-2002. All four blocks were divided into three treatment units encompassing an area of 4.02 acres each. The experimental unit consisted of treatment unit conditions prior to and following partial harvests which maintained residual basal areas of 70, 50, and 30 square feet. These partial harvest intensities were replicated once in treatment units for each of the three "treatment areas."

¹Graduate Research Assistant, Professor, Professor, and Professor Emeritus, College of Forest Resources, Mississippi State University, Mississippi State, MS, respectively.

Citation for proceedings: Connor, Kristina F., ed. 2004. Proceedings of the 12th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS–71. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 594 p.

The partial harvest activity occurred during September of 2002. These harvests were conducted by Kynard Logging contracted by Springdale Land Management. Large diameter trees were felled with chainsaws then topped, delimbed, and cut to length prior to being skidded. Stems with large canopies used for pulpwood were topped and limbed in place. Smaller sized stems used for pulpwood were topped and delimbed at the loading decks. Wheeled, articulated skidders were used to move raw wood materials to loading decks located at the east boundaries of the treatment blocks. Weather conditions during the time of harvest were dry.

DATA COLLECTION AND ANALYSIS

Pre-harvest overstory basal area was inventoried on the site. All overstory stems were measured for diameter at breast height (d.b.h.). Basal area for individual stems was determined using the formula:

basal area = $d.b.h.^2 * 0.005454$.

These data allowed for estimations of basal area for all species occupying treatment units and served as a basis for determining trees to be removed to achieve desired basal areas following partial harvests.

Oak regeneration sampling was conducted on 108 0.01-acre sample plots. Oak seedlings within these plots were tallied by species and height classes. Height classes included seedlings less than 1 foot in height and 1-3 feet in height. Sampling was conducted prior to harvesting activity during April 2002 and following harvesting activity in November 2002. Harvesting damage occurring on residual stems was also quantified. Damage was categorized by height classes and damage area or size classes. Height classes occurred at three-foot intervals starting at ground level. Size classes had a range of 40 square inches (e.g., 0-40, 41-80, etc.).

RESULTS AND DISCUSSION

Regeneration sampling data indicated that the majority of oak seedlings were in the height classes of < 1 foot and 1-3 feet for both pre- and post-harvest measurements. Preharvest measurements were analyzed to determine the impact of applying midstory/understory control. These data compared the three treated blocks against the control block. Results derived from sampling data showed that oak regeneration generally increased following the midstory/ understory control treatment. Results for block 1 indicate that oak regeneration decreased by 5.57 percent compared to control. Blocks 2 and 3 increased by 71.08 percent and 79.44 percent, respectively, compared to control. The increased amount of oak regeneration can be attributed to greater amounts of sunlight penetration reaching the forest floor. The increased abundance of sunlight results from the control of live stems occupying the midstory/understory level of the stand.

The intensity of harvesting activity also impacted the amount of oak regeneration following partial harvest. Evaluations to determine the impact harvesting intensity had on oak regeneration were derived by comparing pre-harvest measurements to post-harvest measurements for the nine treatment units on blocks 1 to 3. Three treatment units suffered the greatest decreases in oak regeneration due to excessive skidding activity. These units were located adjacent to logging decks. These blocks were excluded in the analysis to determine initial oak regeneration responses to partial harvesting intensity. The two treatment units that had residual basal areas of 50 square feet showed the largest increases in initial oak regeneration. This intensity of partial harvest promoted greater regeneration due to the combination of harvest activity and light penetration. Individual treatment unit percentages of increase/decrease are depicted in table 1.

Another observation made from regeneration data was that water and willow oak seedlings increased in abundance more than other oak species. This response was observed on the four treatment units that had increased percentages of initial oak regeneration. These two species had a bumper acorn crop during the fall of 2001. Partial harvest activity occurred following this year, which may have promoted the increase of seedlings from these two species. Findings from this study tend to indicate that conducting harvesting activity following good seed crop years can increase oak regeneration success.

Analysis to determine the impact of harvesting intensity on damage occurring to residual stems showed similar trends between treatments. The majority of damage occurred below 3 feet on the butt logs of the residual trees. Greater than half of the residual trees were undamaged on all treatment units. This indicated that regardless of the partial harvest intensity applied, the majority of residual stems will be undamaged following harvest activity. Treatment units with residual basal areas of 70 square feet had the lowest percentage of damaged residual stems among the three treatments. The percentages of damage occurrence for the three residual basal areas are illustrated in table 2.

Table 1—Harvesting impacts on regeneration

	Treatment— residual basal area	Oak seedling change	Adjacent to loading deck
	square feet	percent	
Block one	50	-59.50	Yes
	30	-3.05	No
	70	-13.45	No
Block two	30	-53.10	Yes
	70	30.22	No
	50	101.00	No
Block three	70	-32.07	Yes
	50	115.93	No
	30	8.15	No

Table 2—Percentage of damage occurrence on residual stems by height class

Height	Basa	Basal area (square feet)				
classes	30	30 50 70				
feet		percent -				
0–3	87.91	77.95	86.51			
3–6	9.89	18.90	11.11			
6–9	2.20	3.15	2.38			

The size of damaged areas on residual stems also showed a similar pattern between treatments. Generally, smaller "damage zones" occurred more frequently than larger "damage zones." The majority of stems had damage areas of less than 80 square inches in size. This can be attributed to proper logging techniques employed by the loggers. The percentages of damage area size compared to residual basal area are presented in table 3.

Table 3—Percentage of residual stems exhibiting various amounts of harvest damage by treatments

		square inches		
Basal area	No damage	< 80	81-160	> 161
square feet		perc	cent	
30	50.57	36.78	9.77	2.87
50	54.89	32.34	5.96	6.81
70	74.84	19.31	2.82	3.04

SUMMARY

The two types of silvicultural treatments conducted during this experiment tended to show positive effects on initial oak regeneration. Midstory/understory control promoted regeneration of desirable species while removing less desirable stems and increasing sunlight penetration to the forest floor. Oak regeneration reached the highest levels on the treatment units that received partial harvests leaving a residual basal area of 50 square feet. The amount and type of initial regeneration was dependent on species present in the stand. This was clearly evident in the case of water and willow oak. Results showed that harvesting in conjunction with a good mast crop tended to enhance initial regeneration. This study also found that there was a strong negative association between skidder activity and oak regeneration. Areas closer to loading decks had a larger reduction in the amount of visible initial regeneration due to increased usage by large machinery. These areas of high impact may, however, regenerate in the future. The majority of damage resulting from harvesting activity will generally tend to be located between the ground and the first 3 feet of butt logs on residual trees. Smaller damaged areas will be more prevalent than larger areas on these logs as well. To reduce the amount of damage occurring to regeneration and residual stems, land managers may opt to either consolidate or spread skid trails out over harvest areas while also ensuring proper logging techniques are applied to hardwood stands.

ACKNOWLEDGMENTS

This project was supported by Delaney Development, Inc. Assistance was also given by Springdale Land Management for contracting of the logging company as well as aiding in the research process.

LITERATURE CITED

- Ezell, A.W.; Lowery, J.; Leopold, B.; Minogue, P.J. 1999. Use of imazapyr injection to promote oak regeneration and wildlife stand improvement in bottomland hardwood stands. In: Haywood, J.D., ed. Proceedings of the tenth biennial southern silvicultural research conference; 1999 February 16-18; Shreveport, LA. Gen. Tech. Rep. SRS-30. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 151-153.
- Gardiner, E.S.; Hodges, J.D. 1998. Growth and biomass distribution of cherrybark oak (*Quercus pagoda* Raf.) seedlings as influenced by light availability. Forest Ecology and Management. 108: 127-134.
- Hodges, J.D.; Gardiner, E.S. 1993. Ecology and physiology of oak regeneration. In: Loftis, D.L.; McGee, C.E., eds. Oak regeneration: Serious problems, practical recommendations. Symposium proceedings; 1992 September 8-10; Knoxville, TN. Gen. Tech. Rep. SE-84. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 54-65.

Great Trinity Forest Management Plan

HARDWOOD SILVICULTURE

Herbicide Release of 4 Year old, Naturally Regenerated Bottomland Oaks-10 Year Results

HERBICIDE RELEASE OF 4 YEAR OLD, NATURALLY REGENERATED BOTTOMLAND OAKS – 10 YEAR RESULTS

Larry E. Nix¹

Abstract—In 1989 two clearcut, naturally regenerated bottomland hardwood stands near the Congaree River in South Carolina were found to contain nearly 2000 red oak (*Quercus pagoda* Raf. *and Q. shumardii* Buckley) seedlings per acre. By age 4, these oak seedlings were quickly being overtopped by competitive sprouts, vines, and fast growing pioneer species. The number of remaining seedlings had been reduced to 450 per acre. In order to improve the competitive status of these desirable oak seedlings, a number of directed and broadcast pine herbicide release treatments at full and half strength were applied late in the fourth growing season. It was hoped that the overtopping competition would be inhibited or killed, but simultaneously shield the oak seedlings from the herbicides. Two year measurements indicated that oak seedling diameter growth was nearly doubled by some of the herbicides, but there was little difference in the number or canopy position of the oaks. Remeasurements 10 years after applications indicated that at least four of the simulated aerial herbicide broadcasts and one of the directed sprays resulted in 14-year-old oak saplings nearly an inch larger in diameter and three times more numerous in the dominant canopy position than those in the untreated areas. One herbicide and method of application in particular resulted in 4-inch diameter oaks, 36 feet tall with nearly 400 per acre remaining in at least a codominant status.

INTRODUCTION

In the past 30 years, oaks (Quercus spp.) have been decreasing in the bottomland hardwood stands of the Southern United States (Johnson 1984). Cherrybark oak (Quercus pagoda Raf.) and Shumard oak (Q. shumardii Buckl.) are two high quality bottomland oak species (Miller and Lamb 1985) that are not being replaced in these stands. The ecology and biology of bottomland oaks have been investigated to determine why the number of oaks are decreasing in new stands of the bottomland forests (Clatterbuck and Hodges 1988, Guldin and Parks 1989, Hodges and Janzen 1987, Johnson 1979). Because of their slow early growth, newly established oak seedlings cannot compete successfully with sprouts and fast-growing pioneer species. Oak seedlings often grow and die back for 4 to 6 years after germination before they reach a stage of rapid height growth.

The slow growth of oak seedlings after germination is compounded by animal predation and overtopping by briars, sprouts, and early successional trees and brush (Gingrich 1979, Hannah 1987, Nix 1989, Watt 1979). However, if released, oak seedlings can respond and achieve a dominant position in the developing canopy (Nix 1989). Some type of release of oak seedlings after clearcutting is usually needed to have oaks in the dominant canopy in a reasonable period with guality, size, and grade (Beck 1970, Hannah 1987). Release should be accomplished 3 to 5 years after clearcutting to allow the seedling oaks to guickly catch up with the other woody species on the site (Beck 1970). Early release should allow the oaks to be a dominant component of the canopy by the time the stand is 25 years old. Early release may also shorten the normal oak rotation by 10 to 15 years for quality oak products (Clatterbuck and Hodges 1988, Clatterbuck and others 1985).

Release of crop trees with herbicides has been accomplished successfully for many years in the management of

pine stands (Bacon and Zedaker 1987, Clason 1978, Nelson and Cantrell 1990, Nelson and others 1985). Therefore, it should be possible to release oak seedlings with careful application of the same herbicide provided the oaks are somehow sheltered from the herbicides. During the winter of 1987 and 1988, two bottomland hardwood stands with a 40-percent oak component (basal area basis) were clearcut along the Congaree River in South Carolina. These stands apparently had a good seed crop prior to harvest, as from 3,000 to 5,000 oak seedlings per acre were established (Nix and Lafave 1993). When first examined, in the winter of 1989, these oak seedlings were overtopped by a dense canopy of blackberry, herbaceous composites, woody seedlings, and sprouts. This study was implemented to evaluate several herbicides and application methods for releasing overtopped oak seedlings developing in dense young clearcuts.

PROCEDURE

Forty-two plots approximately 18- to 22-feet wide and 25to 130-feet long were located along 2-foot wide transects cut through the vegetation of 3- and a 4-year-old clearcuts, approximately 1/2-mile apart in the first bottom of the Congaree River near Columbia, SC. The transects were placed where two or three cherrybark or Shumard oak seedlings in each of three height classes were found in an overtopped condition. The plots were placed at least 15 feet apart to reduce the chance of one herbicide treatment influencing any adjoining treatment. Each plot contained six to nine seedlings located along either side of the transect. Seedlings were chosen within approximately 7 to 9 feet of the transect, but at least 2 feet away so that the cut transect did not affect the seedlings' competitive status. The seedlings were selected to be approximately 6+ feet apart to accommodate the directed herbicide treatments. The length of the plot was determined by the distribution of the seedlings and the width of the plot was determined by the overspray width and the spray apparatus boom height. The

¹ Professor, Department of Forest Resources, Clemson University, Clemson SC 29634-1003.

Citation for proceedings: Connor, Kristina F., ed. 2004. Proceedings of the 12th biennial southern silvicultural research conference. Gen. Tech. Rep. SRS–71. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 594 p.

beginning and end of each plot center line was marked by a white, 4-foot, plastic pipe driven into the ground.

The treatment plots were blocked for replication according to preliminary estimates of total competition based on a height/distance index (Daniels 1976) on the plots (high, medium or low). Three replications of the treatments were arranged with each treatment plot randomly placed in each of the three blocks. The oak seedlings chosen for measurement were statistically similar in height on all plots. The preferred condition of the seedlings for release was overtopped, but at least 12 inches in height. Each plot contained at least two seedlings and, if possible, three seedlings in each of the three following height classes: 1-Foot Class (1 to 1.9 feet), 2-Foot Class (2 to 2.9 feet), 3-Foot Class (3 feet and taller). Each seedling chosen was marked by a wire flag and metal tag bearing an embossed coded number.

Preliminary treatments with a quick-browning herbicide in 20 percent increments of the recommended release rates for pines were applied to plots adjacent to the study area after full foliage development (midsummer) to refine the rates that were used in the treatments. The methods of treatment, herbicides, and rates which were applied in late July to early August were as follows:

- 1. Control-no treatment.
- 2. Directed (Streamline)—Application of 20 percent Garlon 4 (a triclopyr herbicide), 10 percent Cide-Kick II surfact-ant (JLB 1984), and 70 percent diesel fuel applied to the lower portion of all competing stems in a 3+ foot radius around the oak seedlings (Dow 1988, McLemore and Cain 1988, Yeiser and others 1989). This application was accomplished with a Solo piston backpack sprayer equipped with a D2 orifice disc tip for stream application.
- 3. Directed (Spot gun)-Application of Velpar L (a hexazinone herbicide) or Escort (a metsulfuron benzoate herbicide) applied in a grid pattern of six 1 mL spots placed approximately 3.5 feet from the oak seedlings to be released. The spots of Velpar L consisted of a mixture of Velpar L and water that was 50 percent of the manufacturer's product strength. The application mixture of Escort consisted of 1 ounce of Escort mixed in 2 gallons of water. Due to the fine texture of the soils on the study site, the manufacturer's rate of application for use on this texture of soil indicated that there would be a 1.5- to 2-foot non-toxic zone around the treated oak seedlings (Du Pont 1987, 1988; Rachal and others 1988). These applications were accomplished with a standard spot gun applicator calibrated at 1 mL per spot application.
- 4. Broadcast (Overspray)—Simulated aerial broadcast, an overspray applied with a boom apparatus, applied at 1/2 and full label-recommended application rate of commercial herbicides for releasing pines with 3.2 ounces per acre of TimberSurf 90 surfactant (0.25 percent v/v commonly recommended rate in release of pines) (Timberland ca.1989) at 10 gallons total mix per acre (Dow 1988) for each of the herbicides in table 1. The simulated aerial broadcast spray was applied with a carbon dioxide pressurized backpack apparatus with a

Table 1—Simulated aerial broadcast applications (oversprays) that were used at 100 and 50 percent of standard pine release rates for oak release in 3-and 4-year-old clearcut bottomland hardwood stands in South Carolina

	Standard pine
Herbicide	release rate
Garlon 4	1.5 ^a
Escort	1 ^b
Accord	1.5 ^a
Velpar L	3 ^a
Arsenal AC/Escort	8 / 0.25 ^{b c}

^a Quarts per acre.

^b Ounces per acre.

^c The rate of 0.25 ounces per acre of Escort remains the same for both the 100 and the 50 percent rate plots.

boom extended well above the vegetation (16 feet above ground) and equipped with a single horizontally aligned flood jet nozzle. The actual area treated was calculated by using the length of the plot, the boom height, and the nominal spray width within which competing vegetation would be uniformly treated (18 to 22 feet).

The spray time, spray pressure, boom height, and amount of herbicide to be used were predetermined. Measurements of the sprav time and pressure were recorded for each plot so the actual rate of application could be calculated. To determine effectiveness of the release treatments after two years, herbaceous competition, woody competition, total competition, seedling height, and ground line diameter (GLD) were measured before and after treatment. Data were analyzed using a complete random block design and PC-SAS, Anova procedure. Means were tested at the 0.01 level of probability. The results of the initial reduction of competition in this study were reported in an earlier paper (Thompson and Nix 1993). For the purpose of this paper which reports 10 year response to the herbicide treatments only diameter at 4.5 feet, total height, and total number of at least codominant red oaks per plot and acre were measured. Water and willow oaks (Q. nigra and Q. phellos) were included as desirable oak crop trees along with the cherrybark and Shumard oaks previously measured.

RESULTS AND DISCUSSION

The effectiveness of any crop tree release treatment can be judged early by the subsequent tree height, diameter growth, and presence of sufficient numbers of future crop stems in the dominant canopy. In this study, oak seedling numbers, height and diameter growth 2 and 10 years after treatment were the major criteria for evaluating the effectiveness of the herbicide release treatments. After 10 full growing seasons, 4 herbicide treatments resulted in a significant increase in diameter growth of released oak seedlings (table 2). Overtopped seedlings released by the

Treatment	Tree per	Tree per	TPA std.		D.b.h. std.
method	plot	acre	dev.	D.b.h.	dev.
	nur	nber		inch	
AVL	4.0	170	118	3.04	0.49
AVH	8.3	365	130	3.54	0.24
AArEL	5.0	219	46	3.92	0.27 ^a
AArEH	8.0	321	138	3.55	0.84
AAcH	11.7	537	223	2.86	0.29
AAcL	8.7	356	278	3.75	0.74
Control	4.7	221	171	2.45	0.04
SE	6.0	220	32	2.93	0.17
AEL	6.3	272	173	2.81	0.16
AEH	5.7	242	89	4.05	0.51 ^a
AGL	7.0	266	56	4.23	0.62 ^a
AGH	8.7	363	361	3.98	0.12 ^a
SV	7.7	336	45	3.74	1.50
SG	9.3	391	198	3.87	1.17

Table 2—Ten year response of 4-year-old red oak seedlings to herbicide release in two clearcuts in Red River bottomland hardwoods in South Carolina

TPA = tree per acre; A = at the beginning denotes aerial application; V = Velpar; Ar = Arsenal; E = Escort; Ac = Accord; G = Garlon 4; H = standard pine release rates; L = one-half rates; SE = spot applied Escort; SV = spot applied Velpar; SG = streamlined applied Garlon 4.

^a Treatments that are significantly different from the control at the 0.01 level of *p*.

aerial broadcast application of the low rate of Garlon, the high rate of Garlon, the high rate of Escort, and the low rate of Arsenal plus Escort were significantly larger in diameter than the untreated seedlings. Numbers of at least codominant saplings, although satisfactory for future stocking at 250 per acre, did not differ from those of the control, 221 per acre. Height of released oak saplings averaged 35 feet but did not differ statistically from the 28 feet height of the control saplings.

One possible reason for the lack of significant height growth response is that the released oak saplings appeared, by casual observation, to have larger and fuller crowns than the control saplings or oak saplings that had less reduction in competition (Thompson and Nix 1993). Another reason for less height growth by released seedlings may be the vine entanglement or lack of close trainer trees in the more extreme directed treatments. The directed treatments, Velpar and Escort spot gun and Garlon streamline application were very effective where splash or flashback did not kill the released oak saplings. However, such treatments were totally impractical in the dense tangled young clearcuts. Again, from casual observation, the oak seedlings that were released enough by the directed herbicide sprays to obtain full sunlight, but that had close (within 3 feet), taller saplings, seemed to achieve more height growth than other oak seedlings.

Vines, particularly the grape (*Vitus* spp.), and honeysuckle (*Japonicum* spp.), were especially onerous; they suppressed many of the released seedlings which eventually died or lost dominance as saplings. In addition wherever vigorous lob-lolly pine (*Pinus taeda*) saplings and sweetgum (*Liquidambar*

styraciflua) sprouts occurred in any of the treatment plots they totally dominated the canopy and suppressed even the released oak seedlings. Furthermore, it is suspected that as many as 20 percent of the taller oak seedlings, greater than 4 to 6 feet, were killed or inhibited by the herbicide overspray treatments as they projected through the sheltering main canopy at 4 years of age (Thompson and Nix 1993).

CONCLUSIONS

It is possible to release oak seedlings with careful application of selected herbicides in young (3- to 4-year-old) clearcuts without serious losses, especially if the oak seedlings are at least several feet tall and in an overtopped, sheltered condition. This overtopping, which is very common in such clearcuts, seems to shield the seedlings from a lethal amount of the broadcast herbicides. Although only 4 treatments induced a significant increase in seedling diameter 10 years after treatment, most of the 13 treatments appeared to increase diameter growth and the presence of desirable oaks in the dominant canopy. It may be too early to ascertain the full effect of this release at 14 years of age, but some of these treatments appear to have potential for establishing more oaks in the main canopy at an earlier age, thereby increasing the oak component and basal area of the young stand. It is apparent that vines must be controlled on these better bottomland sites at some time during this critical developmental period, or many of the released oak seedlings will succumb to suppression.

These are 10 year results and because of the oak developmental pattern, the codominant oak saplings need perhaps 10 more growing seasons before the full treatment effects can be evaluated. Economic analysis of the treatments will better assess the full practicality of such herbicide release of young overtopped oaks in recently clearcut bottomland hardwood stands.

ACKNOWLEDGMENTS

The author gratefully acknowledges the assistance of former graduate student John F. Thompson who did most of the work in establishing and documenting this study in the initial 2 year measurements. Able assistance was also provided by former graduate student Charles Carlisle and current undergraduate assistant Elijah Blue Hozey in the subsequent 10 years of work. We are all grateful to the Millaree Hunt Club and Mr. Marion Burnsides, chairman, for the hospitality shown us and to the property owners, the Biedler Family, and to Angus Lafaye, consulting forester of Milliken Forestry who first called the bountiful oak regeneration to our attention.

LITERATURE CITED

- Bacon, C.G.; Zedaker, S.M. 1987. Third-year growth response of loblolly pine to eight levels of competition control. Southern Journal of Applied Forestry. 11: 91-95.
- Beck, D.E. 1970. Effect of competition on survival and height growth of red oak seedlings. Res. Pap. SE-56. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 7 p.

Clason, T.R. 1978. Removal of hardwood vegetation increases growth and yield of young loblolly pine stands. Southern Journal of Applied Forestry. 2: 96-97.

- Clatterbuck, W.K.; Hodges, J.D. 1988. Development of cherrybark oak and sweetgum in mixed even-aged bottomland stands in central Mississippi, U.SA. Canadian Journal of Forest Research. 18: 12-18.
- Clatterbuck, W.K.; Hodges, J.D.; Burkhardt, E.C. 1985. Cherrybark oak development in natural mixed oak-sweetgum stands -Preliminary results. In: Shoulders, E., ed. Proceedings of the third biennial southern silvicultural research conference. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 438-444.
- Daniels, R.F. 1976. Competition indices and their correlation with annual loblolly pine tree growth. Forest Science. 22: 454-456.
- Dow. 1988. Garlon 4 herbicide: Specimen label. The Dow Chemical Company. 38322-L6.
- Du Pont. 1987. Velpar[®] L herbicide: Specimen label. E.I. Du Pont De Nemours and Co., Inc. Ag-2508.
- Du Pont. 1988. Escort[®] herbicide: Specimen label. E.I. Du Font De Nemours and Co., Inc.. AG-2989.
- Gingrich, S.F. 1979. Stocking levels and regeneration potential. In: Proceedings: Regenerating oaks in the upland hardwood forests. John S. Wright forestry workshop. West Lafayette, IN: Purdue University: 27-29.
- Guldin, J.M.; Parks, T. 1989. Development of cherrybark oak in an uneven-aged stand in west Tennessee. In: Miller, J.H., comp. Proceedings of the fifth biennial southern silvicultural research conference. Gen. Tech. Rep. SO-74. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 327-331

- Hannah, P.R. 1987. Regeneration methods for oaks. Northern Journal of Applied Forestry. 4: 97-101.
- Hodges, J.D.; Janzen, G. 1987. Studies on the biology of cherrybark oak: Recommendations for regeneration. In: Phillips, D.R., ed.
 Proceedings of the fourth biennial southern silvicultural research conference. Gen. Tech. Rep. SE-42. Asheville, NC: U.S.
 Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 133-139.
- JLB. 1984. Cide-Kick II spray adjuvant for pesticides: Specimen label. JLB International Chemical Inc. Vero Beach, FL.
- Johnson, R.L. 1979. Adequate oak regeneration—a problem without a solution? In: Proceedings, seventh annual hardwood symposium. Hardwood Research Council: 59-65.
- Johnson, R.L. 1984. Direct-seeded cherrybark and Shumard oaks battle natural regeneration through 10 years. Southern Journal of Applied Forestry. 8: 266-230.
- McLemore, B.F.; Cain, M.D. 1988. A test of basal sprays for controlling hardwood brush and trees. In: Proceedings of the southern weed science society, 41st annual meeting. 41: 180-186.
- Miller, H.A.; Lamb, S.H. 1985. Oaks of North America. Happy Camp, CA: Naturegraph Publishers, Inc. 327 p.
- Nelson, L.R.; Cantrell, R.L. 1990. Herbicide prescription manual for southern pine management. EC 659. Clemson, SC: Clemson University, Cooperative Extension Service and U.S. Department of Agriculture. 53 p.
- Nelson, L.R.; Zutter, B.R.; Gjerstad, D.H. 1985. Planted longleaf pine seedlings respond to herbaceous weed control using herbicides. Southern Journal of Applied Forestry. 9: 236-240.
- Nix, L.E. 1989. Early release of bottomland oak enrichment plantings appears promising in South Carolina. In: Miller, J.H., comp. Proceedings of the fifth biennial southern silvicultural research conference. New Orleans, LA: Gen. Tech. Rep. SO-74. U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 378-383.
- Nix, L.E.; Lafaye, A. 1993. Successful regeneration of quality oaks in a red river bottomland hardwood stand in South Carolina. In: Brissette, J.C., ed. Proceedings of the seventh biennial southern silvicultural research conference. New Orleans, LA: Gen. Tech. Rep. SO-93. U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 81-85.
- Rachal, J.M.; Karr, B.L.; Janzen, G.C. 1988. Spot applications of hexazinone for pine release-fifth year results. In: Proceedings of the southern weed science society. 41st annual meeting. 41: 163-172.
- Thompson, J.F., Jr.; Nix, L.N. 1993. Early release of naturallyregenerated cherrybark and Shumard oak seedlings with herbicides. In: Brissette, J.C., ed. Proceedings of the seventh biennial southern silvicultural research conference. New Orleans, LA: Gen. Tech. Rep. SO-93. U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 445-451.
- Timberland. ca.1989. Timbersurf 90: Specimen label. Monticello, AR: Timberland Enterprises, Inc.
- Watt, R.F. 1979. The need for advance regeneration in oak stands. In: Proceedings: Regenerating oaks in the upland hardwood forests. John S. Wright Forestry workshop. West Lafayette, IN: Purdue University: 11-17.
- Yeiser, J.L.; Reed, D.J.; Kidd, F.A. 1989. Streamlining with triclopyr and fluroxypyr for hardwood control. In: Proceedings of the southern weed science society. 42nd Annual Meeting. 42: 197-201.

Great Trinity Forest Management Plan

HARDWOOD SILVICULTURE

Oak Regeneration: Serious Problems Practical Recommendations

(USDA Forest Service Southeastern Forest Experiment Station General Technical Report SE - 84)

United States Department of Agriculture

Forest Service



Southeastern Forest Experiment Station

General Technical Report SE-84

Oak Regeneration:

Serious Problems Practical Recommendations



SYMPOSIUM PROCEEDINGS Knoxville, Tennessee September 8-10, 1992

- Loftis, David; McGee, Charles E. eds. 1993. Oak Regeneration: Serious problems, practical recommendations. Symposium Proceedings; 1992 September 8-10; Knoxville, Tennessee. Presented by the Center for Oak Studies. Gen. Tech. Rep. SE-84. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 319 pp.
- Twenty six papers dealing with the problems and opportunities associated with artificial and natural regeneration of the oak resources are presented. Subject matter, titles of papers, and authors were carefully selected to provide the best available coverage of the state-of-the-art of oak regeneration.

Remarks about pesticides appear in some of the papers contained in these Proceedings. Publication of these statements does not constitute endorsement or recommendation of them by any of the conference sponsors, nor does it imply that uses discussed have been registered. Use of most pesticides is regulated by State and Federal law, and applicable regulations must be obtained from appropriate regulatory agencies.

Caution: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish and other wildlife—if they are not handled and applied properly. Use all pesticides selectively and carefully. Follow recommended practices given on the label for use and disposal of pesticides and pesticide containers.

The use trade or company names of products or services in the Proceedings is for the benefit of the reader. Such use does not constitute an endorsement or approval of any service or product by the symposium sponsors to the exclusion of others that may be suitable.

Oak Regeneration:

Serious Problems, Practical Recommendations

Symposium Proceedings

Knoxville, Tennessee September 8-10, 1992

Editors David L. Loftis and Charles E. McGee

Sponsored by USDA Forest Service: Southern Region, Southeastern Forest Experiment Station, and Southern Forest Experiment Station

> Presented by: Center for Oak Studies Department of Forestry, Wildlife, and Fisheries University of Tennessee, Knoxville

Published by: Southeastern Forest Experiment Station, Asheville, N.C.

Page 55 of 404

Preface

These Proceedings are in partial fulfillment of a Technology Transfer Plan on oak agreed to by a number of parties in 1988. The plan called for a symposium on oak regeneration. Oral presentations based upon the papers in these proceedings were delivered on September 9th and 10th, 1992, in Knoxville, Tennessee.

A Core Team was selected to administer and implement the Technology Transfer Plan, and this Core Team served as the Steering Committee for the symposium. An outline for the symposium on oak regeneration was developed and approved by the Core Team, which then approved the subject for each paper and decided upon the most appropriate author to write and present the paper. The primary objective was to carefully structure a program would problems that address the and opportunities associated with oak regeneration. A secondary objective was to evaluate the procedures used in this effort as means of achieving technology transfer.

Each of the papers in the Proceedings received technical and editorial review; all reviewers' comments were made in the form of suggestions, however. Therefore, the content and accuracy of each paper is the responsibility of the author. Overall, the Core Team feels that these papers accurately reflect the state of the art for oak regeneration today. The Team suggests that each reader take into account that some conclusions and recommendations are reached and made in a very dynamic environment. It is expected that this symposium fills the need for "results now."

The Core Team wishes to thank the sponsors, authors, and reviewers. The Team especially appreciates the moderators who not only added their own expertise and credibility to the program but did an excellent job keeping the demanding schedule intact. Moderators included Jack Pitcher, Hardwood Research Council; Bob Rogers, University of Wisconsin; Bill Mahalak, Michigan Department of Natural Resources; Tony Parks, Anderson Tully Company; Gary Schneider, University of Tennessee; Randy Rousseau, Westvaco Corporation; and Charles E. McGee, Center for Oak Studies.

The Core Team especially thanks Bill Hamilton (USFS-Retired) and Tami Steppleton (USFS), for their dedicated, persistent efforts to design, assemble, and format these Proceedings.

Core Team

Presentation Committee

Dan Sims, Co-chair USDA Forest Service

Ed Buckner, Co-chair University of Tennessee

Kim Coder University of Georgia

William Gardner North Carolina State University

Planning Committee

John Toliver, Chairman USDA Forest Service

Harvey Kennedy USDA Forest Service

David Graney USDA Forest Service

Tom Lloyd USDA Forest Service

Wayne Clatterbuck Tennessee Division of Forestry

Editorial and Proceedings Committee

David Loftis, Chairman USDA Forest Service

John Hodges Mississippi State University

Dave Smith Virginia Polytechnic Institute

Paul Johnson USDA Forest Service

Symposium Coordinator: Charles E. McGee-Center for Oak Studies

iv

Contents

THE OAK REGENERATION PROBLEM

An Historical Perspective of Oak Regeneration
Causes of the Oak Regeneration Problem 14 Craig Lorimer
Oak Regeneration: The Scope of the Problem 40 David Smith
THE BIOLOGY OF OAK REGENERATION
Ecology and Physiology of Oak Regeneration 54 John Hodges and Emile Gardiner
The Role of Fire in Oak Regeneration 66 David H. Van Lear and Janet M. Watt
Flowering and Oak Regeneration
Acorns and Oak Regeneration
Insects and Diseases Affecting Oak Regeneration Success
Sources of Oak Reproduction
PREDICTING AND MODELING OAK REGENERATION
Predicting Oak Regeneration—State of the Art 134 David Loftis
Predicting Quantity and Quality of Regeneration in the Uplands
Prediction of Oak Regeneration in Bottomland Forests
Decision-Making for Natural Regeneration in the Northern Forest Ecosystem

NATURAL REGENERATION OF OAKS

Regenerating Oaks in the Central States	174
Regenerating Oaks in the Bottomlands	184
Relative Growth of Oaks and Pines in Natural Mixtures on Intermediate to Xeric Piedmont Sites Tom Lloyd and Tom Waldrop	196
Regenerating Northern Red Oak on High-Quality Sites in the Southern Appalachians	202
Regenerating Oaks in the Central Appalachians	211
ARTIFICIAL REGENERATION OF OAKS	
A Historical Perspective of Planting and Seeding Oaks Phil Pope	224
Artificial Regeneration of Bottomland Oaks Harvey Kennedy	241
Artificial Regeneration of Oaks in the Uplands	250
Oak Plantation Establishment Using Mechanical, Burning, and Herbicide Treatments	264
Collection and Care of Acorns	290
The Use of Tree Shelters and Underplanting for Oak Regeneration	298
SUMMARY PAPERS	
Oak Regeneration—Where We Go From Here? Bob Kellison	308
Oak Regeneration: A Summary Charles McGee and David Loftis	316

The Oak Regeneration Problem

Page 61 of 404

An Historical Perspective of Oak Regeneration

F. Bryan Clark, USDA Forest Service, Retired, Annandale, VA 22003

ABSTRACT Concerns about oak management in the middle ages led to forest ordinances in France designed to ensure oak establishment. Oak was an early export from the American colonies because it was scarce and expensive in England. Probably the first government sponsored forest research in the United States was attempts in Florida in the early 1800's to establish live oak (Quercus virginiana) for shipbuilding. With the advent of technical forestry in this country there were references in early papers and textbooks to oak regeneration and the shelterwood method. This interest was primarily for academic study and followed European descriptions, but some of the earliest oak regeneration research related to concerns about areas without advance reproduction.

> In the late 1930's understory oaks were described in relation to past treatment. In the 1950's and 1960's researchers and ecologists began to quantify oak reproduction under various stand conditions on different sites. Mostly they found lots of understory oaks, but there were exceptions. Researchers thought these exceptions might be important and started to look more carefully at the response of oaks to harvest cutting. Even-aged cuttings quickly showed that on good oak sites without substantial numbers of vigorous advance oak the stands of the future would be quite different. The new stands generally contained less oak. At the same time observers noted that oak types were changing in some areas. More recently, comparisons of repeat forest surveys show a decline of oak types in some states. Increased prices for oak timber suggest that accelerated harvesting is adding to the ecological changes taking place.

> We find ourselves with a growing knowledge about oak ecology and silviculture but without a convincing history of being able to prescribe with certainty ways to increase the amount of oak in new stands. Considering the tremendous value of oak forests there is an urgency to improve oak management and to give high priority to long-term research.

INTRODUCTION

Is oak regeneration really a problem? Experts believe that on many sites now occupied by oak, regeneration will become increasingly difficult. How can that be? Oaks are the most prevalent forest types in the United States and dominate much of the East's landscape. Volume growth exceeds commercial removals. It is even more confusing to remember that the present stands were born out of very harsh efforts to either convert forest lands to agriculture through fire and grazing or to

mine them of timber with little or no concern for future crops. But it was these very treatments plus the loss of chestnut (*Castanea dentata*) in the Appalachian region that favored oak. Some recommended practices of the last 50 years favor other species at the expense of oak, especially on moist sites where oak quality is best. Paradoxically, we find oak on some of the areas we thought were mismanaged in the past. I believe there is a pressing need for practical alternatives to create conditions to favor oaks on the right sites without detriment to the land or other forest resources.

The very physiologic and genetic characteristics that make oak difficult to establish are the traits that have sustained it through centuries of insults. We have been too impatient in dealing with oak, but there is emerging a better understanding of oak culture. At the same time forest managers have joined users and researchers in common concerns about the future of the oak resource. To understand the complexities of oak regeneration, the status of current prescriptions, and the best direction for future research and development, the Symposium organizers decided to look at the past as prologue for this conference. I have drawn on selected literature along with my own recollections. Consequently, this paper is a collage of facts and personal experience.

To the first farmers of Europe thousands of years ago oaks and other trees were the enemy, something to be girdled with stone axes and burned to make way for crops and a better way of life. In contrast, native Americans living in eastern forests did not rely as much on agriculture and clearing. Their biggest impact on the oak forests related to fire they used or caused. In Europe, by Roman times, oak coppice management regimes developed to improve wood supplies. By the Middle Ages concerns about oak forests led to the French forest ordinances in the 13th Century that mandated practices to establish oak seedlings (Thirgood 1971). Have the French known for hundreds of years what we have been "discovering" over the past 30 or 40 years? Probably, but remember the first American foresters were European-trained and undoubtedly were familiar with oak practices and problems in France, Germany, and Great Britain. Oak was an early export of the American colonies because it was becoming scarce and expensive in Great Britain.

Early American textbooks reflected European silvicultural systems including shelterwoods and coppice. But, there were few references to oak management problems in American forest literature until after the 1930's. There are a number of plausible reasons for this lack of concern: (1) Historically much of the hardwood forest was an unwanted barrier to agricultural development; (2) After several hundred years of harsh treatment and neglect oak trees were still plentiful and there were no serious timber supply problems; (3) Protection and reestablishment were the primary early forestry concerns; and (4) The very practices of indiscriminate burning and overcutting that fostered the forest conservation movement favored oak. It was not so much that we ignored oak regeneration problems in the United States; we are following historical precedence of abundance. We wait to respond to a problem that has emerged over a long time until it has the potential to reduce future supplies. Considering the ecological, economic, and social forces involved we must realistically expect a decline in the oak forest types. However, it is certainly not "too late" to develop and extend the necessary technology to maintain healthy and useful stands throughout the range of oak.

CURRENT CONCERNS NOT NEW

Specific concerns for oak regeneration in the United States go back at least to the early 1800's. Perhaps the first government sponsored forest research in this country was to plant and tend live oak (*Quercus virginiana*) on public land in Florida (Hough 1878). At that time there was great concern about adequate supplies of oak timber for shipbuilding.

Early forestry authors Leffelman, Hawley, and Korstian recognized the kinds and importance of advance oak regeneration (Leffelman and Hawley 1925, Korstian 1927, Hawley 1946). Hawley's 5th edition text, *Practice of Silviculture*, described a successful 1902 shelterwood cutting in Connecticut that favored oak; so there must have been a shortage of advance oak in that stand. Hawley's first edition was printed in 1921 and likely followed European practice. Smith (1962) made a good review of oak regeneration shelterwoods in his silviculture textbook. He also suggested that the term shelterwood is better than the term clearcutting when advance reproduction is involved. A "one-cut" shelterwood for oak *required* that advance reproduction must be established before the final harvest. This excellent perspective is still the conventional wisdom for extensive management.

Liming and Johnson (1944) described oak reproduction in the Missouri Ozarks in 1933 when fire protection started on the new National Forests. Reproduction appeared to be sparse and in poor condition. But within a few years sprout stands emerged from existing roots. Seedling numbers also increased. The authors predicted that with time and protection the forests would improve. And so they have. The change has been dramatic even since 1949 when I first worked under Franklin Liming. As the Ozark forests grew so did the concerns that the next generation of forest stands might be different. A similar "sudden" appearance of oak advance regeneration took place in the late 1800's in southern Wisconsin with the cessation of wildfire as described by Curtis (1959). Crow (1988) provides an excellent historical review and bibliography on oak forests and savannas before European settlement. He cites numerous authors to show how fire and logging favored oak in many parts of the East. In the 1950's Scholz, Arend, Johnson, and others worked on oak regeneration problems as they developed in southern Wisconsin and adjacent Lake States areas.

Information was also accumulating in the Central States. Ecologists Potsger and Friesner (1934) compared a virgin southwestern Indiana oak-hickory forest with adjacent areas cut in the late 1800's and found succession to be a return to oak-hickory and not beech-maple. But from the species lists it is obvious that the sites were dry. Kuenzel and McGuire (1942) reported on clearcut plots in southern Indiana following a bumper crop of chestnut oak (Q. prinus) seed. Ten years later 97 percent of the new stand was from stump sprouts. This important information probably did not cause much of a stir among managers but it was picked up by researchers. Illinois plots clearcut in 1935 had good oak 27 years later due to desirable advance reproduction (Bey 1964). In the late 1940's Leon Minckler began a series of compartment studies on the Kaskaskia Experimental Forest in Illinois and showed how oak, especially white oak (Q. alba), grew and developed in group selection openings (Minckler 1989).

Studies of oak seed production in North Carolina and Georgia (Downs and McQuilken 1944) also showed the importance of advance reproduction in that area. They suggested shelterwoods were best to regenerate oaks and thought that small group selection openings might also work. By 1958 Merz and Boyce showed that

in southeast Ohio the amount of oak in the new stands was directly related to the amount of reproduction established before cutting. In the mid-fifties Ivan Sander and I remeasured plots on the Berea College Forest in Kentucky. The plots were cut in 1923 by the old Appalachian Experiment Station. Sugar maple (Acer saccharum) dominated the understory but red oak (Q. rubra) dominated the overstory. In 1923 sugar maple was assumed to be a good species for these relatively thin soils. But the more than 30 years growth on these plots showed that red oak was a much better crop than sugar maple in both quality and volume growth (Sander and Williamson 1957). These results made an early impression on me because a maple understory was developing under many mixed oak stands in the Midwest as fire control took hold.

Also, in the mid-fifties I made my first visit to West Virginia and was surprised to hear complaints about "too much" yellow-poplar (*Liriodendron tulipifera*) showing up in the forest landscape even though yellow-poplar is an excellent species. The concerns were about future oak timber and mast supplies. Then, as clearcutting became more prevalent we began to see more yellow-poplar along with red maple (*A. rubrum*). Later, Beck (1988) gave an excellent synthesis and forecast for continued increases of yellow-poplar and other fast growing species at the expense of oaks unless special efforts are made to favor oak establishment and early growth.

In a classic West Virginia study Carvell and Tryon (1961) stated: "The deficiency of oak regeneration beneath mature oak stands is of grave concern to the forest manager since oak regeneration is virtually impossible to obtain quickly." They too found that the composition of the new stand depended largely on the composition of the understory before harvest. However, they counted more oak seedlings than expected in a survey of 59 areas in eight counties. The greater the past disturbance from fire, logging, and grazing, the more oak they found.

Studies in West Virginia (Weitzman and Trimble 1957, Carvell and Tryon 1961) and from other locations showed that moist sites are more difficult to regenerate to oak than are dry sites. The basic problem on moist sites including bottomlands is understory competition.

In 1960 the Division of Forest Management Research of the old Central States Forest Experiment Station under A.G. Chapman prepared a comprehensive problem analysis, Guidelines for Forest Management Research in the Central It was prepared by committees and represented the Hardwood Region. conventional wisdom, but not necessarily agreement, of a diverse group of hardwood researchers. It was never published. Some of the conclusions related to oaks and cutting practices 32 years ago were: (1) Reproduction follows all kinds of harvests; (2) Even-age silviculture was better for central hardwoods than selection silviculture; (3) Nearly all clearcutting trials had adequate oak; (4) There was strong evidence that oak consistently followed partial cutting; (5) Major objections to clearcutting related to concerns about sprout stand quality and increased competition of undesirable species; and (6) The quality and importance of advance reproduction was in a state of confusion because of insufficient data. When this analysis was written in the early 1960's there were a large number of active Forest Service oak silviculture studies. Many were new and inconclusive. Highest priority for new research was to find out how to ensure adequate, vigorous regeneration for upland oak. A few years later Forest Service oak research in

Ohio, Kentucky, and Indiana was discontinued. Later the same fate befell most silvicultural research in Illinois and Iowa.

Natural hardwood regeneration research dating from the 1950's was summarized by Sander and Clark (1971). More than 200 cutting plots in Ohio, Kentucky, Indiana, and Illinois were analyzed. This publication provided quantitative data for some of the tenets of central hardwood regeneration at that time. Cutting method had little effect on the amount of regeneration except for yellow-poplar; very few oaks were established after the cuts. The importance of advance oak was stressed, but there were no specific concerns in the text about a lack of advance oaks. However, the statistical tables showed that the Indiana plots—all in one location on the Hoosier National Forest—had a lot fewer advance oak than in other states.

In the late 1960's and early 1970's interests in oak regeneration and management grew as the controversy over harvest cutting methods developed. The forestry community responded to this interest by organizing the first Oak Symposium in 1971 at Morgantown, West Virginia. This followed a special session on hardwood silviculture at the 1968 Society of American Foresters Convention in Philadelphia. I gave papers on regeneration at both meetings at the height of my "oak missionary days" (Clark 1970 and Clark and Watt 1971). Although no longer a research doer, I was part of the movement to get more research results into practice. At the 1971 Oak Symposium, Richard Watt and I synthesized our regeneration recommendations from the work of several researchers including our own. We stated that most maturing stands had enough advance oak but some did not. We had no data on the importance of this problem. Our synthesis has held up fairly well. Since then a number of technical issues are now better resolved through continuing research and new information. Yet, in those 20 years since the first Symposium, progress and proof seem slow. It is the nature of oak.

SOLVING THE Many attempts have been made to plant oaks but successful plantations were rare in the past. Midwest forest tree nurseries have grown and shipped oaks since the PROBLEMS 1930's. I personally checked a number of old oak planting sites from the records of the Hoosier National Forest; all sites were old fields, all were failures. In the late 1950's Bob Williams and I included oaks in our hardwood nursery practices and planting research in Indiana. Most of our time was spent on black walnut (Juglans nigra). We had good success with walnut, some encouraging results with red oak, and poor results with white oak. We were convinced that site selection, stock quality, and competition control were very important for all hardwoods. Following a series of studies started in the 1960's, Johnson (1985) reported some success with oak underplanting in Missouri. More recent oak planting studies have produced positive results on upland sites in the Central and Lake States, the Mid-South, and the Northeast. There are also reports of successful planting on bottomland sites. The goal is to substitute planted oak on problem sites without natural oak regeneration. The outlook for operational planting is now much better, but risks and costs are still high.

Early researchers recommended that advance oak must be big or vigorous enough to compete in the new stands, but specifics on size were not available until the 1970's. Perhaps the classic study of the response of different kinds and sizes of oak regeneration was on the Kaskaskia Experimental Forest. Sander (1972) concluded from 12-year data that only advance oak 4.5 ft. or taller could compete in the new stand. He suggested it would be better to have 6-to-8 ft. regeneration before final cutting. Some of the more esoteric results of that research deserve careful restudy by serious oak researchers and managers. Only the regeneration types with older, well-developed root systems grew to the recommended height during 12 years in partial cut stands with only 29 percent overstory stocking. This is a bit chilling when considering shelterwood schedules in areas with scarce advance oak.

Sander, Johnson, and Watt (1976) developed specific guidelines to evaluate the adequacy of advance oak. This work continues to be refined but applies mainly to Missouri. Carvell (1979) did some original work on the importance of recognizing not only the size but also the vigor of advance oak. He too stressed the importance of adequate advance oak based on his earlier work and suggested how to make preharvest assessments (Carvell 1988). Loftis (1988) used an approach similar to Sander and associates to determine the oak regeneration potential for areas in the Appalachians.

Most authors stressed that it is essential to have large numbers of well-developed oaks in the understory prior to the final harvest. In sharp contrast, Johnson and others (1989) reported on a mesic site clearcut in southwestern Wisconsin that was successfully reproduced with new red oak seedlings. In this general area oak is being replaced through successional pressure of more tolerant species. In this case study, competition was greatly reduced by treating the understory with herbicides 2 successive years, and then removing the overstory after a good seedfall. The authors also cited examples of both success and failures with fewer and smaller advance red oaks than usually recommended. Johnson and others (1989) suggested that competition control with herbicides may substitute for the long regeneration period suggested in the literature. But they caution that a shelterwood is safer than a clearcut and that understory treatments may vary among ecosystems. Understory control has been suggested for many years as a potential solution to competition problems, but we still need specific prescriptions supported by practical demonstrations for a variety of problem areas.

RECENT HISTORY— PROBLEMS GROW

In 23 selectively cut private woodlands in southern Indiana, 95 percent of the openings created were too small to stimulate advance oak growth (Callahan and Fischer 1982). Oak was a substantial overstory component in most of the woodlands studied, but oak reproduction more than 6 ft. tall were found on only 2.4 percent of the transects. While the 23 areas appeared to be productive, the authors found them to be seriously understocked with desirable species. Callahan and Fisher concluded: "The present prospects appear dim that many oak trees will grow in future stands on upland hardwood sites."

Responding to concerns about cutting practices on the Hoosier National Forest, George and Fischer (1989) reported on an intensive regeneration survey on five 17year-old clearcuts in southern Indiana. Their data showed that the new stands have more yellow-poplar and less oak than the stands harvested. They point out that long-term development trends remain unanswered. One important question is will the more mesic species fade with time and drought as reported by Hilt (1985) in southeast Ohio? Interestingly, the yellow-poplar/oak trends in a large number of commercial clearcuts discussed by George and Fischer (1989) were consistent with data found in Sander's and Clark's 1971 report for southern Indiana plots and the increased incidence of yellow-poplar elsewhere.

With the advent of fire and grazing protection, oak stands in Iowa and Missouri are in various stages of conversion to sugar maple and other hardwoods, according to Countryman and Miller (1989). Recently, McGee (1989) found that an old stand of mixed hardwood with a substantial oak component is suddenly experiencing rapid overstory decline. In spite of the fact that this middle Tennessee old-growth on an excellent site had survived for decades, there is little chance that oak will be in the new stand under natural succession. Some eastern stands with severe gypsy moth mortality are not returning to oak but are now occupied by such low-value species as red maple that dominated the understory when the oak overstory declined. The potential loss of the oak type through gypsy moth is devastating.

Tracing the historical development of 46 red oak stands in northern Wisconsin provides strong evidence that these stands replaced other species and associations following past heavy cuts and fire (Nowacki, Abrams, and Lorimer 1990). The authors suggest that northern red oak in the study area may be limited to one generation and will be replaced by the tolerant red and sugar maples on all but the driest sites. The authors provide a brief overview of the mostly 1980's ecological literature to support the growing belief "that a high proportion of stands are on the threshold of a dramatic change in structure and composition." They too suggest that dry sites may be an exception to the ecological "instability" of eastern forest oak stands. These are strong and sobering opinions shared by a growing number of ecologists and foresters. Questions of oak stand dynamics have been around for a long time but evidence accumulates with the time it takes for ecological trends to manifest.

Crow (1991) concludes that in the Upper Midwest landscapes "future forests will differ in composition and structure from past and present forests." Crow (1988) also refers to the present abundance of oak in eastern forests as "an artifact of disturbance regimes that are no longer common." Spencer and Kingsley (1991) give us definitive insight to oak resource problems from the perspective of forest inventory type changes. Their analysis covered the seven-state Upper Midwest from Indiana to Minnesota. They compared the latest oak forest inventories with previous inventories. The average time between inventories was 15 years. During that "average" period oak types lost 6.5 percent in area but increased in total volume. Four of the seven states decreased in oak area. The increased area in the other three states was due primarily to a classification anomaly of wooded pastures shifting to timberland between inventories. Indiana had the biggest oak loss, 36 percent for the 19 years between surveys. That was an average annual conversion of 1.9 percent of oak types into the more mesic maple-beech type. This report also shows that only 17 percent of the oak types are in seedling-sapling stands, and is further proof that many oak stands are not replacing themselves. Spencer and Kingsley suggest that red oak is being overcut in Illinois and Indiana. Research economists in the Lake States found that the value of red oak lumber and logs has been increasing 6 to 8 percent a year above the inflation rate since the 1970's (North Central Forest Experiment Station 1991). Stumpage prices for both domestic and export logs are up dramatically, suggesting a growing scarcity of

"available" high quality trees. History repeats itself, only this time it is not a shortage of oak for shipbuilding and fuel.

Nyland (1992) has expressed strong concerns about potentially serious consequences of selective and diameter limit cuts in eastern hardwoods. He suggests that exploitive harvests following the development of export markets "portends a long-term conversion in composition that promises lower market values in the future, especially among oak communities." Ralph Nyland, one of the countries leading hardwood silviculturists, calls on practicing foresters to be more aggressive in promoting good forestry to stop the second great exploitation of the eastern forests.

Spencer and Kingsley's (1991) analysis of what is happening in the Upper Midwest is a dramatic picture of recent "historical" trends in oak ecology. Considering the Indiana data, the situation is cause for serious concern for those who want to maintain oak on good sites. Are we really witnessing successional changes taking place at a rapid rate? Yes, *but* it is not all "natural" succession. Selectively harvesting oak is probably most responsible for type conversion changes from one inventory to the next. Inventory forest types relate to the overstory, so removing the oak and leaving most of the other species automatically changes the type. What is left, the residual type, does not bode well for regenerating a new oak stand. Other harvest methods, such as clearcutting without vigorous advance oak regeneration, can also result in type conversion.

AND WHAT OF THE FUTURE?

Practicing field foresters in the Northeast ranked oak regeneration guidelines as the eighth most important research need out of 46 research problems (Broderick and others 1991). The authors stressed that the 660 respondents were well qualified to give an accurate picture of research needs. Yet the authors point out that "the literature does not lack for such information" and cite work done in the Central States and Missouri. They suggest that foresters may be "unaware of existing research or they are not satisfied with it" and that there may be problems in technology transfer. All of the above is true. But we need to understand that existing oak regeneration guides are based on data from limited areas and should only be extended with caution, validation, and perhaps additional research and testing to account for regional ecological differences and past treatments. In my experience, the call for more research is often based on an inadequate knowledge of information that already exists. But there are problems with the transfer of oak technology, and my definition of oak technology includes information for understanding oak ecology and ecosystems with oak. Too few managers and researchers are well founded in oak ecology; I include my own generation as part of the problem. Oak silviculture and management is far too complex to relegate to "standard" prescriptions.

Just as the present stands changed over time, so will the next rotation change under different treatments and environmental influences. Unfortunately, a lot of past long-term oak research has been discontinued due to costs and changing priorities. With new technology to handle data it is not too late to salvage some of this work, especially on areas with good understory records. A new look at some of the older studies may answer a very important question: Will oak outlast some of the more

Page 69 of 404

mesic species? Recently there has been a call for more financial support for longterm, basic ecological research. What could be more "basic" than understanding what is happening to the largest forest types in the United States surrounding a majority of the country's population? Like forester-turned-pathologist-turned Nobel Peace Prize winner, Norman Borlaug, I believe: "Our research must be good, but it must be good for something."

PUTTING HISTORY TO WORK In the past 10 years there have been at least six major conferences, from Pennsylvania to Minnesota and now Tennessee, that have addressed oak problems and solutions. From an historical perspective this unprecedented interest and concern for oak is both encouraging and further evidence that there are serious management problems over a broad area. Published prescriptions, guides, and texts are now more definitive but with room for improvement as new information emerges. The vast extent of the oak forests with their inherent and acquired variations makes them very difficult to know and manage. With that difficulty comes the challenge and excitement of working with a resource that has provided for the needs of both ancient and modern mankind from food to esthetics.

There is no easy way to summarize the history of oak regeneration problems except to say that it is lengthy, complex, paradoxical, and often perplexing. But do not forget the first canon of oak management, *regeneration is a process, not an event*. I commend to both researchers and managers a reading of some of the older literature. It is educational, sobering, and sometimes humbling. New and helpful interpretations are possible as pieces fit together and we learn the value of the knowledge base we inherited, and appreciate those who bequeathed it.

LITERATURE CITED

- Beck, D.E. 1988. Regenerating cove hardwood stands. In: Smith, H.C.; Perkey, A.W.; Kidd, W.E., Jr., eds. Guidelines for regenerating Appalachian hardwood stands. SAF Publ. 88-03. Morgantown, W. VA: 156-165.
- Bey, C.F. 1964. Advance oak reproduction grows fast after clearcutting. Journal of Forestry. 62:339-340.
- Broderick, S.H.; Kittredge, D.B.; Decker, D.J.; Connelly, N.A. 1991. Practicing foresters identify research needs. Journal of Forestry. 89(8):27-35.
- Callahan, J.C.; Fischer, B.C. 1982. Economics and silvicultural potential of 23 harvested Indiana woodlands. Bull. 362. W. Lafayette, In: Purdue University, Agriculture Experiment Station. 23 pp.
- Carvell, K.L.; Tryon, E. H. 1961. The effect of environmental factors on the abundance of oak regeneration beneath mature oak stands. Forest Science. 7:98-105.
- Carvell, K.L. 1979. Factors affecting the abundance vigor and growth response of understory oak seedlings. In: Regenerating oaks in upland hardwood forests. 1979 John S. Wright Forestry conference. Lafayette, IN: Purdue University: 23-26.
- Carvell, K.L. 1988. Field guide for analyzing hardwood stand regeneration. In: Smith, H.C.; Perky, A.W.; Kidd, W.E., Jr., eds. Guidelines for regenerating Applachian hardwood stands. SAF Publ. 88-03. Morgantown, W. VA: 148-155.

- Clark, F.B. 1970. Measures necessary for natural regeneration of oaks, yellow-poplar, sweetgum and black walnut The silviculture of oaks and associated species. Res. Pap. NE-144. U. S. Department of Agriculture, Forest Service: 1-16.
- Clark, F.B.; Watt, R. F. 1971. Silvicultural methods for regenerating oaks. In: Proceedings of the oak symposium. Morgantown, WV. Upper Darby, PA: U. S. Department of Agriculture, Forest Service: 34-43.
- Countryman, D.W.; Miller, H. R. 1989. Investment analysis of upland oak stands with sugar maple understories: management for oak vs. conversion to sugar maple in Iowa and Missouri. Northern Journal of Applied Forestry. 6:165-169
- Crow, T.R. 1988. Reproductive mode and mechanisms for self-replacement of northern red oak (Quercus ruba) a review. Forest Science. 34:19-40.
- Crow, T.R. 1991. Oak in a changing landscape [Abstract]. In: The oak resource in the upper Midwest conference. Publ. No. NR-BU5663-3. St. Paul, MN: University of Minnesota. 4 pp.
- Curtis, J.T. 1959. The vegetation of Wisconsin. Madison: University of Wisconsin Press. 657 pp.
- Downs, A.A.; McQuilken, W. E. 1944. Seed production of southern Appalachian oaks. Journal of Forestry. 42:913-920.
- George, D.W.; Fischer, B. C. 1989. The effects of site and age on tree reproduction in young upland hardwood clearcuts. In: Rink, G.; Budelsky, C.A., eds. Proceedings of the seventh central hardwood conference. NC-132. Carbondale, IL: U. S. Department of Agriculture, Forest Service: 40-473.
- Hawley, R.C. 1946. The practice of silviculture. Fifth ed. N.Y.: John Wiley and Sons, Inc. 354 pp.
- Hilt, D. E. 1985. Where has my yellow-poplar gone? Northern Journal of Applied Forestry. 2:67-69.
- Hough, F.B. 1878. Report upon forestry. Washington: G.P.O. 650 pp.
- Johnson, P.S. 1985. Regenerating oaks in the Lake States. P. 98-109 In: Johnson, J.E., ed. Proceedings of the challenges in oak management and utilization. Madison: Coop. Extension Service, University of Wisconsin: 161 pp.
- Johnson, P.S.; Jacobs, R. D.; Martin, A. J.; Godel, E. D. 1989. Regenerating northern red oak: three successful case histories. Northern Journal of Applied Forestry. 6:174-178.
- Korstian, C.F. 1927. Factors controlling germination and early survival in oak. Sch. For. Bull. 19. Yale University. 115 pp.
- Kuenzel, J.G.; McGuire, J. R. 1942. Response of chestnut oak reproduction to clear and partial cutting of overstory. Journal of Forestry. 40:238-243.
- Leffleman, L.J.; Hawley, R. C. 1925. Studies of Connecticut hardwoods; the treatment of advance growth arising as a result of thinnings and shelterwood cuttings. Yale University Sch. For. Bull. 15.
- Liming, F.G.; Johnston, J. P. 1944. Reproduction in oak-hickory forest stands of the Missouri Ozarks. Journal of Forestry. 42:175-180.
- Loftis, D.L. 1988. Regenerating oaks on high-quality sites, an update. In: Smith, H.C.; Perky, A.W.; Kidd, W.E, Jr., eds. Proc. Guidelines for regenerating Appalachian hardwood stands. SAF Publ. 88-03. Morgantown, WV: 199-209.

- McGee, C.E. 1989. Oldgrowth dynamics and mast production. In: McGee, C.E., ed. Proc. of the Workshop: Southern Appalachian Mast Management. Knoxville, TN: University of Tennessee: 37-41.
- Merz, R.W.; Boyce, S. G. 1958. Reproduction of upland hardwoods in southeastern Ohio. Tech. Pap. 155. U. S. Department of Agriculture, Forest Service, CSFES. 24 pp.
- Minckler, L.S. 1989 Intensive group selection silviculture in central hardwoods after 40 years. In: Rink, G.; Budelsky, C.A., eds. Proc. of the Seventh Central Hardwood Conference. NC-132. Carbondale, IN: U. S. Department of Agriculture, Forest Service: 35-39.
- North Central Forest Experiment Station. 1991. Do a good turn for your oak forest. NEFES News 1991 July. St. Paul, MN: U. S. Department of Agriculture, Forest Service.
- Nowacki, G.J.; Abrams, M. D.; Lorimer, C. G. 1990. Composition, structure and historical development of northern red oak stands along an edaphic gradient in North-Central Wisconsin. Forest Science. 36:276-292.
- Nyland, R.D. 1992. Exploitation and greed in eastern hardwood forests. Journal of Forestry. 90:33-37.
- Potzger, J.E.; Friesner, R. C. 1934. Some comparisons between virgin forests and adjacent areas of secondary succession. Butler University Botanical Studies. 3:85-98.
- Sander, I.L.; Williamson, M. J. 1957. Response of a mixed hardwood stand in eastern Kentucky to a harvest cutting. Journal of Forestry. 55:291-293.
- Sander, I.L.; Clark, F. B. 1971. Reproduction of upland hardwood forests in the Central States. Ag. Handbook No. 405. U. S. Department of Agriculture, Forest Service. 25 pp.
- Sander, I.L. 1972. Size of oak advance reproduction: Key to growth following harvest cutting. Res. Pap. NC-79. St. Paul, MN: U. S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 6 pp.
- Sander, I.L.; Johnson, P. S.; Watt, R. F. 1976. A guide for evaluating the adequacy of oak advance reproduction. Gen. Tech. Rep. NC 23. St. Paul, MN: U. S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 7 pp.
- Smith, D.M. 1962. The practice of silviculture. New York: John Wiley and Sons, Inc. 578 pp.
- Spencer, J.S., Jr.; Kingsley, N. P. 1991. The oak resource in the Upper Midwest: an overview. In: The oak resource in the Upper Midwest Conference. Publ. No. NR-BU-5663-3. St. Paul, MN: University of Minnesota: 11-17.
- Thirgood, J.V. 1971. The historical significance of oak. In: Oak symposium proceedings. Morgantown, WV. Upper Darby, PA: U. S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 1-17.
- Weitzman, S.; Trimble, G. R., Jr. 1957. Some natural factors that govern the management of oaks. Pap. No. 88. Upper Darby, PA: U. S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 40 pp.

Causes of the Oak Regeneration Problem

Craig G. Lorimer, Department of Forestry, School of Natural Resources, University of Wisconsin, Madison, WI 53706

ABSTRACT

Historical records indicate that oak species dominated much of the central hardwood forest at the time of settlement. However, many oak stands harvested in the past 40 years on average or good sites are now dominated to various degrees by other hardwood species. Several possible causes are discussed, including acorn predation, climatic change, damage to seedlings by insects and deer, and excessive competition resulting from decreased fire frequency. A number of these factors appear capable of reducing the success of oak regeneration or causing nearly complete failures. However, mature oak stands usually have several hundred to several thousand oak seedlings per acre present in the understory. The crux of the problem often appears to be the failure of these seedlings to survive and increase in vigor, even when released from competition. Slow juvenile growth occurs even where deer browsing is uncommon and appears to be an inherent trait of northern red oak and white oak on mesic and dry-mesic sites. Experimental evidence shows that the dense understory of shade-tolerant species on mesic sites is a major limiting factor to adequate oak seedling development, and that seedling development is markedly improved if the understory is removed. Historical evidence suggests that many of our existing oak stands on mesic sites either developed after fire or were periodically subjected to fire and other disturbances removing understory and subcanopy trees. Historical evidence on changes in fire frequency is reviewed, and problems with the fire hypothesis are discussed.

INTRODUCTION

If a panel of practicing foresters and researchers were asked to "round up the usual suspects" implicated in causing oak regeneration failures, it would probably not take long to produce a hefty list. Erratic seed production, predation of acorns by insects, rodents, and deer, defoliation and browsing of established seedlings by animals, and intense competition with other vegetation have all been implicated. Foresters also wonder about recent changes in the environment. Fires and grazing are no longer a dominant force, woodlands are fragmented in many areas, chestnut (*Castanea dentata*) is no longer a major component of the canopy, predator control has altered animal populations, new insects have been introduced from other continents, and the climate may be changing.

Are all these factors having a major impact on oak regeneration? It may be useful to make a distinction between an "aggravating factor" and a "limiting factor." An

aggravating factor may reduce the frequency of seedling establishment or the rate of seedling growth, but is not ordinarily capable by itself of causing complete regeneration failures. A limiting factor, on the other hand, can cause nearly complete regeneration failures even if no other aggravating factors are present. It is possible, of course, for certain factors to be merely aggravating at some times or places and limiting in others.

Because of this variability, we need to view "the oak regeneration problem" as having both local and regional aspects. Oaks, like other species, can have local regeneration failures in response to chance factors such as seed crop failure or locally intense browsing. But it is the relatively sudden and repeated regeneration failures over widely separated areas that have been the cause of recent concern. Questions naturally arise: Is there a common underlying cause behind the regional problem? Is the regional problem caused by the cumulative impact of a host of aggravating factors or primarily by one or more limiting factors?

The ideal solution to resolving these questions would be to conduct controlled experiments that would simultaneously evaluate a number of factors and would be replicated in several geographical areas. One study underway in the Northeast may, in a few years, help sort out the importance of various factors in that region (Galford and others 1991). In the meantime, a review of causal factors over the central hardwood region must rely on a combination of limited experimental evidence, observational data or empirical correlations, and historical evidence.

Explaining the Facts of the Case Faced with a host of possible causal factors and limited evidence, a reasonable place to start is to list a number of facts concerning the oak regeneration problem that must be explained by any convincing hypothesis. The following list includes some of the major considerations:

(1) Oaks have dominated much of the central hardwood region for over 6,000 years. At the end of the glacial episode, much of central hardwood region was occupied by a boreal forest of spruce and pine. About 6,000-9,000 years ago, this boreal forest was replaced by an oak-pine-mixed hardwood forest, which has retained dominance ever since (Craig 1969, Watts 1979, Delcourt and Delcourt 1985). This evidence from the pollen record is supported by independent historical evidence on the presettlement forest (ca. 1600-1850) from early travelers and government land survey records. For example, species of oaks comprised 40-80 percent of the witness trees in sizable areas of New Jersey, Pennsylvania, Virginia, Kentucky, and Illinois (Spurr 1951, Russell 1981, Leitner and Jackson 1981, Bryant and Martin 1988, Abrams and Downs 1990). Thus, while disturbances that accompanied European settlement may have increased oak dominance in some places, particularly at the margins of its range (Whitney 1987, Nowacki and others 1990), oak was already dominant over large sections of the landscape and maintained by natural regeneration long before European settlement.

(2) Oak regeneration failures appear to have been a widespread problem only in the last 50 years or so. This trend is seen most clearly by the fact that oak stands on mesic sites clearcut after about 1930 are now frequently dominated by species such as maples (*Acer rubrum* and *A. saccharum*), yellow-poplar (*Liriodendron tulipifera*), hickories (*Carya* spp.), and black cherry (*Prunus serotina*), while stands

originating prior to 1930 on similar sites are usually dominated by oaks (Johnson 1976, Heiligmann and others 1985, Beck and Hooper 1986, Hix and Lorimer 1991, Abrams and Nowacki 1992).

(3) Oak regeneration is often successful on the drier or below-average sites. The site specific nature of oak regeneration problems has been known for some time (e.g., Weitzman and Trimble 1957). Until more detailed habitat classifications are available, site index can serve as a rough guide. In many areas, sites with an oak SI of 60 ft. or less are likely to support stable, self-perpetuating oak forests, while those with SI > 60 are likely to convert wholly or in part to more mesic species (Hilt 1985, Hill 1987, Loftis 1988b, Zaldivar-Garcia and Tew 1991). Stable oak forests usually have a sizable component of the more xerophytic species, such as chestnut oak (*Quercus prinus*) and black oak (*Q. velutina*), whereas the stands most susceptible to regeneration failures usually occur on the more mesic sites dominated by northern red oak (*Q. rubra*) and white oak (*Q. alba*). A convincing hypothesis should be able to explain this clear habitat-related variability. For example, if insects or deer are proposed as limiting factors, is there evidence that animal damage is consistently less serious on SI 60 lands than on SI 70?

(4) The oak regeneration problem is geographically widespread. Pinpointing the geographical extent of oak regeneration failures is difficult because regeneration surveys taken in the first few years after harvest can be misleading (Bey 1964, Oliver 1978). Whether or not oak will be a major part of the canopy can only be determined with reasonable confidence after the forest has entered the "pole stage" of development, when the forest has developed a closed canopy and stratified into crown classes (e.g., after age 25). Using these criteria, unintended conversions of oak forests to other species have been documented most clearly on average and good sites in the Ridge and Valley section of central Pennsylvania (Abrams and Nowacki 1992), cove forests of the southern Appalachian Mountains (Beck and Hooper 1986), upland sites in southern Michigan (Gammon and others 1960), and upland sites of southwestern Wisconsin (Johnson 1976, Hix and Lorimer 1991). A similar study on an average site in southeastern Ohio suggests a significant reduction in the oak component (Heiligmann and others 1985). Succession of oak forests to sugar maple and other shade-tolerant species is less certain but still likely to occur on mesic sites examined in eastern Tennessee, central Indiana, southern Illinois, and the River Hills area of Missouri (McGee 1984, Parker and others 1985, Schlesinger 1989, Pallardy and others 1988).

From this evidence it is clear that oak regeneration problems are not confined to northern states at the margin of the range, which might otherwise be most affected by climatic change. Also, problems are occurring far outside the original range of American chestnut, as well as in areas such as the southern Appalachian Mountains where the forest is not highly fragmented.

REVIEW OF EVIDENCE ON POSSIBLE CAUSES

Because successful oak regeneration usually depends on the existence of seedlings in the understory before harvest (Sander 1972, McQuilkin 1975), poor seed crops and high rates of consumption by animals can have significant impacts on the ability of oaks to compete with other species. Unfavorable weather and insect damage can both lead to poor acorn crops (Cecich 1991). Most of the eastern

Page 75 of 404

Limited Production and Heavy Consumption of Acorns Upland oak species, however, have good seed crops at intervals of 3-5 years (Burns and Honkala 1990). Intervals between good seed years in white oak may be longer (Rogers 1990), and local factors might occasionally lead to regeneration failures from this cause. Smith (1993) has reported that average intervals between good acorn crops may be as long as 10 years in the central Appalachians.

Destruction of acorns by insects, rodents, and deer is probably a more important factor in most areas; a loss of 90 percent of the current crop is typical (Arend and Scholz 1969, Marquis and others 1976, Galford and others 1991). In a recent study in Pennsylvania, rodents removed virtually every unprotected acorn on the ground surface and 78 percent of the buried acorns. Insects destroyed 63 percent of the surface acorns protected from rodents (Galford and others 1991).

Marquis and others (1976) found that acorn production and the activity of insects and rodents varied greatly in different stands for unknown reasons. Compared to a stand with abundant advance regeneration, a stand with few oak seedlings (96 per acre) not only produced fewer acorns, but twice as many were damaged by insects, and pilferage by rodents was three times as high.

Such evidence suggests that destruction of acorns by animals can potentially be limiting factors in some locations. Whether it is a major cause of the regional problem is not as clear. Thorn and Tzilkowski (1991) point out that by burying acorns in well-distributed caches, small mammals may actually facilitate seedling germination. In that study, small mammal activity resulted in 28,000 well-distributed caches per acre. Thus, an acorn that disappears from a seed trap should not necessarily be considered "destroyed." More importantly, a review of many studies in the central hardwood region shows that most mature stands have several hundred to several thousand oak seedlings per acre at any one time (Carvell and Tryon 1961, Nowacki and Abrams 1992, Merritt and Pope 1991). Thus, seedling germination is not a limiting factor in many areas. Beck and Hooper (1986) documented an oak regeneration failure that occurred even though more than 5,000 oak seedlings per acre were present at the time of overstory removal.

Damage to Seedlings by Animals (Johnson 1985). A newly established cohort of seedlings in a mature stand on a good site in the Southern Appalachians had a 10-year survival rate of 10 percent, with negligible growth of the survivors (Loftis 1988a). Could repeated defoliation and other animal damage be a factor in causing high mortality and slow growth?

Evidence on seedling damage by insects is very fragmentary. An insect with a potentially serious impact on oak seedlings is the Asiatic oak weevil (*Cyrtepistomus castaneus*) because larvae feed on fine root hairs and adults feed on leaves (Triplehorn 1955, Roling 1979). The weevil, introduced from Japan and first recorded in New Jersey in 1933, is now distributed in most of the eastern states as far west as Missouri and Kansas. In Missouri, the weevil has a strong preference for oak and apparently has little impact on other hardwoods (Ferguson 1987). Linit and others (1986) reported 46 insect species, including the Asiatic oak weevil, associated with planted oak seedlings in Missouri. Leaf area losses averaged about 22 percent over a season. The impact on growth was not measured, but was not necessarily considered to be severe.

In addition to obtaining more evidence on the impact of low to moderate defoliation on growth and vigor, more evidence is needed on insect activity across habitat types. The introduction of the Asiatic oak weevil in about 1930 did coincide with the start of major oak regeneration failures, but for that to be considered a principal limiting factor, evidence would probably be needed that weevil activity is concentrated on mesic sites where most oak regeneration failures occur.

The potential for deer browsing to block the development of competitive oak advance regeneration is better documented than insect damage. The deer population on the Allegheny Plateau in Pennsylvania (often more than 30 per square mile) is sufficient to create open, park-like stands with little undergrowth. Where deer populations are high, browsing can occur on oak seedlings that are as short as 15 in. high (Galford and others 1991). Sometimes there is so little advance regeneration of hardwoods that clearcuts revert to grass and scattered shrubs (Marquis 1974).

Published evidence on the effects of deer browsing on oak is very limited outside of Pennsylvania. Similarly severe effects have been documented elsewhere, but the problem is often quite localized. High deer populations (34-59 per square mile) in a game preserve in Massachusetts have created savanna-like conditions, but in the surrounding region the deer average only 3-8 per square mile and browsing is limited (Healy and Lyons 1987). The intensity of deer browsing appears to vary greatly from place to place. Four underplanting trials of northern red oak in southern Wisconsin have shown little browsing in two counties with average deer populations of 18 per square mile (Pubanz and Lorimer 1992), but destructive levels of browsing in counties with average deer densities of 25-35 per square mile (Pubanz and Lorimer, personal observations). In the mountains of West Virginia, moderate-sized clearcuts (e.g., 20 acres) develop so much vegetation that the ability of deer to modify the outcome is limited (H. C. Smith, personal communication). While deer browsing was observed on oak seedlings in a southern Appalachian cove stand, and may have contributed to the slow growth rate (Beck 1970), the problem of slow growth persisted long after the deer density had greatly diminished (D. E. Beck, personal communication).

Deer browsing is clearly a limiting factor for oak regeneration in some places, and the substantial growth of deer populations that occurred in many areas around the 1930's does coincide with the beginning of widespread oak problems. However, the occurrence of oak regeneration failures in places where deer are not especially numerous makes a number of researchers feel that deer are generally more of an aggravating factor than a primary limiting factor. We need more evidence, however, on the effects of moderate deer browsing on growth rates, especially where deer may be browsing oak in preference to other species (George and others 1991). Furthermore, the deer problem seems to be getting progressively worse. Deer populations in the lower Midwest were historically low (USDA Forest Service 1970) but appear to have increased substantially in recent years (e.g., Ishmael 1990). As one forester in Indiana remarked, complaints about deer damage are becoming more numerous, and establishing a hardwood plantation is often like "setting the table for deer."

Excessive Shade and Competition Another explanation for the slow growth and high mortality of understory oak seedlings besides insect and deer damage is the detrimental effects of dense understory vegetation. Northern red oak and white oak both appear to have a growth strategy in which photosynthate is diverted to root growth at the expense of shoot development (Crow 1988, Dickson 1991). Seedlings may, therefore, develop a stout taproot and persist for many years despite repeated shoot dieback (Merz and Boyce 1956). When an opening occurs, such "seedling sprouts" are often capable of rapid growth (Bey 1964, Sander 1972).

This strategy is effective on dry-mesic or xeric sites where moisture may be limiting and a moderate amount of light reaches the forest floor, but it is a poor strategy on mesic sites where light levels are much lower (Dickson 1991). Shade-tolerant species such as maples have an important advantage over oaks because they can make significant height growth under a closed canopy, steadily increasing in both size and number until a nearly continuous subcanopy or a multi-storied layer of vegetation develops (Lorimer 1984). These added layers of foliage beneath a closed upper canopy intercept so much light that often less than 1 percent of full sunlight reaches the seedling layer (Hanson 1986, Pubanz and Lorimer 1992). Hanson (1986) demonstrated a negative carbon balance for northern red oak seedlings growing under a heavy canopy. As a result, seedlings often die once acorn reserves are exhausted, and even among the survivors a vigorous root system doesn't ordinarily develop (Crow 1988).

The ability to persist under dense shade appears to vary among oak species. White oak and chestnut oak, for example, are often considered to be moderately shade-tolerant (McGee 1981, McQuilkin 1990). However, the shade tolerance of oaks is markedly less than for many of its mesic competitors. The average 5-year mortality rate for large, overtopped saplings in a dry-mesic stand in southern New York was 45 percent for northern red oak and 26 percent for chestnut oak, but only 11 percent for red maple (Lorimer 1981). On a dry-mesic site in central Massachusetts, overtopped red oak had a 19-year mortality rate of 90 percent compared to only 16 percent for red maple (Lorimer 1983).

A recent field experiment in southwestern Wisconsin showed that even vigorous, nursery-grown northern red oak seedlings are mostly unable to survive when underplanted in mature, undisturbed oak forests on mesic and dry-mesic sites. After 5 years, over 70 percent of the planted seedlings had died on both sites. However, in plots on the same sites where the understory layer was removed, planted seedlings not only had 93 percent survival, but seedlings doubled in height and had an average of 35 leaves after 5 years (figures 1 and 2).

Such evidence suggests that the presence of a dense understory of shade-tolerant species is often sufficient to prevent the development of vigorous oak advance regeneration, whether or not other limiting or aggravating factors are present. The small oak seedlings typically present on average and good sites grow very slowly, and usually seem incapable of rapid growth even if the overstory is removed (Beck 1970, McQuilkin 1975, Sander 1972, Beck and Hooper 1986). A typical harvest operation on mesic sites creates a situation in which scattered oak seedlings, usually less than a foot tall and capable of growing only a few inches in height per year, must compete with thousands of tall saplings of other species capable of growing 18 to 24 in. in height per year (Beck and Hooper 1986; Hix and Lorimer 1990, 1991). Early spring frosts, which McGee (1986) found especially damaging to oak

Page 78 of 404

seedlings that originated beneath a canopy, could place oak seedlings at a further disadvantage.

In contrast to mesic sites, where one cohort of oak seedlings may largely disappear before the next one germinates, drier sites often have oak seedlings that may persist for up to 30-50 years, developing a strong root system and often a tall shoot as well (Liming and Johnston 1944, Sander 1972, Johnson 1991). Even if saplings of shade-tolerant competitors are present on these sites, they may not be numerous enough or vigorous enough to present a serious challenge. Development of vigorous oak seedlings on mesic sites is feasible, but it has only been demonstrated in cases where understory vegetation has been eradicated before or at the time of overstory removal (Johnson and Jacobs 1981, Johnson and others 1989, Lorimer 1989, Loftis 1990, Nowacki and others 1990). In the following section, evidence is reviewed on historical conditions that may have permitted the expansion of oaks onto mesic sites where they ordinarily would not be competitive.




Figure 1—Before (top) and after (bottom) understory removal and light thinning of the canopy on a mesic site dominated by northern red oak in southwestern Wisconsin. These treatments increased natural oak seedling establishment by 400 percent and greatly stimulated the growth and vigor of underplanted oak seedlings (Pubanz and Lorimer 1992).



Figure 2—(Left) Northern red oak seedlings underplanted in an undisturbed mature oak forest in southwestern Wisconsin had 74 percent mortality after only 5 years, and survivors had low vigor and no net height growth. (Right) Seedlings underplanted in stands where the understory was removed (see figure 1, bottom) had 93 percent survival and doubled in size. This experiment illustrated the inability of even vigorous, nursery-grown oak seedlings to survive beneath the shade-tolerant understory of a typical mature forest (Pubanz and Lorimer 1992).

HISTORICAL FACTORS

Origin of Existing Oak Stands If oak regeneration on mesic sites is so difficult, how did the existing stands originate? Detailed stand records rarely go back as far as 70 or 80 years, but comments by contemporary observers, supplemented by field evidence on stand history, can give us a general idea of stand origins.

The most thorough evidence comes from the Midwest, because it was the last part of the central hardwood region to be settled and has the best records. Many midwestern oak stands occur on sites that supported oak savannas at the time of settlement. These savannas were usually dominated by the more fire-resistant species such as bur oak (Q. macrocarpa), white oak, and black oak, and were "covered with trees about as far apart as in a common orchard" (Bayley 1954). Oak savannas are estimated to have covered between 13-32 million acres in eight states (Nuzzo 1986). Based on 19th century land survey records, Curtis (1959) estimated that southern Wisconsin at that time was approximately 45 percent oak savannas, 20 percent prairie and sedge meadow, 25 percent maple and floodplain forest, and 10 percent oak forest. There is little question that these oak savannas were fire-maintained and fire-dependent (Gleason 1913, Muir 1913, Curtis 1959, Grimm 1984). For example, in 1750, Father Vivier wrote that in the Ozark region, "trees are almost as thinly scattered as in our public promenades. This is partly due to the fact that the savages set fire to the prairies toward the end of autumn, when the grass is dry; the fire spreads everywhere and destroys most of

Page 80 of 404

the young trees" (Johnson 1992). In Wisconsin, the more fire-sensitive northern red oak was often present only as brushy sprouts or "grubs" among the prairie grasses (Curtis 1959). When the fires ceased, "the grubs grew up into trees, and formed tall thickets so dense that it was difficult to walk through them and every trace of the sunny *openings* vanished" (Muir 1913). Many of the red oak stands being harvested today appear to be the first—and only—generation of closed-canopy oak forest to follow the savannas. Stands harvested in recent decades have been displaced by the shade-tolerant species that quickly developed as an understory beneath the oaks (Johnson 1976, Jokela and Sawtelle 1985, Hix and Lorimer 1991).

Existing oak stands in states farther to the east, in the Piedmont-Appalachian region, originated after 50-100 years of influence by European settlers, and in a number of different ways:

(1) Some of our existing oak stands on mesic sites originated after intense fire and are even-aged (Brown 1960, Ward and Stephens 1989, Nowacki and others 1990).

(2) Some eastern oak stands apparently originated in a manner similar to those in the Midwest, after the cessation of repeated fires. In the southeastern Piedmont, Hammond (1880) noted that "since the discontinuation of spring and autumn fires, [a deer] could not be seen at fifteen paces, because of the thick growth of oak and hickory that has taken over the land." Grazing was also widespread at the time (Greeley and Ashe 1907) and may have had an effect similar to fire.

(3) Probably millions of acres of existing oak forest, both in the Northeast and Southeast, originated as an understory beneath a pine canopy (Hammond 1880, Oosting 1942). In central New England, 65-70 percent of the land had been cleared for agriculture at some point, but much was subsequently abandoned. Much of this land reverted to "old-field" pine. As can be seen today, old field pine stands usually have a relatively sparse understory, but oak saplings 3-15 ft. tall are common. Data gathered by McKinnon and others (1935) in central New England show an average of 578 saplings per acre released after the harvest of old-field pine on good sites, of which 215 were oak. The oaks typically became the dominant species in the subsequent stands (Oosting 1942, Lutz and Cline 1947).

(4) Many oak stands in the mid-Atlantic region appear to have originated after repeated, intensive clearcutting to produce charcoal for the iron and brick industries. Because stem quality was not an important consideration, small stems as well as large could be utilized, and clearcutting was often done at short intervals of 30-40 years (Schnur 1937, Raup 1938, Abrams and Nowacki 1992). Approximately 300 acres of forest had to be cleared each year to supply a single iron furnace (Stout 1933). Fires often followed. All of these factors would have favored vigorous sprouters such as oaks and hickories. The coppice nature of most stands was such a prominent feature of the landscape that Hawley and Hawes (1912) mapped a large area of southern New England, New York, and New Jersey as the "Sprout Hardwoods Region." These historical circumstances are much different from the prevailing situation today, when small trees are in little demand, and both logging and natural blowdown remove only the larger overstory trees and leave the small tree layer largely intact.

Although our existing oak forests on mesic sites have clearly originated in a variety of ways, the common denominator in all of these scenarios is that historical events resulted in a "low competition" environment in which sizable understory trees were either absent, sparse, or periodically eradicated.

Fire Effects on Oak Regeneration Most oak species have biological traits, such as relatively thick bark and protection against rootkill, that suggest an adaptation to periodic fire. Perhaps the most significant impact of fire is its reduction in the density of shade-tolerant saplings, which as noted above, is probably the most common inhibiting factor in oak regeneration. In a study in central New York, mortality of saplings caused by accidental spring fires ranged from 35 percent in ironwood (Ostrya virginiana) to 93 percent in hemlock (Tsuga canadensis), but only 12 percent of the oaks and hickories were root-killed (Swan 1970). Others have reported similar selective discrimination of fire against shade-tolerant competitors (Brown 1960, Carvell and Maxey 1969, Kruger 1992). Fire may also result in a major reduction in shrub cover (Nyland and others 1982, Abrams 1988, Reich and others 1990). As a result of this selective reduction of competitors, oaks may increase in relative density even if the absolute density remains unchanged.

Evidence for positive effects of fire on oak seedling establishment, on the other hand, is currently rather inconclusive. Several studies of prescribed burns have shown little or no increase in oak seedling establishment after the burn; in several cases, the fire resulted in a great influx of other species or resprouting of competitors as multiple-stemmed clumps (Sims 1932, Johnson 1974, McGee 1979, Nyland and others 1982, Wendel and Smith 1986, Merritt and Pope 1991). Others have produced evidence that burning can substantially increase oak numbers (Keetch 1944, Carvell and Tryon 1961, Carvell and Maxey 1969, Niering and others 1970, Thor and Nichols 1973, Little 1973).

There may be several reasons for these discrepancies. Many of the studies reporting discouraging results are based on short-term observations (2 years or less) after a single, low-intensity prescribed burn on good sites. If a good acorn crop doesn't occur during this period, a substantial improvement in oak seedling establishment is not likely. A single burn will often not accomplish much, and may even be counterproductive on mesic sites, where it may simply aid the establishment of competing vegetation. A number of researchers have suggested that repeated burning over a period of time may be necessary (Van Lear and Waldrop 1988, Loftis 1990, Johnson 1992). As Komarek (1974) once commented in relation to southern pine: "All too often investigators in the past have expected miraculous results with one light burn after years of fire exclusion." Nearly all of the studies reporting increases in oak seedling establishment involved several burns. Unfortunately, however, most of the reported successes with fire have occurred on dry sites. Clearly we need more evidence on the long-term effects of repeated burns on mesic sites.

The effect of intense wildfire on oak regeneration has received less attention, but it is this type of fire that has produced the most dramatic results on mesic sites. Carvell and Maxey (1969) reported that an October wildfire in a young stand of mixed cove hardwoods in West Virginia increased the number of dominant oaks from 93 to 293 per acre. Fifty-five years after an intense surface fire in a Connecticut mixed hardwood stand, burned sections increased oak and hickory

Page 82 of 404

dominance by 240 percent compared to unburned sections (Ward and Stephens 1989). Similar results were reported by Brown (1960) in Rhode Island. In the northern hardwood region, northern red oak has sometimes displaced the hemlock-maple type after intense slash fires (Nowacki and others 1990). In such cases where intensive site preparation or competition control has taken place, surprisingly rapid growth of newly established oak seedlings sometimes occurs that would not be expected under other circumstances (Johnson and others 1989, Lorimer 1989).

Climatic Change Oaks are usually considered to be favored by relatively warm and dry climates, and paleocologists often use abundant fossil oak pollen as one indicator of this type of climate. Oak regeneration failures could in principle be related to a shift to cooler and moister conditions. This change would probably have an effect of a southward shift in the boundary between northern (beech-maple) and central hardwoods (oak-hickory). In other regions, it might also cause an expansion of mesophytic species onto sites that were once relatively warm and dry, with a corresponding shrinkage of microhabitats that were formerly stable oak sites.

The issue of possible climatic effects needs further study, but several important lines of evidence seem rather inconsistent with the hypothesis that climatic change has been a major causal factor. There is little evidence in the pollen record that a rather dramatic cooling episode known as the "Little Ice Age" (ca. 1400-1850) had any substantial impact on oak dominance (Davis 1958, Watts 1979, Patterson and Backman 1988). Indeed, most early explorers in the Northeast described oak as the dominant forest type. Second, data show that the climate has been warmer (and drier) in recent decades compared to the late 19th and early 20th centuries (Wahl 1968), a trend that should not be favorable for the shade-tolerant competitors. Third, a relatively warm and dry climate does not seem to facilitate oak regeneration on mesic sites or hinder the development of mesic competitors to any noticeable degree. This relationship is apparent from the development of dense understories of sugar maple on mesic sites in southern Illinois, southwestern Wisconsin, and the River Hills section of Missouri, all of which were at or close to the original prairie border (Schlesinger 1989, Pallardy and others 1988, Hix and Lorimer 1991).

Historical Changes in Disturbance Frequency If a major cause of the regional oak regeneration problem has been the development of dense understories of competing species on mesic and dry-mesic sites, some evidence would be helpful in clarifying whether or not the magnitude of factors inhibiting understory development was sufficient prior to 1930 to account for the existing widespread occurrence of mature oak on mesic sites, and whether these factors have subsequently decreased.

The Post-Settlement Era (1750-1930) Most or all of the factors involved in the creation of existing oak stands, and discussed in previous sections, were widespread in the post-settlement era. Numerous sources, such as Van der Donck (1656), Kalm (1770), Harris (1805), and Hammond (1880) reported that the early settlers repeatedly burned the woods to improve hunting prospects and pasturage for cattle. As Hammond (1880) noted for the South Carolina Piedmont: "The early settlers in this region were stock raisers and kept up the Indian practice of burning off the woods during the winter." A more systematic survey of burning frequency was obtained by Hough (1882). Nearly all of his correspondents throughout the central hardwood region reported frequent, intentional burning, and three correspondents in Pennsylvania and Virginia volunteered estimates of the average percentage of forest land burned in a typical year. These estimates ranged from 2-14 percent of the land burned annually, equivalent to an average fire rotation period of 11 years (range of 7 to 50).

Frequent burning persisted until relatively recent times. In southern New England, Buttrick (1912) remarked that "many tracts are subject to recurring fires at frequent intervals, often annually." Graves and Fisher (1903) agreed that "woodlots which do not show some traces of fire are scarce." A major systematic survey of the southern Appalachian region was conducted by Ayres and Ashe (1905), who reported fire scars on 80 percent of the 6.5 million acres examined. Even as late as the 1920's, Logan (1975) described annual burning of woods by farmers in southwestern Wisconsin.

The advent of fire suppression programs such as the "Dixie Crusaders" and the Smokey Bear program in the 1930's and 1940's coincides well with the beginning of widespread oak regeneration problems. Based on fire scar evidence in the Great Smoky Mountains, Harmon (1982) showed a change in the fire rotation period of 10 years or less in the period prior to 1940 to over 2,000 years for the period 1940-1979. Current fire rotation periods for national forests in the central hardwood region range from 900-6,000 years (Haines and others 1975).

Fires, along with widespread grazing, led to open, park-like stands with little undergrowth (Greely and Ashe 1907, Korstian 1927). Schnur's (1937) data show that average 80-90 year old even-aged stands on site index 70 lands in the 1920's had only 6-8 trees per acre in the 2- to 4-in. diameter classes of all shade-tolerant species combined. The data of Gevorkiantz and Scholz (1948) in southwestern Wisconsin show similar trends, with an average of only 18-26 suppressed trees per acre in the 1- to 4-in. d.b.h. classes. McGill (1991) resampled fully-stocked, even-aged oak stands in the same geographical region and documented a dramatic increase in the density of shade-tolerant understory trees. Whereas stands in the earlier survey had bell-shaped size distributions, the current stands have an irregular, steeply descending distribution with the largest number of trees in the smallest size classes.

The Presettlement and Archaic Periods While disturbances caused by European settlers and subsequent generations are probably sufficient to account for the widespread occurrence of oak forests on sites where they don't ordinarily appear to be competitive, a plausible hypothesis is needed to explain how oaks were able to dominate the landscape for thousands of years before settlers arrived. It is a difficult issue because fire would probably have had to occur more than once during the lifespan of an oak stand to prevent succession to other species, particularly if the fires were of low or moderate intensity. Yet, the available evidence suggests that Indian populations were relatively small and lightning fires relatively uncommon (Schroeder and Buck 1970, Barden and Woods 1973).

The hypothesis that fire was a major factor in accounting for the dominance of oak forests on mesic sites does not presuppose that the entire landscape had to be Page 84 of 404

burned at frequent intervals. First, as already noted, many dry sites and some average sites appear to be able to maintain stable oak forests in the absence of fire. Second, sizable areas of the central hardwood region are known to have been covered by non-oak forest in presettlement times. Beech-maple forest, mixed with hemlock in some areas, covered large areas of New York, northern and central Pennsylvania, and northern Ohio. Mixed-mesophytic forest also occupied many of the good sites in West Virginia, eastern Kentucky, and central Tennessee (Sargent 1884, Braun 1950, Kuchler 1964, Whitney 1990). In the oak savanna regions of the Midwest, the role of fire is already established beyond reasonable doubt. Thus, the areas in which the frequency and impact of fire are most in question are the eastern Piedmont, the Ridge and Valley province, and the Southern Appalachian Mountains. The following is a brief overview of the available evidence in those areas and a discussion of problems that remain.

There are indeed many references by early travelers and colonists to the widespread practice of woods burning by Indians. In fact, few writers of the time who included any significant account of the vegetation failed to mention the practice or its effects on the forest. A number of these accounts, but by no means all, were reviewed by Maxwell (1910), Day (1953), and Lorimer (1985). The following description of coastal Massachusetts by Wood (1634) has many of the elements that occur frequently in these accounts:

There is no underwood saving in swamps and low grounds that are wet . . . for it being the custome of the Indians to burne the wood in November, when the grass is withered, and the leaves dried, it consumes all the underwood, and rubbish, which otherwise would over grow the Countrey, making it unpassable, and spoyle their much affected hunting; so that by this meanes in those places where the Indians inhabit, there is scarce a bush or bramble, or any cumbersome underwood to bee seene in the more champion ground. . .

Morton (1632) implied that the practice was so widespread in coastal Massachusetts that trees not scorched or damaged by fire were difficult to find:

So that hee that will looke to finde large trees, and good tymber. ...must seeke for them, (as I and others have done) in the lower grounds where the grounds are wett when the Country is fired ... for the Salvages by this Custome of theirs, have spoiled all the rest: for this Custome hath bin continued from the beginninge....

Since the fires were often set or accidentally started during periods of dry weather, and usually no attempt was made to control them, they could spread over large areas. Loskiel (1794) noted that "these fires run on for many miles," and Morton (1632) described the typical fire as "burning continually night and day, until a shower of rain falls to quench it." In New York, Van der Donck (1656) described the woods burning as a "yearly custom" and that some fires were quite intense, "for it frequently spreads and rages with such violence, that it is awful to behold."

Russell (1983) has challenged the idea of widespread intentional burning by Indians. She notes, with some justification, that early writers did not necessarily imply widespread burning of the landscape, but only "in those places where the Page 85 of 404 Indians inhabit" (Wood 1634) and "in all places where they come" (Morton 1632). She also feels that the writers may have been biased, and exaggerated the openness of the woods in order to attract more colonists. However, the descriptions of fire use and open woods with little undergrowth are consistent among many groups of people including priests and missionaries (Loskiel 1794, Rights 1947), surveyors (Lindestrom 1656, Byrd 1728), botanists (Bartram 1791), early historians (Norris 1890, Logan 1859), and early explorers and travelers with no specific land claims in the areas described (Smith 1616, Lawson 1709, Grant 1946). In Massachusetts, independent evidence can be found which supports the accounts of Indian burning related by Morton and Wood. Colonial records of Massachusetts indicate that in 1677 the court was "informed that great damage hath happened to several persons in the outskirt plantations by Indians kindling fires in the woods in the latter part of the yeare", and passed a law limiting the times during which Indians could set fires (Hough 1882).

A more difficult problem than writer bias is probably the fact that these early accounts do not provide a systematic view of the landscape. Early explorers often traveled along rivers, where evidence of Indian activities was more likely to be seen (e.g., Lawson 1709, Bartram 1791), or used Indian paths and trading paths connecting different villages. Also, early colonists tended to settle first in those areas that had been cleared by Indians (H. S. Russell 1980) which might not be representative of the general landscape.

The diary of Col. William Byrd is therefore of particular interest because in 1728 he was involved in surveying the boundary line between Virginia and North Carolina, and was therefore confined to a compass line. The frequent references to fire and fire effects do give the impression that fire was a major factor on the landscape in that region. Byrd made several references to extensive thickets of saplings that appeared to be of fire origin, as when he reported that they "Scuffled thro' a mighty thicket, at least three miles long. The whole was one continued tract of rich high land, the woods whereof had been burnt not long before. It was then overgrown with Saplings of Oak, Hiccory, and Locust "When the Byrd party reached the mountains, near a major Indian trail, they encountered a "great fire" which Byrd attributed to Indians. Most importantly, Byrd made general remarks about fire frequency, stating that "the woods are not there burnt every year, as they generally are amongst the Inhabitants. But the dead Leaves and Trash of many years are heapt up together, which being at length kindled by the Indians that happen to pass that way, furnishing fewel for a conflagration that carries all before it." The contrast in fire regimes between these mountain forests and areas more densely settled by Indians, which Byrd considered to be subject to widespread burning each year, provides independent confirmation of accounts elsewhere by Morton (1632), Wood (1634) and others.

The low importance of shade-tolerant species over extensive areas of the Piedmont and Ridge and Valley provinces in presettlement times provides indirect but important evidence on presettlement fire frequency. Tolerant species appear quite capable of dominating the overstories on many sites (e.g., Abrams and Downs 1990, Abrams and Nowacki 1992), so we must consider why these strong successional trends did not also take place in presettlement times. If fire was indeed the principal factor restricting the occurrence of these species (cf. Kline and Cottam 1979 and Grimm 1984 for midwestern examples), then the rarity of Page 86 of 404 late-successional forests on the uplands suggests the influence of fire may have been widespread and pervasive.

Scientific Evidence on Fire Frequency. Modern scientific studies can provide independent data for comparison with historical accounts. Studies of fire scars on living trees are unfortunately very limited because by the 20th century old-growth oak forest was already very rare. Apparently the only evidence for the eastern Piedmont/Appalachian region is based on a cross-section of a single large white oak in the New Jersey Piedmont, which, however, had several fire scars between the period 1641 and 1711 (Buell and others 1954). Some indication of presettlement fire frequency in other areas is given by a study by Lutz (1930) of a hemlock-beech forest on the Allegheny Plateau. In spite of the typically low fire frequency in this forest type, and the fact that the stand was located on an east-facing slope in a remote area near the headwaters of a small stream, Lutz found evidence of at least five major fire dates for the presettlement period 1725-1795. Fire scar studies should be done soon in the remaining old-growth Appalachian oak stands before the demise of old trees that were alive in presettlement times.

A promising approach to investigating presettlement fire frequency is the analysis of charcoal fragments in sediment cores from bogs and ponds. Although few studies of charcoal have yet been done, Watts (1979) and Patterson and Backman (1988) found charcoal throughout the sediment profiles in nine different sites in western Virginia, central Pennsylvania, coastal New Jersey, New York, and Massachusetts. In a number of cases there was no indication that charcoal was less abundant in the several thousand years prior to European settlement than afterward. In a site adjacent to the Little Tennessee River, Cridlebaugh (1984) also found charcoal throughout the profile. However, in this case charcoal abundance increased by an order of magnitude after A.D. 1000, when Indian agricultural activities intensified, and showed another dramatic rise at the time of European settlement.

Indian Settlement Patterns. How widespread were Indian effects on vegetation? Archaeological and historical evidence indicates that although villages were generally located near streams and rivers, Indians themselves used a wide range of environments and the pattern of use frequently shifted. Indians in the Archaic period (ca. 8000-1000 B.C.) were migratory hunter-gatherers and made the greatest use of upland habitats. In North Carolina, Ward (1983) noted that signs of the Archaic cultures "covered the Piedmont landscape, leaving a network of tracks that is hard to miss . . . The broad alluvial valleys, the rolling upland hills, and the banks of small streams were all occupied, visited, or utilized at some point during the 6,000 to 7,000-year span of the Archaic period." After the widespread adoption of agriculture (ca. A.D. 1000), there was a dramatic shift of village sites to the floodplains of major rivers and streams (Purrington 1983, Thornton 1990). However, upland sites continued to be used, perhaps as temporary base camps for deer hunting and nut harvesting (Ward 1983). In the Smoky Mountain region, 40-50 percent of food processing sites continued to occur in upland zones, including benches, coves, and saddles between mountains (Purrington 1983).

Knowledge of Cherokee villages and population levels is probably more accurate than for some other tribes, and can perhaps give some indication of the probable impact on the landscape. The Cherokee population in the early 1600's, prior to contact with European diseases, is estimated to have been between 22,000-30,000 Page 87 of 404 people distributed over approximately 52,000 square miles, with an average population density of about 16 per 25 square miles (Thornton 1991). The population was concentrated in at least 60 villages along major branches of the Hiwassee, Little Tennessee, and Tugaloo rivers. Goodwin's (1977) map of known villages (ca. 1700) shows that even villages on different rivers were usually not more than 15-20 miles apart, while villages on the same river were usually only a few miles apart. Because of this dispersion of villages across the region, it is likely that Indians could have had a major impact on the intervening upland areas. As Bishop Spangenberg commented in 1752: "The woods are full of Cherokees, and we see their signs wherever we go. They are out hunting." (Rights 1947).

It is important to keep in mind that Indians, even after the advent of agriculture, were migratory and hence would have had more impact than their numbers alone might suggest. In the early 1600's, John Smith wrote that when the Indians of coastal Virginia went hunting, "they leave their habitations, and reduce themselves into companies, as the Tartars doe, and goe to the most dessert [uninhabited] places with their families, where they spend their time in hunting and fowling up towards the mountains, by the heads of their rivers, where there is plentie of game . . . Having found the Deere, they environ them with many fires, and betwixt the fires they place themselves.". Loskiel (1794) likewise noted that Indians in the mid-Atlantic region commonly set out for long hunts lasting 3-4 weeks, and often several months. In addition, warfare and trade often prompted Indians to travel astonishingly long distances. For example, it was not uncommon for the Iroquois of central New York to travel to South Carolina to make war against the Catawbas and into Florida to fight the Creeks (Byrd 1728, Myer 1971). Long-distance travel for hunting, warfare, and trade was made possible by a complex and surprisingly Maps of known Indian trails, such as those in dense network of trails. Massachusetts (Russell 1980), Pennsylvania (Wallace 1965), and the Southeast (Myer 1971) show that relatively few stands would have been located more than 15 miles from the nearest Indian trail (figure 3). This would have rendered many stands in otherwise sparsely settled areas subject to occasional accidental or intentional fires.

A final resolution of the Indian fire question is not possible at the present time. It is probably reasonable to state, however, that the available historical and scientific evidence is consistent with the hypothesis that Indian fires were widespread and had a major impact on the landscape. More evidence on charcoal deposits in sediments and fire scars in old-growth stands may help clarify the issue of local fire frequency. Also, by integrating evidence on fire size distributions in the absence of suppression (e.g., Hough 1882) with evidence on Indian populations size and distribution, it might be possible to judge whether or not fires could have been frequent enough to account for the original distribution of oaks on sites where they don't ordinarily appear to be competitive.

CONCLUSIONS A number of factors, including poor seed production, consumption of acorns, damage to seedlings by animals, adverse spring weather, and poor development of oak seedlings under dense canopies, are known to contribute to oak regeneration failures. Each of these factors can probably lead to local oak regeneration failures at certain times and places. The degree to which the regional oak regeneration problem is caused by interactions of all of these factors is still not well known.

Page 88 of 404




Figure 3—The network of recorded Indian trails in Tennessee and surrounding areas in the early colonial period, compiled by W. E. Myer in 1923 (Myer 1971).

Relatively little evidence, for example, is available on ambient levels and resulting impacts of insect defoliation of oak seedlings or deer browsing over large areas.

The principal bottleneck to successful oak regeneration, nevertheless, does not usually seem to occur in the seedling germination stage. Most mature oak stands already have hundreds or thousands of oak seedlings per acre in the understory. A major limiting factor is that these seedlings usually have slow juvenile growth and low survival rates on good sites, even after the overstory is removed. Studies have shown that the development of a shade-tolerant understory layer is particularly detrimental to adequate development of oak advance regeneration, and historical evidence suggests that most oak stands developed in situations where fire, grazing, and other disturbances periodically removed this small-tree layer. The hypothesis that a major factor in the regional oak regeneration problem has been the widespread development of shade-tolerant saplings on mesic sites after alteration of the disturbance regime is consistent with life-history traits of several of the common upland oak species, and helps explain several observed features of the regional problem. For example, the disturbance hypothesis can account for the relatively sudden increase in oak regeneration failures after the 1930's and 1940's, the geographically widespread nature of the problem, and the fact that oak Page 89 of 404

regeneration has usually not been a problem on drier sites. Although limited quantitative evidence on fire frequency in presettlement times is perhaps the weakest aspect of the hypothesis, there is considerable historical and some scientific evidence that fire frequencies were much higher in presettlement and postsettlement times than at present. Additional field studies on fire history, and controlled experiments on factors affecting oak regeneration, will help resolve some of these uncertainties.

LITERATURE CITED Abrams, Marc D. 1988. Effects of prescribed fire on woody vegetation in a gallery forest understory in northeastern Kansas. Transactions of the Kansas Academy of Science. 91(3-4):63-70.

- Abrams, Marc D.; Downs, Julie A. 1990. Successional replacement of old-growth white oak by mixed mesophytic hardwoods in southwestern Pennsylvania. Canadian Journal of Forest Research. 20(12):1864-1870.
- Abrams, Marc D.; Nowacki, Gregory J. 1992. Historical variation in fire, oak recruitment, and post-logging accelerated succession in central Pennsylvania. Bulletin of the Torrey Botanical Club. 119(1):19-28.
- Arend, John L.; Scholz, Harold F. 1969. Oak forests of the Lake States and their management. Res. Pap. NC-31. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 36 pp.
- Ayres, H.B.; Ashe, W.W. 1905. The southern Appalachian forests. U.S. Geol. Surv. Prof. Pap. 37. 291 pp.
- Barden, L.S.; Woods, F.W. 1973. Characteristics of lightning fires in southern Appalachian forests. In: Proceedings of the annual tall timbers fire ecology conference: 345-361.
- Bartram, William. 1791. Travels of William Bartram. 1955 ed. New York: Dover Publications. 414 pp.
- Bayley, C.C. 1954. Western trip: an address given at Manlius, N.Y. Wisconsin Magazine of History. 37:237-239.
- Beck, D.E. 1970. Effect of competition on survival and height growth of red oak seedlings. Res. Pap. SE-56. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 7 pp.
- Beck, D.E.; Hooper, R.M. 1986. Development of a Southern Appalachian hardwood stand after clearcutting. Southern Journal of Applied Forestry. 10:168-172.
- Bey, C.F. 1964. Advanced oak reproduction grows fast after clearcutting. Journal of Forestry. 62:339-340.
- Braun, E.L. 1950. Deciduous forests of eastern North America. New York: Hafner. 596 pp.
- Brown, J.H. 1960. The role of fire in altering the species composition of forests in Rhode Island. Ecology. 41:310-316.
- Bryant, William S.; Martin, William H. 1988. Vegetation of the Jackson Purchase of Kentucky based on the 1820 General Land Office Survey. In: Snyder, David H. Proceedings of the first annual symposium on the natural history of lower Tennessee and Cumberland River valleys; 1988 March 11-12; Land Between the Lakes. Clarksville, TN: Center for Field Biology of Land Between the Lakes: 264-276.
- Buell, M.F.; Buell, H.F; Small, J.A. 1954. Fire in the history of Mettler's Woods. Bulletin of the Torrey Botanical Club. 81:253-255.

- Burns, Russell M.; Honkala, Barbara H. 1990. Silvics of North America. Vol. 2, Hardwoods. Agric.
 Hand. 654. Washington, DC: U.S. Department of Agriculture, Forest Service. 877 pp.
- Buttrick, P.L. 1912. The effect of forest fires on trees and reproduction in southern New England. Forestry Quarterly. 10:13.
- Byrd, William; Bassett, J.S., ed. 1728. The writings of Colonel William Byrd. New York: Burt Franklin, (1970). 461 pp.
- Carvell, K.L.; Tryon, E.H. 1961. The effect of environmental factors on the abundance of oak regeneration beneath mature stands. Forest Science. 7:98-105.
- Carvell, K.L.; Maxey, W.R. 1969. Wildfire destroys! Western Virginia Agriculture and Forestry. 2(2):4-5, 12.
- Cecich, Robert A. 1991. Seed production in oak. In: Laursen, Steven B.; DeBoe, Joyce F. The oak resource in the upper Midwest: implications for management; 1991 June 3-6; Winona, MN. St. Paul, MN: Minnesota Extension Service, University of Minnesota: 125-131.
- Craig, Alan J. 1969. Vegetational history of the Shenandoah Valley, Virginia. Geological Society of American Special Paper. 123:283-296.
- Cridlebaugh, P. A. 1984. American Indian and Euro-American impact on Holocene vegetation in the Lower Little Tennessee River Valley, East Tennessee. Knoxville: University of Tennessee. 225 pp. Dissertation.
- Crow, T.R. 1988. Reproductive mode and mechanisms for self-replacement of northern red oak (Quercus rubra)--a review. Forest Science. 34:19-40.
- Curtis, J.T. 1959. The vegetation of Wisconsin. Madison: University of Wisconsin Press. 657 pp.
- Davis, Margaret B. 1958. Three pollen diagrams from central Massachusetts. American Journal of Science. 256:540-570.
- Day, G.M. 1953. The Indian as an ecological factor in the northeastern forest. Ecology. 34:329-346.
- Delcourt, Hazel R.; Delcourt, Paul A. 1985. Quaternary palynology and vegetational history of the southeastern United States. In: Bryant, V.M.; Holloway, R.G., eds. Pollen records of late-Quaternary North American sediments. American Association of Stratigraphic Palynologists Foundation: 1-28.
- Dickson, Richard E. 1991. Episodic growth and carbon physiology in northern red oak. In: Laursen, Steven B.; De Boe, Joyce F. The oak resource in the upper Midwest: implications for management; 1991 June 3-6; Winona, MN. St. Paul, MN: Minnesota Extension Service, University of Minnesota: 117-124.
- Ferguson, Carol S. 1987. Distribution and host plant preference of the Asiatic oak weevil, *Cyrtepistomus castaneus* (Roelofs), in a Missouri oak-hickory forest. Columbia: University of Missouri-Columbia. 125 pp. Dissertation.
- Galford, Jimmy; Auchmoody, L. R.; Smith, H. C.; Walters, R. S. 1991. Insects affecting establishment of northern red oak seedlings in central Pennsylvania. In: McCormick, Larry H.; Gottschalk, Kurt W. Proceedings of the 8th central hardwood forest conference; 1991 March 4-6; University Park, PA. Gen. Tech. Report NE-148. Radnor, PA: Northeastern Forest Experiment Station: 271-280.
- Gammon, A.D.; Rudolph, V.J.; Arend, J.L. 1960. Regeneration following clearcutting of oak during a seed year. Journal of Forestry. 58:711-715.
- George, David W.; Bowersox, Todd W.; McCormick, Larry H. 1991. Effectiveness of electric deer fences to protect planted seedlings in Pennsylvania. In: McCormick, Larry H.; Gottschalk, Kurt

W. Proceedings of the 8th central hardwood forest conference; 1991 March 4-6; University Park, PA. Gen. Tech. Report NE-148. Radnor, PA: Northeastern Forest Experiment Station: 395-401.

- Gevorkiantz, S. R.; Scholz, H. F. 1948. Timber yields and possible returns from the mixed-oak farmwoods of southwestern Wisconsin. Publ. No. 521. U.S. Department of Agriculture, Forest Service, Lake States Forest Experiment Station. 72 pp.
- Gleason, H.A. 1913. The relation of forest distribution and prairie fires in the Middle West. Torreya. 13:173-181.
- Goodwin, Gary C. 1977. Cherokees in transition: a study of changing culture and environment prior to 1775. Chicago: Department of Geography, University of Chicago. 207 pp.
- Grant, W. L., ed. 1946. Voyages of Samuel de Champlain 1604-1618. New York: Barnes and Noble. 374 pp.
- Graves, Henry S.; Fisher, R. T. 1903. The woodlot: a handbook for owners of woodlands in southern New England. Bulletin No. 42. Washington, D.C.: U.S. Department of Agriculture, Bureau of Forestry. 89 pp.
- Greeley, W.B.; Ashe, W.W. 1907. White oak in the southern Appalachians. Circular 105. Washington, D.C.: U.S. Department of Agriculture, Forest Service.
- Grimm, E.C. 1984. Fire and other factors controlling the Big Woods vegetation of Minnesota in the mid-nineteenth century. Ecological Monographs. 54:291-311.
- Haines, D.A.; Johnson, V.J.; Main, W.A. 1975. Wildfire atlas of the northeastern and north central states. Gen. Tech. Rep. NC-16. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 25 pp.
- Hammond, H. H. 1880. Tenth U.S. Census, South Carolina. Washington, D.C.: U.S. Government Printing Office. 63 pp.
- Hanson, P. J. 1986. Studies of Quercus rubra L. seedling dry matter accumulation, morphological development, and carbon dioxide exchange under controlled conditions. University of Minnesota. 169 pp. Dissertation.
- Harmon, M. 1982. Fire history of the westernmost portion of Great Smoky Mountains National Park. Bulletin of the Torrey Botanical Club. 109:74-79.
- Harris, T.M. 1805. The journal of a tour into the territory northwest of the Alleghany Mountains, made in the spring of the year 1803. In early western travels 1748-1846, Vol.3 (ed. by R.G. Thwaites). Cleveland, Ohio: Clark Co: 307-382.
- Hawley, R.C.; Hawes, A.F. 1912. Forestry in New England. New York: Wiley and Sons. 479 pp.
- Healy, William M.; Lyons, Paul J. 1987. Deer and forests on Boston's municipal watershed after 50 years as a wildlife sanctuary. In: symposium proceedings, deer, forestry, and agriculture: interactions and strategies for management; 1987 June 15-17; Warren, PA: Allegheny Society of American Foresters: 3-21.
- Heiligmann, R.B.; Norland, E.R.; Hilt, D.E. 1985. 28-year-old-reproduction on five cutting practices in upland oak. Northern Journal of Applied Forestry. 2:17-22.
- Hill, D.B. 1987. Oak-hickory regeneration in eastern Kentucky. In: Hay, R.L.; Woods, F.W.; De Selm, H., ed. Proceedings of the central hardwood forest conference VI. Knoxville, TN.: University of Tennesse: 111-118.
- Hilt, D.E. 1985. Species composition of young central hardwood stands that develop after clearcutting. In: Dawson, J.O.; Majerus, K.A., ed. Proceedings of the fifth central hardwood forest conference. Urbana-Champaign, IL: University of Illinois: 11-14.

- Hix, David M.; Lorimer, Craig G. 1990. Growth-competition relationships in young hardwood stands on two contrasting sites in southwestern Wisconsin. Forest Science. 36(4):1032-1049.
- Hix, David M.; Lorimer, Craig G. 1991. Early stand development on former oak sites in southwestern Wisconsin. Forest Ecology and Management. 42:169-193.
- Hough, Franklin B. 1882. Report on forestry, submitted to Congress by the Commissioner of Agriculture. Washington, D.C.: U.S. Government Printing Office.
- Ishmael, William E. 1990. Counting heads in the herd. Wisconsin Natural Resources. 14(5):21-22, 27-28.
- Johnson, Paul S. 1974. Survival and growth of northern red oak seedlings following a prescribed burn. Res. Note NC-177. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 3 pp.
- Johnson, Paul S. 1976. Modal development of regeneration in clearcut red oak stands in the Driftless Area. In: Fralish, James S.; Weaver, George T.; Schlesinger, Richard C. Proceedings of the central hardwood forest conference; 1976 October 17-19; Carbondale, IL: Department of Forestry, Southern Illinois University: 455-475.
- Johnson, Paul S. 1985. Regenerating oaks in the Lake States. In: Johnson, James E. Proceedings, Challenges in oak management and utilization; 1985 March 28-29; Madison, WI: Cooperative Extension Service, University of Wisconsin-Extension: 98-109.
- Johnson, Paul S. [in press]. New perspectives and the silviculture of oak-dominated forests in the central and eastern states. In: New Perspectives. Corvallis, OR: Oregon State University Press.
- Johnson, P.S.; Jacobs, R.D. 1981. Northern red oak regeneration after preherbicided clearcutting and shelterwood removal cutting. Res. Pap. NC-202. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station.
- Johnson, Paul S.; Jacobs, Rodney, D.; Martin, A. Jeff; Godel, Edwin D. 1989. Regenerating northern red oak: three successful case histories. Northern Journal of Applied Forestry. 6:174-178.
- Jokela, J. J.; Sawtelle, R. A. 1985. Origin of oak stands on the Springfield Plain: a lesson on oak regeneration. In: Dawson, Jeffrey O.; Majerus, Kimberly A. 5th central hardwood forest conference; 1985 April 15-17; Publ. 85-05. Urbana-Champaign, IL: Society of American Foresters: Department of Forestry, University of Illinois: 181-188.
- Kalm, P. 1770. Travels into North America (transl. by Forster, J.R.). Barre, Mass: Imprint Soc., (1972). 514 pp.
- Keetch, J.J. 1944. Sprout development on once-burned and repeatedly-burned areas in the Southern Appalachians. Tech. Note 59. U.S. Forest Service, Appalachian Forest Experiment Station. 3 pp.
- Kline, Virginia M.; Cottam, Grant. 1979. Vegetation response to climate and fire in the Driftless Area of Wisconsin. Ecology. 60(5):861-868.
- Komarek, E. V. 1974. Effects of fire on temperate forests and related ecosystems: Southeastern United States. In: Kozlowski, T. T.; Ahlgren, C. E., eds. Fire and ecosystems. New York: Academic Press. Chapter 8.
- Korstian, C.F. 1927. Factors controlling germination and early survival of oaks. Bulletin 19. Yale University School of Forestry. 115 pp.
- Kruger, Eric. 1992. Survival, growth and ecophysiology of northern red oak (*Quercus rubra* L.) and competing tree regeneration in response to fire and related disturbance. University of Wisconsin-Madison. Dissertation.

- Kuchler, A. W. 1964. Potential natural vegetation of the conterminous United States. Spec. Publ. 36. American Geographical Society.
- Lawson, John. 1709. A new voyage to Carolina. 1967 ed.; Lefler, Hugh T., ed. Chapel Hill, NC: University of North Carolina Press. 305 pp.
- Leitner, L.A.; Jackson, M.T. 1981. Presettlement forests of the unglaciated portion of southern Illinois. American Midland Naturalist. 105:290-304.
- Liming, Franklin G.; Johnston, John P. 1944. Reproduction in oak-hickory forest stands of the Missouri Ozarks. Journal of Forestry. 42:175-180.
- Lindestrom, Peter. 1656. Geographia Americae. (Johnson, A., transl.). Philadelphia: Swedish Colonial Soc., (1925). 418 pp.
- Linit, M. J.; Johnson, P. S.; McKinney, R. A.; Kearby, W. H. 1986. Insects and leaf area losses of planted northern red oak seedlings in an Ozark forest. Forest Science. 32(1):11-20.
- Little, S. 1973. Eighteen-year changes in the composition of a stand of *Pinus echinata* and *Pinus rigida* in southern New Jersey. Bulletin of the Torrey Botanical Club. 100:94-102.
- Loftis, David L. 1988a. Regenerating oaks on high-quality sites; an update. In: Workshop proceedings: Guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV: 199-209.
- Loftis, David L. 1988b. Species composition of regeneration after clearcutting southern Appalachian hardwoods. In: Miller, James H. (comp.) Fifth biennial southern silviculture research conference; 1988 Nov. 1-3; Memphis, TN. Gen. Tech. Rep. SO-74. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 252-257.
- Loftis, David L. 1990. A shelterwood method for regenerating red oak in the southern Appalachians. Forest Science. 36(4):917-929.
- Logan, Ben. 1975. The land remembers. Heartland Press. 281 pp.
- Logan, John H. 1859. History of the upper country of South Carolina from the earliest period to the close of the War of Independence. Charleston, SC: S. G. Courtenary & Co. Vol. I. 521 pp.
- Lorimer, Craig G. 1981. Survival and growth of understory trees in oak forests of the Hudson Highlands, New York. Canadian Journal of Forest Research. 11:689-695.
- Lorimer, Craig G. 1983. A test of the accuracy of shade-tolerance classifications based on physiognomic and reproductive traits. Canadian Journal of Botany. 61:1595-1598.
- Lorimer, Craig G. 1984. Development of the red maple understory in north-eastern oak forests. Forest Science. 30:3-22.
- Lorimer, Craig G. 1985. The role of fire in the perpetuation of oak forests. In: Johnson, J.E., ed. proceedings, challenges in oak management and utilization. Madison, WI: Coop. Extension Service, University of Winconsin: 8-25.
- Lorimer, Craig G. 1989. The oak regeneration problem: new evidence on causes and possible solutions. Forest Resource Analyses No. 8, Publ. R3484. Madison, WI: Department of Forestry, University of Wisconsin-Madison. 31 pp.
- Loskiel, George H. 1794. History of the mission of the United Brethren among the Indians in North America. Trans. London: by C. I. La Trobe. 159 pp.
- Lutz, H. J. 1930. The vegetation of Heart's Content, a virgin forest in northwestern Pennsylvania. Ecology. 11:2-29.

- Lutz, R.J.; Cline, A.C. Results of the first thirty years of experimentation in silviculture in the Harvard Forest, 1908-1938. I. The conversion of stands of old field origin by various methods of cutting and subsequent cultural treatments. Harvard Forest Bulletin 23. 182 pp.
- Marquis, D. A. 1974. The impact of deer browsing on Allegheny hardwood regeneration. Res. Pap. NE-308. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 8 pp.
- Marquis, D.A.; Eckert, P.L.; Roach, B.A. 1976. Acorn weevils, rodents, and deer all contribute to oak regeneration difficulties in Pennsylvania. Res. Pap. NE-356. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.
- Maxwell, Hu. 1910. The use and abuse of forests by the Virginia Indians. William and Mary College Quarterly Historical Magazine. 19(2):73-103.
- McGee, Charles E. 1979. Fire and other factors related to oak regeneration. In: Proceedings of the 1979 John S. Wright Forestry Conference, Regenerating oaks in upland hardwood forests: West Lafayette, IN: Purdue University: 75-81.
- McGee, Charles E. 1981. Response of white oak to release. Res. Note SO-273. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 pp.
- McGee, Charles E. 1984. Heavy mortality and succession in a virgin mixed mesophytic forest. Res. Pap. SO-209. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 9 pp.
- McGee, Charles E. 1986. Budbreak for twenty-three upland hardwoods compared under forest canopies and in recent clearcuts. Forest Science. 32:924-935.
- McGill, David W. 1991. Stocking relationships of northern red oak in Wisconsin. University of Wisconsin-Madison. 50 pp. Thesis
- McKinnon, F.S.; Hyde, G.R.; Cline, A.C. 1935. Cut-over old field pine lands in central New England: a regional study of the composition and stocking of the ensuing volunteer stands. Harvard Forest Bulletin. 80 pp.
- McQuilkin, R.A. 1975. Growth of four types of white oak reproduction after clearcutting in the Missouri Ozarks. Res. Pap. NC-116. St.Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 5 pp.
- McQuilkin, Robert A. 1990. Chestnut oak. In: Burns, Russel M.; Honkala, Barbara H. Silvics of North America. Agriculture Handbook 654. Washington, D.C.: U.S. Department of Agriculture, Forest Service: 721-726.
- Merritt, Clair; Pope, Phillip E. 1991. The effect of environmental factors, including wildfire and prescribed burning, on the regeneration of oaks in Indiana. Station Bull. 612. West Lafayette, IN: Department of Forestry and Natural Resources, Purdue University. 52 pp.

Merz, Robert W.; Boyce, Stephen G. 1956. Age of oak "seedlings". Journal of Forestry. 54:774-775.

Morton, Thomas. 1632. New English Canaan. Charles Green. 125 pp.

Muir, John. 1913. The story of my boyhood and youth. New York. 294 pp.

Myer, William E. 1971. Indian trails of the Southeast. Nashville, TN: Blue and Gray Press. 123 pp.

Niering, W.A.; Goodwin, R.H.; Taylor, S. 1970. Prescribed burning in southern New England: Introduction to long-range studies. In: Proc. 10th annual tall timbers fire ecology conference. Tallahassee, FL: Tall Timbers Research Station: 267-286.

Norris, J. E. 1890. History of the lower Shenandoah Valley. 1972 ed. Chicago: A. Warner. 812 pp. Page 95 of 404

- Nowacki, G. J.; Abrams, M. D.; Lorimer, C. G. 1990. Composition, structure, and historical development of northern red oak stands along an edaphic gradient in north-central Wisconsin. Forest Science. 36: 276-292.
- Nowacki, Gregory J.; Abrams, Marc D. 1992. Community, edaphic, and historical analysis of mixed oak forests of the Ridge and Valley province in central Pennsylvania. Canadian Journal of Forest Research. 22:790-800.
- Nuzzo, V. A. 1986. Extent and status of midwest oak savanna: presettlement and 1985. Natural Areas Journal. 6:6-36.
- Nyland, R.D.; Abrahamson, L.P.; Adams, K.B. 1982. Use of prescribed fire for regenerating red oak and white oak in New York. In: Proc. Soc. American Foresters National Convention: 163-167.
- Oliver, Chadwick D. 1978. The development of northern red oak in mixed stands in central New England. Bull. 91. New Haven, CT: School of Forestry and Environmental Studies, Yale University. 63 pp.
- Oosting, Henry J. 1942. An ecological analysis of the plant communities of Piedmont, North Carolina. American Midland Naturalist. 28:1-126.
- Pallardy, S.G.; Nigh, T.A.; Garrett, H.E. 1988. Changes in forest composition in central Missouri: 1968-1982. American Midland Naturalist. 120:380-390.
- Parker, G.R.; Leopold, D.J.; Eichenberger, J.K. 1985. Tree dynamics in an old-growth, deciduous forest. Forest Ecology Management. 11:31-57.
- Patterson, William A.; Backman, Andrew E. 1988. Fire and disease history of forests. In: Huntley B.; Webb, T., eds. Vegetation history. Kluwer Academic Publishers: 603-632.
- Pubanz, Dan M.; Lorimer, Craig G. 1992. Oak regeneration experiments in southwestern Wisconsin: two-year results. Res. Rep. R3552. Madison, WI: School of Natural Resources, College of Agricultural and Life Sciences, University of Wisconsin-Madison. 16 pp.
- Purrington, Burton L. 1983. Ancient mountaineers: an overview of prehistoric archaeology of North Carolina's western mountain region. In: Mathis, Mark A.; Crow, Jeffrey J. The prehistory of North Carolina. North Carolina Division of Archives and History. Chapter 3.
- Raup, H.M. 1938. Botanical studies in the Black Rock Forest. Black Rock Forest Bulletin 7. 161 pp.
- Reich, P. B.; Abrams, M. D.; Ellsworth, D. S.; Kruger, E. L.; Tabone, T. J. 1990. Fire affects ecophysiology and community dynamics of central Wisconsin oak forest regeneration. Ecology. 71:2179-2190.
- Rights, Douglas L. 1947. The American Indian in North Carolina. Durham, NC: Duke University Press. 296 pp.
- Rogers, Robert. 1990. White oak. In: Burns, Russell M.; Honkala, Barbara H. Silvics of North America. Agriculture Handbook 654. Washington, D.C.: U.S. Department of Agriculture, Forest Service: 605-613.
- Roling, M. 1979. Biology of the Asiatic oak weevil in central Missouri. University of Missouri-Columbia. 79 pp. Dissertation.
- Russell, E.W.B. 1981. Vegetation of northern New Jersey before European settlement. American Midland Naturalist. 105:1-12.
- Russell, E.W.B. 1983. Indian-set fires in the forests of the northeastern United States. Ecology. 64:78-88.

- Russell, H.S. 1980. Indian New England before the Mayflower. Hanover, NH: University Press of New England. 284 pp.
- Sander, I.L. 1972. Size of oak advance reproduction: Key to growth following harvest cutting. Res. Pap. NC-79. U.S. Department of Agriculture, Forest Service.
- Sargent, C.S. 1884. Report on the forests of North America. U.S. Department Interior, Census Office, 10th Census.
- Schlesinger, R.C. 1989. Dynamics of the sugar maple component of a white oak-yelow-popular community. In: Rink, G.; Budelsky, C.A., eds. Proceedings, 7th central hardwood forest conference. Gen. Tech. Rep. NC-132. Carbondale, IL: U.S. Department of Agriculture, Forest Service: 262-266.
- Schnur, G.L. 1937. Yield, stand, and volume tables for even-aged upland oak forests. Technical Bulletin 560. U.S. Department of Agriculture. 88 pp.
- Schroeder, Mark J.; Buck, Charles C. 1970. Fire weather. Agriculture Handbook 360. Washington, D.C.: U.S. Department of Agriculture, Forest Service. 229 pp.
- Sims, I. H. 1932. Establishment and survival of yellow-poplar following a clearcutting in the southern Appalachians. Journal of Forestry. 30:409-414.
- Smith, H. Clay. 1993. Regenerating oak in the central appalachians. In: Loftis, D.L.; Hodges, J.; Smith, D.; Johnson, P., Proceedings, A Symposium on Oak Regeneration; 1992 September 8-10; Knoxville, TN.
- Smith, J. 1616. A description of New England. In Sailors narratives of voyages along the New England coast 1524-1624. Boston: Houghton-Mifflin, (1905). 212-247 pp.
- Spurr, S.H. 1951. George Washington, surveyor and ecological observer. Ecology. 32:544-549.
- Stout, W. 1933. The charcoal iron industry of the Hanging Rock iron district: its influence on the early development of the Ohio Valley. Ohio State Archaeological and Historical Quarterly. 42:72-104.
- Swan, F.R. 1970. Postfire response of four plant communities in southcentral New York State. Ecology. 51:1074-1082.
- Thor, E.; Nichols, G.M. 1973. Some effects of fires on litter, soil, and hardwood regeneration. In: Proc. 13th annual tall timbers fire ecology conference. Tallahassee, FL: Tall Timbers Research Station: 317-329.
- Thorn, Eugene R.; Tzilkowski, Walter M. 1991. Mammal caching of oak acorns in a red pine and a mixed-oak stand. In: McCormick, Larry H.; Gottschalk, Kurt W. Proceedings, 8th central hardwood forest conference; 1991 March 4-6; University Park, PA. Gen. Tech. Rep. NE-148. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 299-304.
- Thornton, Russell. 1990. The Cherokees: a population history. Lincoln, NE: University of Nebraska Press.
- Triplehorn, C. A. 1955. The Asiatic oak weevil in Delaware. Journal of Economic Entomology. 48:289-293.
- U.S. Department of Agriculture. 1970. White-tailed deer in the Midwest, a symposium; 1968 December 9; Columbus, OH. Res. Pap. NC-39. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 34 pp.
- Van der Donck, A. 1656. A description of the New Netherlands (O'Donnell, T.F., ed.). Syracuse, NY: Syracuse University Press, (1968). 139 pp.

- Van Lear, David H.; Waldrop, Thomas A. 1988. Effects of fire on natural regeneration in the Appalachian Mountains. In: Workshop proceedings: guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV: 56-70.
- Wahl, E. W. 1968. A comparison of the climate of the eastern United States during the 1830s with the current normals. Monthly Weather Review. 96(2):73-82.
- Wallace, P.A.W. 1965. Indian paths of Pennsylvania. Harrisburg, PA: Pennsylvania Historical & Museum Commission.
- Ward, H. Travick. 1983. A review of archaeology in the North Carolina piedmont: a study of change. In: Mathis, Mark A.; Crow, Jeffrey J., eds. The prehistory of North Carolina: an archaeological symposium. Raleigh, NC: North Carolina Division of Archives and History: 53-82.
- Ward, J.S.; Stephens, G.R. 1989. Long-term effects of a 1932 surface fire on stand structure in a Connecticut mixed-hardwoods forest. In: Rink, G.; Budelsky, C.A., eds. Proceedings, 7th central hardwood forest conference, Southern Illinois University. Gen. Tech. Rep. NC-132. Carbondale, IL: U.S. Department of Agriculture, Forest Service: 267-273.
- Watts, W. A. 1979. Late Quaternary vegetation of central Appalachia and the New Jersey coastal plain. Ecological Monographs. 49:427-469.
- Weitzman, S.; Trimble, G.R. 1957. Some natural factors that govern the management of oaks. Station Pap. 88. Northeastern Forest Experiment Station.
- Wendel, G. W.; Smith, H. C. 1986. Effects of prescribed fire in a central Appalachian oak-hickory stand. Res. Pap. NE-594. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 8 pp.
- Whitney, G. G. 1987. An ecological history of the Great Lakes forest of Michigan. Journal of Ecology. 75:667-684.
- Whitney, G. G. 1990. The history and status of the hemlock-hardwood forests of the Allegheny plateau. Journal of Ecology. 78:443-458.
- Wood, W. 1634. New England's prospect. Boston, MA: Prince Soc., (1865). 110 pp.
- Zaldivar-Garcia, Pilar; Tew, D. Thompson. 1991. Ten year regeneration of southern Appalachian hardwood clearcuts after controlling residual trees. In: McCormick, Larry H.; Gottschalk, Kurt W. Proceedings, 8th central hardwood forest conference; 1991 March 4-6; University Park, PA. Gen. Tech. Rep. NE-148. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 332-346.

Oak Regeneration: The Scope of the Problem

David Wm. Smith, College of Forestry and Wildlife Resources, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061

ABSTRACT An evaluation of oak regeneration, based on published literature and direct input from researchers in the field, is presented for 31 oak species native to the United States. The evaluated species include 12 from the white oak group, 18 from the red oak group, and one from the intermediate oak group. There are 25 species from the eastern United States and six from the western United States. Twenty of the species are indigenous to upland sites and 11 are considered bottomland species. A general summary of key silvical characteristics is provided. An attempt has been made to assess regeneration success of each species in terms of geographic location, site quality (two levels), and regeneration type (seedling, seedling sprout, or stump sprout). A qualitative evaluation of the level of research knowledge available about the regeneration of each species is given.

OVERVIEW Quercus is the classical Latin name of the oaks. It is thought to be of Celtic derivation meaning *fine* and *tree* (Little 1979). Oak is the largest tree genus in the United States and is the most important hardwood genus. The oaks are the major component of eastern deciduous forests and dominate stands in central and southern upland forests. In addition, they are a significant component of mixed bottomland hardwoods throughout the eastern and southern United States, and occur on valley and slope sites in the semi-arid regions of the western United States. They are not found in the Great Plains region.

The oaks are of major economic importance for wood products as well as for numerous wildlife, recreation, and aesthetic uses and values in both rural and urban forests. The wood of oak species has long been known for its strength, durability, and beauty. It is widely used in products ranging from fine furniture to rough construction material, railroad cross-ties, various types of cooperage, and mine props. Acorns are a primary source of hard mast and a mainstay in the diets of many animal and bird species. The majestic oaks are known for beauty and are highly sought after for lawn and shade trees.

The oak genus (*Quercus*) is one of the eight genera of the beech family (*Fagaceae*) found worldwide. Of the estimated 500-600 species of oak, most are found in the northern hemisphere. Some 200-250 species are found in North, Central, and

South America, and 300 or more species are scattered throughout Europe, Asia, and North Africa (Little 1979, Harlow and others 1991). There are 58 recognized native oak tree species in the United States and Canada, one naturalized tree species, and about 10 native oak shrubs. There are about 125 species native to Mexico, 45 species native to Central America, five species in Colombia, and one species in Cuba (Little 1979).

For the most part, oaks are durable and relatively long-lived when compared with associated species. Ages up to 700 years have been recorded (Hora 1981). In the United States, extremely large-sized oaks have been documented. A classic example is a white oak (Q. alba L.) that was cut in 1913 near Lead Mine, West Virginia. This giant was 13 ft. in diameter 16 ft. at the base, and 10 ft. in diameter 31 ft. from the base (Clarkson 1964).

Unquestionably, oaks are a highly valued and important species. They have many important uses and values, and the forested land base on which they are growing is under extreme pressure for conversion to alternative land uses, as the nation's and the world's populations continue to grow and the demands for land-based resources continue to intensify.

OAK REGENERATION -IS IT A PROBLEM?

Whenever there is an abundance of any resource or commodity, and that resource or commodity is readily available to people who want it at a price they are willing to pay, then there is no problem with that particular resource. Oak in forests is in high demand for a variety of uses and values. But, the amount of oak that regenerates following a harvest or major disturbance is often less than was present in the parent stand. Or, forestland owners often want oak in their stands even when they did not have it in the previous stand. The mere fact that you want oak, and you do not get it, constitutes a problem regardless of the reasons for not getting it.

Two basic types of problems apply to failures in obtaining the desired level of oak regeneration. A scenario for the first type of problem is that a forest landowner or forest manager wants oak regeneration; however, an analysis of known biological and site information indicates that oak regeneration is highly unlikely. A scenario for the second type is that all of the biological and site data indicate that regeneration should occur, but it does not. Oak regeneration was an objective in both scenarios, but it did not occur; the bottom line in both cases is that it did not occur; therefore, oak regeneration is a problem. As research silviculturists, it is very important that we clearly distinguish between the two basic problem types so that viable research efforts are not misdirected.

In the first problem type, where we can clearly determine that the biotic and abiotic conditions were not conducive to oak regeneration, we need to clearly communicate this to the landowner or manager. It is quite probable that we can suggest appropriate cultural practices to alter the biotic or abiotic conditions so that the probability of obtaining oak regeneration is enhanced.

In contrast, the second problem scenario is more perplexing and problematic to silviculturists because, although the biological and site conditions indicate that successful oak regeneration should occur, it does not, indicating that factors may be coming into play that we do not understand. Only through a careful analysis and thorough evaluation of this second problem type will silviculturists be able to identify and prioritize oak regeneration research problems and subsequently devise research programs that will lead to new knowledge.

If we do not effectively convey the information we already have about oak regeneration, it is not an *oak regeneration problem*, it is a *management (people)* problem; yet it is still a problem. A discussion of oak technology transfer, however, is beyond the scope of this paper, but is being actively addressed by the sponsors and planners of this Oak Regeneration Symposium. In the remainder of this paper, I will briefly review some of the literature that deals with oak regeneration, summarize silvical characteristics of various oak species that are important in the regeneration process, and then evaluate the regeneration problem and the general level of knowledge about 31 oak species native to the United States.

SOME RESULTS OF OAK REGENERATION RESEARCH

A number of authors have identified oak regeneration as a problem in attempts to establish a new stand following harvesting. McLintock (1987) emphasized that obtaining adequate representation of preferred species, especially the oaks, is a high priority research area. Merritt (1979) provided a thorough and thoughtful review of problems associated with oak regeneration in eastern upland forests. Later in the same proceedings, McGee (1979) eluded to the sparse regeneration of northern red oak (O, rubra L.) on higher quality sites in the Southern Appalachians. Others have identified oak regeneration as a management problem in that prescribed practices are not producing the consistent oak regeneration results desired (Beck 1970, Beck and Hooper 1986, Loftis 1983, McGee and Hooper 1970). Without release from faster growing yellow-poplar, the fate of planted or natural seedlings of southern oak species in the small openings of a group selection reproduction harvest was not encouraging after 22 years (Johnson and Krinard 1983). In the semi-arid regions of California, a statewide oak regeneration survey concluded that regeneration often was inadequate to maintain existing stands and that regeneration was highly site-specific. Blue oak (Q. douglasii Hook. Arn.) and valley oak (Q. lobata Née) regeneration tended to be poor (Muick and Bartolome 1987). In another study, Muick (1991) observed that coast live oak (Q. agrifolia Née), interior live oak (Q. wislizeni A. DC.) and canyon live oak (Q. chrysolepis Liebm.) were regenerating more successfully. Certainly, not all reports of oak regeneration indicate failures; there are successes. Johnson and Krinard (1988) reported that 29 years after harvest, cherrybark oak (Q. falcata var. pagodifolia Ell.) was a significant component in a southern bottomland hardwood stand. Earlier evaluations of the stand development had been less optimistic as far as the oak component was concerned.

Clark and Watt (1971) identified two basic principles that must be understood when discussing natural regeneration of upland oaks and that probably also apply to bottomland oaks and to some degree to the Western United States oaks as well:

- 1. The new stand will contain oak in proportion to the advance oak reproduction on the area before the harvest cut.
- 2. Advance oak reproduction must have a well-established root system to compete successfully with other woody vegetation (and with grasses in many Western United States oak sites).

The need to understand these two very basic principles has not changed in the past two decades. This points to a need to understand as much as we can about how oak regeneration occurs and how it gets to the stage of advance reproduction with the associated well-established root system. In other words, what are the important silvical characteristics of oak species that are important in reproduction establishment and subsequent growth and development?

DAK REGENERATION: IMPLICATIONS OF SILVICAL CHARACTERISTICS A review of some of the more important regeneration-related silvical characteristics of oak species provides insights into possible reasons for low regeneration success or for complete failures (table 1). For the most part, eastern upland and bottomland oaks occur in association with a wide array of overstory and understory woody species. Seldom will oak occur in pure stands. On the more arid sites in the West, oak will often occur in clumps or small, pure stands; however, competition from grasses and shrubs is often severe and greatly influences regeneration establishment and early growth.

> The preponderance of oak reproduction results from natural regeneration following a harvest or natural disturbance. Planting, although very limited in application, has enjoyed a wide range of success ranging from complete failures to fully stocked plantations. However, the artificial regeneration process usually requires intense culture, including competition control for one or more growing seasons and is only accomplished at a relatively high cost. Natural regeneration usually is composed of three different reproductive forms: seedlings, seedling sprouts (normally considered as advance regeneration), and stump sprouts. Of the three reproductive forms, seedlings are the slowest growing and least competitive, and stump sprouts are generally the fastest growing and most competitive. Seedling sprouts are unique because their root systems are usually well-developed compared to a seedling root; however, the top often resembles that of seedlings.

> Of the 31 species reviewed in this paper, three are considered tolerant to shade, 11 are intermediate in tolerance to shade, and 17 are intolerant or very intolerant (table 1). None of the bottomland oaks tolerate flooding to any significant degree, especially flooding that occurs during the growing season (table 1). In discussions with colleagues active in bottomland oak regeneration research, there is a strong consensus that flooding is particularly detrimental to newly established seedlings. In contrast to shade and flood tolerance, however, the oaks that inhabit exposed, droughty sites appear to be quite tolerant of low moisture conditions (table 1). In general, regeneration success is relatively good on poorer sites where competition from associated species is often less intense.

> Juvenile growth of oak species is generally slower than or equal to associated competition. Sixteen of the species reviewed grew slower than competitors, eight grew about as fast, and only five had juvenile growth rates that were faster than associated competitors (table 1). Acorn production of oak species is sporadic and unpredictable at best. In general, bumper acorn crops are required to obtain significant seedling establishment. Predation by wildlife and insects coupled with acorn mortality from disease and adverse weather conditions during the intervening years between bumper crops are believed to restrict greatly the acorns that are available to germinate and develop as regeneration. The analysis of available oak regeneration research information suggests that high levels of acorn production generally occur at intervals ranging from 3 to 7 years or more (table 1).

Table 1—Summary of regeneration-related silvical characteristics¹ of 31 oak species native to the United States, noting important uses, values, and relative assessment of the available research knowledge

Scientific Name (common name)	Tolerance			Acorn Production (years)		Juvenile Growth	Uses and	Research Knowl-
	Shade ²	Flood ³	Drought ³	Normal	High	Rate ⁴	Values ⁵	edge ⁶
		W1.4. 0						
Q. agrifolia Née (coast live oak)	Tol ⁷	Subgenus: N/A	Leucobalanus Inte	: White O Unk	Unk Unk	slower	W,O,F,S	Lim
Q. alba L. (white oak)	Inte	N/A	Intol	2-4	4-10	slower	W,O,F, T,S	Ext
Q. bicolor Willd. (swamp white oak)	Inte	Tol	N/A	2-4	3-5	slower	W,O,F,T	Mod
Q. garryana Dougl. ex Hook. (Oregon white oak)	Inte	Tol	Tol	Unk	3-5	slower	W,O,F, T,S	Mod
Q. lyrata Walt. (overcup oak)	Inte	Tol	N/A	1-2	3-4	compet- itive	W,F,T	Lim
Q. macrocarpa Michx. (bur oak)	Inte	N/A	Tol	2-4	2-3	faster	W,F,T,S	Lim
Q. michauxii Nutt. (swamp chestnut oak)	Intol ⁸ Inte	Inte	N/A	2-4	3-5	compet- itive	W,F,T	Mod
Q. prinus L. (chestnut oak)	Inte	N/A	Tol	erratic	4-5	slower	W,F,T,S	Mod
Q. stellata Wangenh. (post oak)	Intol	N/A	Tol	2-4	2-3	slower	W,O,F, T,S	Lim
Q. virginiana Mill. (live oak)	Inte	Inte	Inte	Unk	Unk	Unk	W,O,F,S	Lim
Q. muehlenbergii Engelm. (chinkapin oak)	Intol	N/A	Tol	Unk	Infreq.	slower	W,F,S	Lim
Q. lobata Née (valley oak)	Intol ⁹	Intol	N/A	Unk	2-3	slower	W,O,F	Lim
Q. coccinea Muenchh. (scarlet oak)	Subg V. Inte		h robalanus: R o Inte- Tol	ed and Bla erratic	ck Oaks 3-5	faster	W,O,F, T,S	Mod
Q. douglasii Hook & Arn. (blue oak)	Intol	N/A	Tol	2-3	5-8	slower	W,F,S	Lim
Q. ellipsoidalis E. J. Hill (northern pin oak)	Intol ¹⁰	N/A	Tol	2-4	3-5	compet- itive	W,O,F,S	Lim
Q. falcata Michx. (southern red oak)	Inte	N/A	Tol	Unk	Unk	compet- itive	W,O,F, T,S	Mod
Q. pagoda Raf. (cherrybark oak)	Intol	Inte	N/A	1-2	2-4	faster	W,O,F, T,S	Mod- Ext
Q. ilicifolia Wangenh. (bear oak)	V. Intol ¹⁰	N/A	Tol	Unk	Unk	slower	W,F,S,	Lim

Scientific Name (common name) S		Tolerance - Flood ³		Acorn Production (years)		Juvenile Growth	Uses and	Research Knowl-
	hade ²		Drought ³	Normal	High	Rate ⁴	Values ⁵	edge ⁶
Q. imbricaria Michx. (shingle oak)	Intol	Inte- Tol	Intol	Unk	Unk	compet- itive	W,O,F, T,S	Lim
Q. kelloggii Newb. (California black oak)	Intol	Intol	Intol	2-4	3-5	slower	W,O,F, T,S	Lim
Q. laevis Walt. (turkey oak)	Intol	N/A	Tol	1-2	2-3	compet- itive	W,F,S	Lim
Q. laurifolia Michx. (swamp laurel oak, diamondleaf	Inte- f) Tol	Tol	N/A	Unk	Unk	Unk	W,O,F,	Lim
Q. marilandica Muenchh. (blackjack oak)	Intol ¹⁰	N/A	Tol	Unk	Unk	slower	W,F,S,T	Lim
Q. nigra L. (water oak)	Intol	Inte	N/A	1-2	1-2	slower	W,O,F, T,S	Lim
Q. nuttallii Palmer (Nuttal oak)	Intol Inte	Inte- Tol	N/A	2-4	3-4	faster	W,O,F,T	Lim
Q. palustris Muenchh. (pin oak)	Intol	Inte- Tol	N/A	2-4	4-6	faster	W,O,F,T	Lim
Q. phellos L. (willow oak)	Intol	Intol- Inte	Intol	1-2	1-2	compet- itive	W,O,F,T	Lim
Q. rubra L. (northern red oak)	Inte	N/A	Intol	2-4	2-5	slower	W,O,F, T,S	Ext
Q. shumardii Buckl. (Shumard oak)	Intol	Intol	Intol	2-4	Unk	compet- itive	W,O,F,T	Lim
Q. velutina Lam. (black oak)	Inte	N/A	Inte	2-4	2-3	slower	W,F,T,S	Mod
	Subį	genus: Proto	obalanus: 1	ntermediate	e Oaks			
Q. chrysolepis Liebm. (canyon live oak)	Tol	N/A	Inte	2-4	2-4	slower	W,O,F, T,S	Lim

Table 1—Summary of regeneration-related silvical characteristics¹ of 31 oak species native to the United States, noting important uses, values, and relative assessment of the available research knowledge (Continued)

¹ Unless otherwise indicated, the information is derived from Burns and Honkala 1990; Little 1979; Harlow and others 1991; Elias 1980.

² Five levels of shade tolerance: V. Tol = very tolerant; Tol = tolerant; Inte = intermediate; Intol = intolerant; V. Intol = very intolerant.

³ Three levels of tolerance: Tol = tolerant; Inte = intermediate; Intol = intolerant; N/A = not applicable.

⁴ Growth rate relative to associated species: slower, competitive, faster.

⁵ Uses and values are W = wildlife; O = ornamental; F = fuel; T = timber/fiber; S = site protection/watershed.

⁶ Ext = extensive; Mod = moderate; Lim = limited.

⁷ Muick 1991.

⁸ Johnson and Krinard 1983.

⁹ Standiford 1991.

¹⁰ Author's field observations.

SUMMARY AND CONCLUSIONS

Oak is the largest tree genus in the United States and is the most important hardwood. The oaks enjoy a full array of values and uses, and the demand for these values and uses will only increase in the future. In terms of the level of knowledge available about oak regeneration, we are at a distinct disadvantage. We have a significant pool of knowledge for only three of the 31 species reviewed. We have a modest amount of regeneration information for seven species and only limited information on the remaining 21 species (table 1). For the 27 native oak species that were not reviewed, there is virtually no information available.

The regeneration-related silvical characteristics of the 31 oak species, as a group, indicate that the oaks:

- 1. Are intolerant or intermediate in tolerance to shade
- 2. Are intolerant of flooding or inundation
- 3. Tend to be drought tolerant
- 4. Grow at rates slower than or equal to associated competitors
- 5. Produce acorns at sporadic and unpredictable intervals with bumper crops being produced at 3- to 7-year intervals.

Based on the above summary, there is little question that successful natural oak regeneration is difficult to achieve consistently without silvicultural intervention. Even with silvicultural intervention, the regeneration results will be quite variable because of unpredictable and uncontrollable environmental conditions and a limited amount of knowledge about the processes and interactions associated with oak regeneration (table 2).

Oak regeneration is a problem and the problem is widespread. Many of the problems can be solved by utilizing information that is already available, but there is a cost involved, which will have to be addressed by forest managers and forest landowners. Other problems with oak regeneration will require a major research commitment.

What we learn at this Oak Regeneration Symposium will clarify our vision of the most important problem areas and allow us to initiate research programs to provide much-needed answers.

Species/ Species Group	Site Quality ¹	Regeneration Origin (Probability of Occurrence) ²		Overall Regeneration Problem ³	
		Geedline		SEVERE	
	High	Seedling Seedling sprout	(Med) (Med)	SEVERE	
	mgn	Stump sprout	(Low)	SEVERE	
White Oak		Stump sprout	(LOW)	SEVERE	
Winte Out		Seedling	(High)	SEVERE	
	Low	Seedling sprout	(High)	MODERATE	
		Stump sprout	(High)	MINIMAL	
Chestnut Oak	High Low	Seedling Seedling sprout Stump sprout Seedling Seedling sprout Stump sprout	(Med) (Med) (Med) (High) (High) (High)	SEVERE MODERATE-SEV MODERATE MODERATE-SEV MODERATE MINIMAL	
Bur Oak Post Oak	High	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A	
Chinkapin Oak	Low	Seedling Seedling sprout Stump sprout	(Med) (High) (High)	MODERATE-SEV MODERATE-MIN MINIMAL	

Table 2a.—Evaluation of regeneration for Eastern United States upland white oak species/species groups as a function of site quality and regeneration origin.

¹ High = site index₅₀ 75 and greater; Low = site index₅₀ less than 75.

² High = high probability of regeneration of specified type occuring on specified site; Med = medium probability; Low = low probability; N/A = not applicable, species do not occur on site indicated.

Species/ Species Group	Site Quality ¹	Regeneration Origin (Probability of Occurrence) ²		Overall Regeneration Problem ³	
		Gaadling		OPVEDE	
	High	Seedling	(Med) (Med)	SEVERE MODERATE-SEV	
	nign	Seedling sprout Stump sprout	(Low)	MODERATE-SEV	
Swamp Oak		Stump sprout	(LOW)	MODERATE	
Swamp Chestnut Oak		Seedling	(N/A)	N/A	
I	Low	Seedling sprout	(N/A)	N/A	
		Stump sprout	(N/A)	N/A	
Overcup Oak	High Low	Seedling Seedling sprout Stump sprout Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A) (Med) (Med) (Med)	N/A N/A N/A MODERATE MODERATE MODERATE	
	High	Seedling Seedling sprout Stump sprout	(Med) (Low) (High)	N/A N/A N/A	
Live Oak		Seedling	(Mad)	MODERATE	
	Low	Seedling sprout	(Med) (Med)	MODERATE-MIN	
	20 W	Stump sprout	(High)	MINIMAL	
		Root sprouts	(High)	MINIMAL	

Table 2b.—Evaluation of regeneration for Eastern United States bottomland white oak species/species groups as a function of site quality and regeneration origin.

¹ High = site index₅₀ 75 and greater; Low = site index₅₀ less than 75.

² High = high probability of regeneration of specified type occuring on specified site; Med = medium probability; Low = low probability; N/A = not applicable, species do not occur on site indicated.

Species/ Site Species Group Quality ¹		Regeneration Origin (Probability of Occurrence) ²		Overall Regeneration Problem ³	
Northern Red Oak Black Oak	High	Seedling Seedling sprout Stump sprout	(Med) (Med) (Low)	SEVERE SEVERE-MOD MODERATE-SEV	
Diack Uak	Low	Seedling Seedling sprout Stump sprout	(Med) (Med) (High)	SEVERE MODERATE MINIMAL	
Scarlet Oak	High	Seedling Seedling sprout Stump sprout	(Low) (Med) (Low)	SEVERE SEVERE-MOD MODERATE-SEV	
Southern Red Oak Northern Pin Oak	Low	Seedling Seedling sprout Stump sprout	(Med) (Med) (High)	SEVERE MODERATE-MIN MINIMAL	
Turkey Oak	High	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A	
Shingle Oak Blackjack Oak	Low	Seedling Seedling sprout Stump sprout	(Med) (Med) (High)	SEVERE MODERATE MINIMAL	
	High	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A	
Bear Oak	Low	Seedling Seedling sprout Stump sprout	(High) (Low) (High)	MODERATE-SEV MODERATE MODERATE	

Table 2c.—Evaluation of regeneration for Eastern United States upland red oak species/species groups as a function of site quality and regeneration origin.

¹ High = site index₅₀ 75 and greater; Low = site index₅₀ less than 75.

² High = high probability of regeneration of specified type occuring on specified site; Med = medium probability; Low = low probability; N/A = not applicable, species do not occur on site indicated.

Species/ Species Group	Site Quality ¹	Regeneration Origin (Probability of Occurrence) ²		Overall Regeneration Problem ³	
Cherrybark Oak	High	Seedling Seedling sprout Stump sprout	(Med) (Med) (Low)	SEVERE-MOD MODERATE SEVERE	
Shumard Oak	Low	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A	
Water Oak	High	Seedling Seedling sprout Stump sprout	(Med) (Med) (Med)	SEVERE SEVERE-MOD MODERATE-SEV	
Willow Oak Nuttall Oak	Low	Seedling Seedling sprout Stump sprout	(Med) (Med) (High)	SEVERE MODERATE MINIMAL	
Swamp Laurel Oak or	High	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A	
Diamondleaf Pin Oak	Low	Seedling Seedling sprout Stump sprout	(Med) (Med) (High)	MODERATE MODERATE-MIN MINIMAL	

Table 2d.—Evaluation of regeneration for Eastern United States bottomland red oak species/species groups as a function of site quality and regeneration origin.

¹ High = site index₅₀ 75 and greater; Low = site index₅₀ less than 75.

² High = high probability of regeneration of specified type occuring on specified site; Med = medium probability; Low = low probability; N/A = not applicable, species do not occur on site indicated.

Species/ Species Group	Site Quality ¹	Regeneration O (Probability of (Overall Regeneration Problem ³
Coast Live Oak	High	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A
Oregon White Oak	Low	Seedling Seedling sprout Stump sprout	(Med) (Unknown) (High)	MODERATE-SEV UNKNOWN MINIMAL
Valley Oak	High	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A
Blue Oak	Low	Seedling Seedling sprout Stump sprout	(Med) (Unknown) (Med)	SEVERE UNKNOWN MODERATE
California Black Oak	High	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A
	Low	Seedling Seedling sprout Stump sprout	(Low) (Unknown) (High)	SEVERE UNKNOWN MINIMAL
Canyon Live Oak	High	Seedling Seedling sprout Stump sprout	(N/A) (N/A) (N/A)	N/A N/A N/A
	Low	Seedling Seedling sprout Stump sprout	(High) (Unknown) (High)	MODERATE UNKNOWN MINIMAL

Table 2e.—Evaluation of regeneration for Western United States oak species/species groups

 (subgenuses combined) as a function of site quality and regeneration origin.

¹ High = site index₅₀ 75 and greater; Low = site index₅₀ less than 75.

² High = high probability of regeneration of specified type occuring on specified site; Med = medium probability; Low = low probability; N/A = not applicable, species do not occur on site indicated.

- LITERATURE CITED
- Beck, D.E. 1970. Effect of competition on survival and height growth of red oak. Res. Pap. SE-56, Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 7 pp.
- Beck, D.E.; Hooper, R. M. 1986. Development of southern Appalachian hardwood clearcutting. Southern Journal of Applied Forestry. 10(3):168-172.
- Burns, R.M.; Honkala, B.H., tech. coords. 1990. Silvics of North America: 2 hardwoods. Agriculture Handbook 654. Washington, D.C.: U.S. Department of Agriculture, Forest Service, Vol. 2. 877 pp.
- Clark, F.B.; Watt, R.F. 1971. Silvicultural methods for regeneration oaks. In: White, D.E.; Roach, B.A., eds. Oak symposium proceedings. Upper Darby, PN: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 37-42.

Clarkson, R.B. 1964. Tumult on the mountain. Parsons, WV: The McClain Printing Co. 410 pp.

- Elias, T.S. 1980. The complete trees of North America. New York, NY: Times Mirror Magazines, Inc. 948 pp.
- Harlow, W.M.; Harrar, E.S.; Hardin, J.W.; White, F.M. 1991. Textbook of dendrology: covering the important forest trees of the United States and Canada. 7th edition. New York, NY: McGraw-Hill, Inc. 501 pp.
- Hora, B., consult. ed. 1981. The Oxford encyclopedia of trees of the world. New York, NY: Oxford University Press. 288 pp.
- Johnson, R.L.; Krinard, R.M. 1983. Development of seven hardwood species in small forest openings 22-year results. Southern Journal of Applied Forestry. 7(3):153-156.
- Johnson, R.L.; Krinard, R.M. 1988. Growth and development of two sweetgum-red oak stands from origin through 29 years. Southern Journal of Applied Forestry. 12(2):73-78.
- Little, E.L., Jr., 1979. Checklist of United States trees (native and naturalized). U.S. Department of Agriculture Handbook 541. U.S. Government Printing Office, Washington, D.C. 375 pp.
- Loftis, D.L. 1983. Regenerating southern Appalachians mixed hardwoods with the shelterwood method. Southern Journal of Applied Forestry. 7(4):212-217.
- McGee, C.E. 1979. Fire and other factors related to oak regeneration. In: Holt, H.A.; Fisher, B.C., eds. Regenerating oaks in upland hardwood forests, 1979 John S. Wright Forestry Conference. West Lafayette, IN: Purdue University: 75-81.
- McGee, C.E.; Hooper, R.M. 1970. Regeneration after clearcutting in the southern Appalachians. Res. Pap. SE-70, Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 12 pp.
- McLintock, T.F. 1987. Research priorities for eastern hardwoods. Hardwood Research Council, Memphis, Tennessee. 103 pp.
- Merritt, C. 1979. An overview of oak regeneration problems. In: Holt, H.A.; Fisher, B.C., eds. Regenerating oaks in upland hardwood forests, 1979 John S. Wright Forestry Conference. West Lafayette, IN: Purdue University: 1-9.
- Muick, P.C. 1991. Effects of shade on blue oak and coast live oak regeneration in California annual grass lands. In: Standiford, R.B., tech. coord. 1991. Proceedings of the symposium on oak woodlands and hardwood rangeland management, 1990 October 31-November 2; Davis, California. Gen. Tech. Rep. PSW-126. Berkeley, California: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 21-24.
- Muick, P.C.; Bartolome, J.W. 1987. An assessment of natural regeneration of oaks in California. Report to California Department of Forestry, Forest and Rangeland assessment programs, Sacramento, California. 100 pp.

The Biology of Oak Regeneration

u

Ecology and Physiology of Oak Regeneration

John D. Hodges and Emile S. Gardiner, Department of Forestry, Mississippi State University, Mississippi State, MS 59762

ABSTRACT Obtaining adequate oak regeneration in hardwood stands is a problem complicated by the lack of a lucid understanding of species biology. Under heavy shade, relatively little morphological acclimation, low photosynthetic capacity, and high respiration contribute to an overall low capacity to compete with tolerant species. When released, oaks usually do not respond as quickly as their intolerant competitors, because of relatively lower compatibility with an open environment. Yet, conservative growth characteristics like early root growth and deep root systems, as well as physiological characteristics such as high water use efficiency and stomatal closure only at very low water potentials, enable oaks to compete more successfully on drier sites. Silvicultural practices aimed at establishing oak regeneration must enhance environmental conditions complimentary to oak biology.

INTRODUCTION Other papers in this symposium address the many problems associated with oak regeneration and the silvicultural and management techniques that have been or could be used to alleviate those problems. It is obvious that much progress has been made, but there are unanswered questions and problems to be solved. Two of the major unresolved problems are: (1) Absence of adequate advance regeneration and knowledge of methods for obtaining it, and (2) Slow juvenile growth rate of oaks and their slow response to release (Hodges 1987, Crow 1988). Resolution of such problems requires an understanding of the biology of the species.

This paper will examine the current state of knowledge regarding the ecology and physiology of oak seedlings and relate that information to oak regeneration. A recent paper by Kolb and others (1990) presented results for northern red oak (*Quercus rubra* L.) and an excellent review of oak growth response to resource limitation. They discussed the growth strategy of red oak in relation to ecological requirements for regeneration. We will review much of that material, but the emphasis will be on physiological and morphological adaptations as they relate to regeneration. A basic assumption we have made is that regeneration and subsequent composition of a forest community is determined by: (1) microclimate and edaphic factors, (2) morphological and physiological characteristics of the species present, and (3) interaction between the two.

The relationship between site quality and oak regeneration has been extensively reviewed by other papers in this symposium. In general, obtaining oak regeneration is not a problem, or far less of a problem, on drier or below-average sites than on good sites (Arend and Gysel 1952, Loftis 1988, Hodges unpublished). Therefore, we will primarily discuss oak regeneration on the better upland and bottomland sites. We will consider ecological and physiological adaptations and how they act to make the species or plant more compatible with its environment and thus influence oak regeneration. There are differences between oak species, but most research has been on species of greater occurrence and commercial importance such as northern red oak, white oak (Q. alba L.), and cherrybark oak (Q. pagoda Raf.). That research will be emphasized, but exceptions will be noted.

NATURAL REGENERATION Oaks can be regenerated by natural and artificial means. Some problems encountered are common to both regeneration methods while others are not. We will consider the two methods separately by examining ecological and physiological factors as they relate to the regeneration process and the problems involved.

Germination Acorn production is highly erratic from year to year, and losses due to insects, birds, and mammals can be very high (see other papers in this symposium for references and detailed information on acorn production, acorn losses, storage, and acorn germination). Once acorns are on the ground, deterioration can be quite rapid, usually due to desiccation but also to excessive heat or cold. Also, on bottomland sites flood waters can wash a site virtually clean of acorns. In any event, it is extremely rare for acorns to survive past the first growing season after dispersal.

Despite these losses, mast production over time generally provides sufficient acorns so that germination *per se* is seldom a limiting factor for seedling establishment on most sites (see paper by Lorimer, this symposium). Burial by rodents and birds and coverage by sediment and debris help assure favorable conditions for acorn germination (Darley-Hill and Johnson 1981, Crow 1988, Deen and Hodges 1990). On upland and bottomland sites containing mature oaks, it is not uncommon to find 12,000 or more new germinants per hectare beneath an existing stand, even in deep shade (Hodges unpublished, Carvell and Tryon 1961, Beck and Hooper 1986, Merritt and Pope 1991). However, on mesic upland and floodplain sites, few of these seedlings will survive to the second growing season.

Survival and First-year Growth Growth Growth Factors influencing early survival and growth of seedlings beneath a stand may be very different from those determining growth of older seedlings. Initial survival and early growth of oak seedlings beneath a dense canopy are primarily dependent on stored food reserves in the acorns and not on current photosynthate production (Wassink and Richardson 1951, Grime and Jeffrey 1965, Musselman and Gatherum 1969, Crow 1988). Under low light, first-year oak seedlings usually produce only one growth flush (Phares 1971, Crow 1988), which is determined by stored food in the acorns (Richardson 1956). Once cotyledon reserves are depleted, seedlings must survive on photosynthate produced by new leaves. Light then becomes a limiting factor for survival and growth (Tubbs 1977, Hanson 1986, Crow 1988). In fact, the inability to maintain a positive carbon balance in low light beneath Page 114 of 404

mature stands may be the most likely reason for lack of oak advance regeneration in those stands (Musselman and Gatherum 1969). At low light intensities, CO_2 uptake and photosynthate production may be restricted by a lack of energy for photochemical reactions, and by stomatal resistance to CO_2 diffusion. Wuenscher and Kozlowski (1970) found that net photosynthesis of some oak species was not saturated until light intensity was high enough to cause complete stomatal opening.

Light intensity near the floor of hardwood stands is often at or below the compensation point for oaks. This low light intensity is often due not so much to a dense main canopy as to a nearly continuous mid-canopy or multi-storied layer of tolerant vegetation below the main canopy (Janzen and Hodges 1985, Hodges 1987, Janzen and Hodges 1987, Pubanz and Lorimer 1992). For northern red oak in the first growth flush, Hanson (1986) found that the level of photosynthetically active radiation needed for a positive carbon balance was about 30 μ mol m² s⁻¹. Light levels in dense northern hardwood stands (Hanson 1986, Crow 1988, Pubanz and Lorimer 1992) and southern bottomland hardwood stands (Lockhart 1992, Hodges unpublished) are often below that level, so carbohydrate reserves used in respiration exceed that produced by photosynthesis, and the new seedlings eventually die.

Although light is a dominant environmental factor limiting seedling establishment under dense canopies, soil moisture may also limit establishment. Root growth of oaks, especially white oaks, begins, and may be quite extensive, before the shoot emerges. Root regeneration and root growth are very sensitive to soil moisture stress (Larson and Whitmore 1970, Larson 1980). In northern red oak, root initiation and growth ceased at soil osmotic potentials between -0.4 and -0.6 MPa (Larson and Whitmore 1970). Even on floodplain sites, greater seedling densities and survival may occur on the wetter micro-sites (Jones and Sharitz 1990). However, cherrybark oak seems less sensitive to root competition than many of its competitors (Jones and others 1989).

Juvenile Growth Phase Numerous studies have shown that successful natural regeneration of eastern oaks requires the presence of advance regeneration of a number and size which can compete successfully with other species (Beck 1970, Sander 1972, Johnson 1975, Loftis 1983). Large numbers of oak seedlings are common on good sites, but they are typically small (<30 cm tall) and incapable of rapid height growth after release (Beck 1970, Sander 1972, Johnson 1979, Janzen and Hodges 1987). Johnson (1975) found that Nuttall oak (*Q. nuttallii* Palmer) seedlings can become established beneath a stand and survive for many years with as little as 2 hours sunlight daily, but height growth was extremely slow. Partial cuttings of various types (thinning, improvement, shelterwood, single-tree selection) may greatly increase stocking of oaks, but these practices do not assure growth to a size necessary for competing successfully with associated species (Loftis 1979, 1983; Hill and Dickmann 1988; Martin and Hix 1988).

> The above findings appear anomalous in light of the many studies which have shown photosynthesis is saturated and maximum growth of oak seedlings occurs at 30-50 percent of full sunlight (Kramer and Decker 1944, Bourdeau 1954, Loach 1967, Musselman and Gatherum 1969, Phares 1971, Hodges 1987). For example, maximum biomass growth of cherrybark oak occurred at 27-53 percent of full sunlight, but height growth at 8 percent was as good as at full sunlight (table 1). Page 115 of 404

What then is the reason for the slow juvenile growth rate of oak seedlings? The answer may reside in the ecological and physiological differences between oaks and the species with which they must compete for survival and dominance of the site (Kolb and others 1990). Often the main competition is between oak seedlings and a midstory and understory of more tolerant species (Beck 1970, Sander 1972, Janzen and Hodges 1985, Beck and Hooper 1986, Hodges 1987).

Table 1—Influence of shade on growth and biomass partitioning in seedlings of cherrybark oak¹

Sun	Year 1			Year 2				
(percent)	Height	Stem	Root	R/S ²	Height	Stem	Root	R/S ²
	(cm)		- (g)		(cm)		(g)	
1 00	22.0	1.59	5.05	3.18	49.8	10.88	25.75	2.37
53	32.6	3.19	8.98	2.82	117.4	77.94	96.44	1.24
27	29.7	1.46	3.68	2.52	94.5	47.60	52.92	1.11
8	22.1	.78	1.98	2.53	58.5	9.48	14.98	1.58

¹ Seedlings were grown in shade houses covered with saran shade cloth. Soil moisture was maintained at or near field capacity in all plots. These unpublished data are averages for two studies replicated in time at the same location.

² Root/shoot ratio.

Shade tolerance is the ability of a species to persist in a low light environment. Kramer and Kozlowski (1979) defined shade tolerant species as those which reach a maximum rate of net photosynthesis at relatively low light intensities. Oak species exhibit a range in tolerance from moderately tolerant to intolerant, with white oaks generally more tolerant than red oaks (McQuilkin 1975, McGee 1981). Woody species with which oaks may compete during their life cycle vary in tolerance from very tolerant, for those occurring beneath a mature stand, to intolerant, for those growing only under more open conditions.

Mechanisms of shade tolerance are related to morphological and physiological adaptations. Some authors have suggested that there is a shade-tolerant growth form (morphology) which permits long-term survival of seedlings (Kohyama 1980, Hara 1987). Coupled with this species-specific growth form may be differences in phenology, photosynthesis, respiration, water use efficiency, and growth rates which permit the species to survive and grow under shade conditions (Grime and Jeffrey 1965, Loach 1970, Matsuda and McBride 1986, Matsuda and others 1989).

McGee (1975, 1986) reported that most tolerant species began budbreak earlier under forest canopies than in open clearcuts. This observation applied to six oak species as well. He suggested that the propensity of some species to initiate growth early under a canopy increases their shade tolerance by allowing some leaf development before crown closure.

Morphological acclimations in plants are thought to occur in response to limited resources (Schulze and others 1983, Johnson and Thornley 1987). When grown under shade, seedlings of most woody species will partition their growth to favor leaf growth and/or leaf area ratio as well as shoot/root ratio (Musselman and Gatherum 1969; Kolb and Steiner 1990a, 1990b), which should favor more effective utilization of the limited light resource. Oaks follow this pattern (figure

1, table 1), but the greater partitioning to shoot growth may not occur in the first growing season (Gottschalk 1987, Kolb and Steiner 1990a, Hodges present paper).

Although these morphological changes should improve the light-capturing ability of oaks, they do not seem to improve growth relative to tolerant understory species (Loach 1970). In fact, morphological acclimations of this type seem to be more important for intolerant species such as yellow-poplar (*Liriodendron tulipifera* L.) and American sycamore (*Platanus occidentalis* L.) than for oaks (Loach 1970, Jones and others 1989, Kolb and Steiner 1990a).




Figure 1—Influence of shade on root/shoot dry weight ratios of 2-year-old cherrybark oak seedlings grown in shade houses. Leaves are not included with shoots. Studies were replicated in time at the same location. (Unpublished data, Mississippi State University.)

Physiological acclimations to shade occur within and between species. Studies with oaks have shown acclimations in shade foliage that, when compared with sun foliage, include higher rates of net photosynthesis at low light levels, lower respiration rates, and higher levels of chlorophyll per unit of leaf area (Loach 1967). However, when compared with more tolerant species at low light intensities, oaks generally have: (1) a higher light compensation point for photosynthesis and less efficient use of "light flecks" or short bursts of light (Loach 1967, Woods and Turner 1971, Teskey and Shrestha 1985); (2) about the same or lower rates of net photosynthesis (Grime and Jeffrey 1965, Wuenscher and Kozlowski 1970, Farmer 1980); (3) as high or higher rates of respiration (Went 1957; Loach 1967; Wuenscher and Kozlowski 1970, 1971; Teskey and Shrestha 1985; Collier and others 1992); (4) slower stomatal opening and/or greater stomatal resistance to CO₂ diffusion (Woods and Turner 1971, Wuenscher and Kozlowski 1971, Davies and Kozlowski 1974, Teskey and Shrestha 1985); (5) lower quantum yield (Teskey and Shrestha 1985); (6) higher water use efficiency (Wuenscher and Kozlowski 1971, Teskey and Shrestha 1985); and (7) higher saturating light

Page 117 of 404

intensity for photosynthesis (Wuenscher and Kozlowski 1970, Bazzaz and Carlson 1982, Teskey and Shrestha 1985). These differences in morphological and physiological characteristics explain why oaks generally do not compete successfully in the understory on good sites.

Response of Seedlings to Release Numerous studies have shown that large numbers of advance oak regeneration will not necessarily assure acceptable oak regeneration even if released by complete clearcutting (Beck 1970; Sander and Clark 1971; Sander 1972; Johnson 1979; Loftis 1983, 1988, 1990; Janzen and Hodges 1987). One reason is that, even though the overstory is removed, the oaks may still have to compete with a dense former understory of larger and usually more tolerant seedlings, saplings, and sprouts (Johnson and Jacobs 1981; Janzen and Hodges 1985; Beck and Hooper 1986; Hix and Lorimer 1990, 1991). These species usually have well-developed root systems and ample foliage which enable them to respond faster to release than oak seedlings.

> Another, and likely more important, reason for oak regeneration failure is the slow growth rate of recently released seedlings (Beck 1970, Sander 1972, McQuilkin 1975, Janzen and Hodges 1987, Hix and Lorimer 1990). Northern red oak (Dickson 1991), cherrybark oak (Hodges and Janzen 1987) and probably most other oaks, have a conservative growth strategy in which photosynthate resources of the young seedlings are devoted first to building a root system. If, at the time of release, the seedling does not have a large root system and adequate shoot height (about 1.2 m for northern red oak) (Sander 1972), shoot growth will be slow until the root system develops.

> Physiologically, oaks are not as compatible with conditions in an open environment as are many of their competitors, especially the intolerant species. Bazzaz and Carlson (1982) found physiological flexibility to be much greater in early successional species than in late successional species. Early successional species were found to be capable of responding to light so that they become more like shade plants when grown in the shade, but late successional species were unable to make the converse switch when grown in a high-light environment. Photosynthetic rates for yellow-poplar are higher than for northern red oak except at very low light intensities, yet when released the rates are much higher at high light intensities. Also, photosynthesis in oak is saturated at a much lower light intensity (Loach 1967).

> Work by McGee (1975, 1986, 1988) suggests yet another possible reason for the slow growth of oak seedlings following release. He observed that budbreak in several oak species when grown under shade was earlier than when grown in the open. Thus, early budbreak of the released seedlings will increase the likelihood of freeze damage and retarded development.

Coppice Regeneration Sprouts are an extremely important source of regeneration for most oaks, especially on xeric sites. Paul Johnson (this symposium) gives an extensive discussion of sprouts, their origin, and growth rates. Sprouts can originate in several ways, but the form that is most important for eastern oak regeneration is stump sprouts, that include seedling sprouts. These sprouts originate from dormant buds at or near the root collar. As long as the crown of the tree or seedling is attached, these buds are

suppressed by growth regulators produced in the crown. When the tree or seedling is severed, the resulting sprouts show very rapid height growth. They may produce four or more flushes per year and more than 1 m of growth (Johnson 1979, Reich and others 1980).

Physiological reasons for the more rapid growth of stump sprouts is not fully understood, but in the case of larger stumps it certainly involves greater carbohydrate reserves and better absorption of water and nutrients because of the large root system. In the case of seedling sprouts, it apparently also involves a shift in the allocation of resources to favor shoot growth, an increase in stomatal conductance and carbon dioxide exchange rates, and/or increased hydraulic conductivity of the stem as opposed to intact seedlings.

Severing seedlings at time of release has been shown to partially alleviate the slow response of intact seedlings (figure 2) (Janzen and Hodges 1987, Kruger and Reich 1989, Lockhart 1992). Lockhart (1992) found that sprout shoots of cherrybark oak were the greatest sink for recently produced photosynthates and stored reserves. This change in allocation pattern allowed sprouts to maintain top growth over a longer period of time compared to intact seedlings. Sprout leaves of northern red oak (Kruger and Reich 1989) and cherrybark oak (Lockhart 1992) showed increased stomatal conductance over those from intact seedlings. In northern red oak there was also an increase in carbon dioxide exchange rate, but this was not observed in cherrybark oak.

Vessel size is known to influence water transport through the stem (Abrams 1990). The vascular system of seedling sprouts may be quite different from that of intact seedlings (Blake and Tschaplionski 1986) and may permit faster movement of water through the stem, thus in lower moisture deficits in seedlings.




Figure 2—Height growth of bottomland oaks after removal of competing species in the midstory and understory by herbicidal injection. In the inject + clip treatment, tops of all oak seedlings were cut 2.5 cm above groundline. The asterisk indicates initial height of clipped seedlings (Janzen and Hodges 1987).

Other papers in this symposium emphasize the importance of site quality in Site Quality and Disturbance determining the ease of obtaining oak regeneration and the role of disturbances in the origin of current oak stands. The relative ease with which certain species of oaks can be regenerated on xeric and lower quality sites appears to be a function of differences in morphology and physiology between oaks and their competition (Kozlowski 1949, Wuenscher and Kozlowski 1971, Turner and Jones 1980, Parker and Pallardy 1988, Abrams 1990). Kolb and others (1990) examined the influence of limiting light, moisture, and nutrients on growth of northern red oak and vellow-They concluded that red oak was better adapted to moderately low poplar. resource levels and, following Grime's (1979) terminology, concluded that red oak has a "stress-tolerant" strategy of growth. Conversely, yellow-poplar has a "competitor strategy" in that it was better adapted to more productive environments, where rapid resource capture is critical for survival.

> Morphological adaptations that account for the success of some oaks on xeric sites include deep roots, xeromorphic leaves, and an effective xylem transport system (Abrams 1990). The conservative seedling growth habit of most oaks that emphasizes root growth in the first few years probably imparts a survival benefit to oaks on xeric sites (Dickson and others 1990, Dickson 1991) as does the deep root system of older trees (Bourdeau 1954, Kozlowski 1971, Hinckley and others 1981). Oak leaves possess several traits, such as high stomatal density, thicker mesophyll, and smaller guard cells that may improve water use efficiency (Bidwell 1974, Abrams and Kubiske 1990). The ring-porous vascular system of oaks is composed of large diameter, early-wood vessels and narrower late-wood vessels (Zimmerman and Brown 1977). This anatomy permits rapid water movement when water is plentiful, and reduced but sustained movement during drought (Abrams 1990).

> Numerous physiological characteristics of oaks may impart a survival benefit on dry-mesic or xeric sites. There may be large differences even between different oak species (Hinckley and others 1978, 1979; Reich and Hinckley 1980; Bahari and others 1985; Chambers and Henkel 1989; Abrams and others 1990), but oaks generally appear to be better adapted to droughty sites than co-occurring species. Abrams (1990) presented an excellent review of these physiological adaptations in oaks. As compared to co-occurring species, they include: (1) higher rates of photosynthesis and less decrease in photosynthesis with increasing soil and atmospheric drought, (2) higher water use efficiency, (3) slower stomatal closure as drought progresses and higher leaf conductance, (4) lower water potential for stomatal closure, and (5) greater osmotic and elastic adjustments in the leaves.

> Oaks do occur, sometimes in almost pure stands, on mesic upland sites as well as on better bottomland sites. In light of the above discussions, it can be concluded that oaks are there because of a fortuitous set of circumstances created by nature or incidentally by acts of man (Aust and others 1987, Crow 1988), not by intentional management practices designed to obtain oak regeneration. For example, in essentially pure bottomland oak stands, we could document some form of disturbance, either fire, grazing or mowing, as instrumental in their establishment (Aust and others 1987). The lesson to be learned is that if we want to grow oaks on those sites we must create an environment that meets and favors the biological requirements of the oaks as opposed to the competing species.

ARTIFICIAL REGENERATION

Problems associated with artificially regenerating good sites are essentially the same as those encountered for natural regeneration at the time of final harvest and release, i.e., slow growth response of the small seedlings. In both cases there must be a balance between root and shoot growth, but with planted seedlings, rapid root growth is critical to maintain a balance between absorbing surface and transpiring surface (Parker 1949).

Early attempts at artificial regeneration of oaks, especially on upland sites, were not successful (Olson and Hooper 1968, McGee and Loftis 1986). Lack of success was probably related to quality of the planting stock (less than desirable size), and kind and amount of competition. More recent plantings of northern red oak have demonstrated that artificial regeneration can be successful on good sites in the Ozarks (Johnson 1984). Planting recommendations based on these results call for large planting stock (1-1 transplants) clipped about 15 cm above the root collar, establishment beneath a thinned stand (60 percent stocking), release by clearcutting 3 years later, and competition control at time of planting and at time of overstory removal. This technique should result in large, well established seedlings, with a large root system, that will respond well to release (Johnson 1984). The treatments seem to provide an environment that is compatible with the morphological adaptations and physiology of the species. Interestingly, these treatment recommendations are very similar to what recent research has shown may work well for natural regeneration (Janzen and Hodges 1987, Kruger and Reich 1989, Lockhart 1992).

SUMMARY AND SILVICULTURAL IMPLICATIONS

Acorn production, acorn losses, and poor initial seedling establishment can, in some cases, account for oak regeneration failures, but overall the major cause of failure on good sites seems to be a slow juvenile growth rate of oak seedlings and the inability to respond to release. The problem is one of competition—the inability of oaks to compete efficiently with more tolerant species, especially those in the lower canopies at low light levels, and with well established and/or faster-growing species under open conditions. The differences between species are the result of differences in morphology and physiology and the ability to acclimate to prevailing environmental conditions.


Figure 3 is an attempt to depict what is known about success of oak regeneration as influenced by site and competition with co-occurring species. Oaks are generally not very "flexible", i.e., they do not acclimate morphologically and physiologically well to changing environments, especially light. On good sites under a dense canopy oaks may undergo slight morphological and physiological acclimations, but these are insufficient to enable them to compete effectively with Compared to such species, oaks have a higher light more tolerant species. compensation point, are less efficient at use of "light flecks," have similar or lower rates of net photosynthesis, have higher rates of respiration, have slower stomatal opening and/or greater stomatal resistance, and have lower quantum yield. At the other extreme, these same oak seedlings may not compete well after release even The poor competitive ability reflects physiological and in full sunlight. morphological characteristics of the oaks. Photosynthesis may be saturated near one-third of full sunlight and rates of net photosynthesis are far less than for intolerant co-occurring species. Furthermore, carbohydrate allocation patterns in oak seedlings emphasize root growth rather than shoot growth. Intolerant


Page 121 of 404

competitors allocate more reserves to shoot growth, giving them a height advantage.

On drier mesic or xeric sites, oaks are in a better competitive position than on more mesic sites. This competitive position again reflects differences in morphology and physiology between the oaks and competing species. Morphological and anatomical characteristics of leaves and xylem, as well as carbon allocation patterns that favor root growth, result in better water-use efficiency and less, or delayed, moisture stress in the oaks. Less stress in turn means less reduction in physiological processes, such as photosynthesis, and therefore better growth.

This brief review of oak ecology and physiology helps explain why it is often difficult to obtain satisfactory regeneration of oaks. The major problems are competition with more tolerant and/or faster growing species and the "inflexible" nature of the growth habit of oaks. Oak regeneration efforts can be successful as long as ecological and physiological requirements are understood and an environment is created which favors those requirements.





LITERATURE CITED

- Abrams, Marc D. 1990. Adaptations and responses to drought in *Quercus* species of North America. Tree Physiology. 7:227-238.
- Abrams, Marc D.; Kubiske, M.E. 1990. Leaf structural characteristics of 31 hardwood and conifer tree species in central Wisconsin: influence of light regime and shade tolerance rank. Forest Ecology and Management. 31:245-253.

- Abrams, Marc D.; Schultz, Jack C.; Kleiner, Karl W. 1990. Ecophysiological responses in mesic versus xeric hardwood species to an early-season drought in central Pennsylvania. Forest Science. 36(4):970-981.
- Arend, John L.; Gysel, Leslie W. 1952. Less oak reproduction on better sites. Tech. Note 378. Lake States Forest Experiment Station. 1 pp.
- Aust, W. Michael; Hodges, J.D.; Johnson, R.L. 1987. The origin, growth and development of natural pure, even-aged stands of bottomland oak. Proceedings of the Third Biennial Southern Silvicultural Research Conference; 1984 November 7-8; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 163-170.
- Bahari, Z.A.; Pallardy, S.G.; Parker, W.C. 1985. Photosynthesis, water relations, and drought adaptation in six woody species of oak-hickory forests in central Missouri. Forest Science. 31:557-569.
- Bazzaz, F.A.; Carlson, R.W. 1982. Photosynthetic acclimation to variability in the light environment of early and late successional plants. Oeocologia (Berl). 54:313-316.
- Beck, Donald E. 1970. Effect of competition on survival and height growth of red oak seedlings. Res. Pap. SE-56. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 7 pp.
- Beck, Donald E.; Hooper, Ralph M. 1986. Development of a southern Appalachian hardwood stand after clearcutting. Southern Journal of Applied Forestry. 10:168-172.

Bidwell, R.G.S. 1974. Plant Physiology. New York: Macmillan Publishing Co. 643 pp.

- Blake, T.J.; Tschaplinski, T.J. 1986. Role of water relations and photosynthesis in the release of buds from apical dominance and the early reinvigoration of decapitated poplars. Physiologia Plantarum. 68:287-293.
- Bourdeau, Philippe. 1954. Oak seedling ecology determining segregation of species in Piedmont oak-hickory forests. Ecological Monographs. 24(3):297-320.
- Carvell, K.L.; Tryon, E.H. 1961. The effect of environmental factors on the abundance of oak regeneration beneath mature oak stands. Forest Science. 7:98-105.
- Chambers, Jim L.; Henkel, Mary W. 1989. Survival and growth of natural and artificial regeneration in bottomland hardwood stands after partial overstory removal. Proceedings of the Fifth Biennial Southern Silvicultural Research Conference; 1988 November 1-3; Memphis, TN. Gen. Tech. Rep. SO-74. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 277-283.
- Collier, Donald E.; Cummins, W. Raymond; Villar, Rafael. 1992. Diurnal patterns of respiration in the leaves of four forest tree species. Physiologia Plantarum. 84:361-366.
- Crow, T.R. 1988. Reproductive mode and mechanisms for self-replacement of northern red oak (*Quercus rubra*) -- a review. Forest Science. 34(1):19-40.
- Darley-Hill, S.; Johnson, W.C. 1981. Acorn dispersal by the blue-jay (Cyanocitta cristata). Oecologia (Berl.). 50:232-232.
- Davies, W.J.; Kozlowski, T.T. 1974. Stomatal responses of five woody angiosperms to light intensity and humidity. Canadian Journal of Botany. 52:1525-1534.
- Deen, Robert T.; Hodges, John D. 1991. Oak regeneration in abandoned fields: presumed role of the blue jay. Proceedings of the sixth biennial southern silvicultural research conference; 1990 October 30-November 1; Memphis, TN. Gen. Tech. Rep. SE-70. Asheville, NC: U.S. Department of Agriculture, Forest Service.

- Dickson, Richard E. 1991. Episodic growth and carbon physiology in northern red oak. In: Laursen, Steven B.; De Boe, Joyce F. The oak resource in the upper Midwest: implications for management; 1991 June 3-6; Winona, MN. St. Paul, MN: Minnesota Extension Service, University of Minnesota: 117-124.
- Dickson, Richard E.; Isebrands, J.G.; Tomlinson, Patricia T. 1990. Distribution and metabolism of current photosynthate by single-flush northern red oak seedlings. Tree Physiology. 7:65-77.
- Farmer, R.E., Jr. 1980. Comparative analysis of 1st-year growth in six deciduous tree species. Canadian Journal of Forest Research. 10:35-41.
- Gottschalk, Kurt W. 1987. Effects of shading on growth and development of northern red oak, black oak, black cherry, and red maple seedlings. I. Height, diameter, and root/shoot ratio. In: Dawson, J.O.; Majerus, K.A. Proceedings 5th Central Hardwood Forest Conference; Urbana-Champaign, IL: 189-195.
- Grime, J.P. 1979. Plant Strategies and Vegetation Processes. New York: Wiley. 222 pp.
- Grime, J.P.; Jeffrey, D.W. 1965. Seedling establishment in vertical gradients of sunlight. Journal of Ecology. 53:621-642.
- Hanson, P.J. 1986. Studies of *Quercus rubra* L. seedling dry matter accumulation, morphological development, and carbon dioxide exchange under controlled conditions. University of Minnesota. 169 pp. Dissertation.
- Hara, Masatoshi. 1987. Analysis of seedling banks of a climax beech forest: ecological importance of seedling sprouts. Vegetatio. 71:67-74.
- Hill, John P.; Dickmann, Donald I. 1988. A comparison of three methods for naturally reproducing oak in southern Michigan. Northern Journal of Applied Forestry. 5:113-117.
- Hinckley, T.M.; Aslin, R.G.; Aubuchon, R.R.; Metcalf, C.L.; Roberts, J.E. 1978. Leaf conductance and photosynthesis in four species of the oak-hickory forest type. Forest Science. 24:73-84.
- Hinckley, T.M.; Dougherty, P.M.; Lassoie, J.P.; Roberts, J.E.; Teskey, R.O. 1979. A severe drought: impact on tree growth, phenology, net photosynthetic rate and water relations. American Midland Naturalist. 102:307-316.
- Hinckley, T.M.; Teskey, R.O.; Duhme, F.; Richter, H. 1981. Temperate hardwood forests. In: Kozlowski, T.T.; Water Deficits and Plant Growth. Vol. VI. Academic Press, New York: 153-208.
- Hix, David M.; Lorimer, Craig G. 1990. Growth-competition relationships in young hardwood stands on two contrasting sites in southwestern Wisconsin. Forest Science. 36(4):1032-1049.
- Hix, David M.; Lorimer, Craig G. 1991. Early stand development on former oak sites in southwestern Wisconsin. Forest Ecology and Management. 42:169-193.
- Hodges, John D. 1987. Cutting mixed bottomland hardwoods for good growth and regeneration.
 Proceedings Fifteenth Annual Hardwood Symposium of the Hardwood Research Council; 1987
 May 10-12; Memphis, TN: 53-60.
- Hodges, John D.; Janzen, Greg. 1987. Studies on the biology of cherrybark oak: recommendations for regeneration. Proceedings of Fourth Biennial Southern Silvicultural Research Conference; 1986 November 4-6; Atlanta, GA. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 133-139.
- Janzen, G.C.; Hodges, J.D. 1985. Influence of midstory and understory vegetation removal on the establishment and development of oak regeneration. Proceedings of the Third Biennial Southern Silvicultural Research Conference; 1984 November 7-8; Atlanta, GA. Tech. Rep. SO-54. New

65a Oak Regeneration Symposium Proceedings . . . September 1992—ERRATA (February 1994)

Orleans, LA: U.S. Department of Agricutture, Forest Service, Southern Forest Experiment Station: 273-278.

- Janzen, G.C.; Hodges, J.D. 1987. Development of advanced oak regeneration as influenced by removal of midstory and under-story vegetation. Proceedings of the Fourth Southern Silvicultural Research Conference; 1986 November 4-6; Atlanta, GA. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 455-461.
- Johnson, I.R.; Thornley, J.H.M. 1987. A model of shoot: root partitioning with optimal growth. Annals of Botany. 60:133-142.
- Johnson, Paul S. 1979a. Growth potential and field performance of planted oaks. In: Holt, H.A.; Fischer, B.C. Regenerating oaks in upland hardwood forests. John S. Wright Forestry Conference Proceedings. West Lafayette, IN: Purdue University: 113-119.
- Johnson, Paul S. 1979b. Shoot elongation of black oak and white oak sprouts. Canadian Journal of Forest Research. 9(4):489-494.
- Johnson, Paul S. 1984. Responses of planted northern red oak to three overstory treatments. Canadian Journal of Forest Research. 14:536-542.
- Johnson, Paul S.; Jacobs, Rodney D. 1981. Northern red oak regeneration after preherbicided clearcutting and shelterwood removal cutting. Res. Paper NC-202. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 5 pp.
- Johnson, R.L. 1975. Natural regeneration and development of Nuttall oak and associated species. Res. Paper SO-104. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 12 pp.
- Jones, Robert H.; Sharitz, Rebecca R. 1990. Dynamics of advance regeneration in four South Carolina bottomland hardwood forests. Proceedings of the sixth biennial Southern Silvicultural Research Conference; 1990 October 30-November 1; Memphis, TN. Gen. Tech. Rep. SE-70. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 567-578.
- Jones, Robert H.; Sharitz, Rebecca R.; McLeod, Kenneth W. 1989. Effects of flooding and root competition on growth of shaded hardwood seedlings. American Midland Naturalist. 121:165-175.
- Kohyama, Takashi. 1980. Growth pattern of *Abies mariesii* saplings under conditions of opengrowth and suppression. Botanical Magazine of Tokyo. 93:13-24.
- Kolb, T.E.; Steiner, K.C. 1990a. Growth and biomass partitioning of northern red oak and yellowpoplar seedlings: effects of shading and grass root competition. Forest Science. 36(1):34-44.
- Kolb, T.E.; Steiner, K.C. 1990b. Growth and biomass partitioning response of northern red oak genotypes to shading and grass root competition. Forest Science. 36(2):293-303.
- Kolb, T.E.; Steiner, K.C.; McCormick, L.H.; Bowersox, T.W. 1990. Growth response of northern red-oak and yellow-poplar seedlings to light, soil moisture and nutrients in relation to ecological strategy. Forest Ecology and Management. 38:65-78.
- Kozlowski, Theodore T. 1949. Light and water in relation to growth and competition of Piedmont forest tree species. Ecological Monographs. 19(3):207-231.
- Kozlowski, T.T. 1971. Growth and development of trees. Vol. II. Cambial growth, root growth and reproductive growth. New York: Academic Press. 514 pp.

- Kramer, P.J; Decker, J.P. 1944. Relation between light intensity and rate of photosynthesis of loblolly pine and certain hardwoods. Plant Physiology. 19:350-358.
- Kramer, P.J; Kozlowski, T.T. 1979. Physiology of Woody Plants. New York: Academic Press. 811 pp.
- Kruger, E.L.; Reich, P.B. 1989. Comparative growth and physiology of stem-pruned and unpruned northern red oak. In: Rink, G.; Budelsky, C.A. Proceedings of Seventh Central Hardwood Conference; 1989 March 5-8; Carbondale, IL. Gen. Tech. Rep. NC-132. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 302 pp.
- Larson, M.M. 1980. Effects of atmospheric humidity and zonal soil water stress on initial growth of planted northern red oak seedlings. Canadian Journal of Forest Research. 10:549-554.
- Larson, M.M.; Whitmore, F.W. 1970. Moisture stress affects root regeneration and early growth of red oak seedlings. Forest Science. 16(4)495-498.
- Loach, K. 1967. Shade tolerance in tree seedlings. I. Leaf photosynthesis and respiration in plants raised under artificial shade. New Phytologist. 66:607-621.
- Loach, K. 1970. Shade tolerance in tree seedlings. II. Growth analysis of plants raised under artificial shade. New Phytologist. 69:273-286.
- Lockhart, Brian Roy. 1992. Morphology, gas-exchange, and ¹⁴C- photosynthate allocation patterns in advanced cherrybark oak reproduction. Mississippi State University. 192 pp. Dissertation.
- Loftis, David L. 1979. Northern red oak performs poorly in North Carolina planting. Res. Note SE-277. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 3 pp.
- Loftis, David L. 1983. Regenerating red oak on productive sites in the southern Appalachians: a research proposal. Proceedings of the Second Biennial Southern Silvicultural Research Conference; 1982 November 4-5; Atlanta, GA. Tech. Rep. SE-24. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 144-150.
- Loftis, David L. 1988. Regenerating oaks on high-quality sites; an update. In: Workshop proceedings: Guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV: 199-209.
- Loftis, David L. 1990. A shelterwood method for regenerating red oak in the southern Appalachians. Forest Science. 36(4):917-929.
- Lorimer, Craig G. 1981. Survival and growth of understory trees in oak forests of the Hudson Highlands, New York. Canadian Journal of Forest Research. 11:689-695.
- Martin, A. Jeff; Hix, David M. 1988. Regeneration development in an upland hardwood stand following a shelterwood harvest. Northern Journal of Applied Forestry. 5:46-49.
- Matsuda, Kozue; McBride, Joe R. 1986. Difference in seedling growth morphology as a factor in the distribution of three oaks in central California. Madroño. 33(3):207-216.
- Matsuda, Kozue; McBride, Joe R.; Kimura, Makoto. 1989. Seedling growth form of oaks. Annals of Botany. 64:439-446.
- McGee, C.E. 1975. Change in forest canopy affects phenology and development of northern red and scarlet oak seedlings. Forest Science. 22(2):175-179.
- McGee, Charles E. 1981. Response of overtopped white oak to release. Res. Note SO-273. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 pp.

65c Oak Regeneration Symposium Proceedings . . . September 1992—ERRATA (February 1994)

- McGee, Charles E. 1986. Budbreak for twenty-three upland hardwoods compared under forest canopies and in recent clearcuts. Forest Science. 32(4):924-935.
- McGee, Charles E. 1988. Spring weather, canopy removal, and early budbreak threaten oak seedlings. Journal of the Elisha Mitchell Society. 104(3):108-115
- McGee, Charles E.; Loftis, David L. 1986. Planted oaks perform poorly in North Carolina and Tennessee. Northern Journal of Applied Forestry. 3:114-116.
- McQuilkin, Robert A. 1975. Growth of four types of white oak reproduction after clearcutting in the Missouri Ozarks. Res. Pap. NC-116. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 5 pp.
- Merritt, C.; Pope, P.E. 1991. Environmental effects, including wildfire and prescribed burning, on the establishment of oaks in Indiana. Station Bull. 612, Dept. of Forestry and Natural Resources, West Lafayette, IN: Purdue University. 52 pp.
- Musselman, R.C.; Gatherum, G.E. 1969. Effects of light and moisture on red oak seedlings. Iowa State Journal of Science. 43(3):273-284.
- Olson, D.F.; Hooper, R.M. 1968. Early survival and growth of planted northern red oak in the Southern Appalachians. Res. Note SE-89. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 3 pp.
- Parker, J. 1949. Effects of variations in the root-leaf ratio on transpiration rate. Plant Physiology. 24:739-743.
- Parker, William C.; Pallardy, Stephen G. 1988. Leaf and root osmotic adjustment in droughtstressed Quercus alba, Q. macrocarpa, and Q. stellata seedlings. Canadian Journal of Forest Research. 18:1-5.
- Phares, Robert E. 1971. Growth of red oak (Quercus rubra L.) seedlings in relation to light and nutrients. Ecology. 52(4):669-672.
- Pubanz, Dan M.; Lorimer, Craig G. 1992. Oak regeneration experiments in southwestern Wisconsin: two-year results. Res. Rep. R3552. Madison, WI: School of Natural Resources, College of Agriculture and Life Sciences, University of Wisconsin-Madison. 16 pp.
- Reich, P.B.; Hinckley, T.M. 1980. Water relations, soil fertility and plant nutrient composition of a pygmy oak ecosystem. Ecology. 61:400-416.
- Richardson, S.D. 1956. On the role of the acorn in root growth of American oak seedlings. Medelelingen van de Landhouwhogeschool te Wageningen-Nederland. 56(12):1-18.
- Sander, Ivan L. 1972. Size of oak advance reproduction: key to growth following harvest cutting. Res. Pap. NC-79. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 6 pp.
- Sander, Ivan L.; Clark, F.B. 1971. Reproduction of upland hardwood forests in the Central States. Agricultural Handbook 405. Washington, D.C.: U.S. Department of Agriculture, Forest Service. 25 pp.
- Schulze, E.D.; Schilling, K.; Nagarajah, S. 1983. Carbohydrate partitioning in relation to whole plant production and water use of Vigna unguiculata (L.) Walp. Oecologia (Berl.). 58:169-171.
- Smith, H. Clay. 1993. Regenerating oaks in the central Appalachians. In: Proceedings, A Symposium on oak regeneration; 1992 September 8-10; Knoxville, TN.
- Teskey, Robert O.; Shrestha, Rajesh B. 1985. A relationship between carbon dioxide, photosynthetic efficiency and shade tolerance. Physiologia Plantarum. 63:126-132

- Tubbs, C.H. 1977. Natural regeneration of northern hardwoods in the northern Great Lakes region. Res. Pap. NC-150. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 20 pp.
- Turner, N.C.; Jones, M.M. 1980. Turgor maintenance by osmotic adjustment: a review and evaluation. In: Turner, N.C.; Kramer, P.J. Adaptations of Plants to Water and High Temperature Stress. New York. John Wiley and Sons. 87-103 pp.
- Wassink, E.C.; Richardson, S.D. 1951. Observations on the connection between root growth and shoot illumination in first-year seedlings of *Acer pseudoplatanus* L. and *Quercus borealis maxima* (Marsh) Ashe. Koninklijke Nederlandse Akademie van Wetenschappen Proceedings Series C. 54:503-510.
- Went, F.W. 1957. The experimental control of plant growth. Chronica Botany. 17:257.
- Woods, David B.; Turner, Neil C. 1971. Stomatal response to changing light by four tree species of varying shade tolerance. New Phytologist. 70:77-84.
- Wuenscher, J.E.; Kozlowski, T.T. 1970. Carbon dioxide transfer as a factor in shade tolerance of tree seedlings. Canadian Journal of Botany. 48:453-456.
- Wuenscher, J.E.; Kozlowski, T.T. 1971. Relationship of gas-exchange resistance to tree-seedling ecology. Ecology. 52(6):1016-1023.
- Zimmerman, M.H.; Brown, C.L. 1977. Tree structure and function. New York: Springer-Verlag. 336 pp.

The Role of Fire in Oak Regeneration

David H. Van Lear, Department of Forest Resources, Clemson University, Clemson, SC 29634 Janet M. Watt, South Carolina Forestry Commission, Newberry, SC 29108

ABSTRACT

Fire has played a dominant role in sustaining oak forests. Oak species have biological adaptations, such as thick bark, a tenacious ability to resprout repeatedly following top-kill, and resistance to rot, which enable them, better than their competitors, to withstand a regime of frequent fire. Fire functions to encourage establishment of oak regeneration by: (1) creating favorable conditions for acorn caching by squirrels and bluejays, (2) reducing populations of insects which prey on acorns and young oak seedlings, (3) xerifying mesic sites through consumption of surface organic materials and exposure of the soil to greater solar radiation, and (4) reducing understory and midstory competition from fire-intolerant species. The ability of oaks to continually resprout when numbers of other sprouting hardwoods have been reduced by fire may allow oak to accumulate in the advance regeneration pool. Improved root/shoot ratios resulting from frequent top-kill should enhance response of oak seedling/sprouts to release and enable them to dominate when stand-level disturbances create conditions favorable for rapid growth. Based on biological adaptations of oak to fire, ecological functions of fire, and fire history, tentative guidelines are presented for using fire to promote oak regeneration on better sites. Effects of wildfires and intense fires in logging debris on establishment and development of oak-dominated stands are discussed.

INTRODUCTION Oaks are often replaced by other species when mature stands are harvested, especially on better quality sites (Sander and others 1983, Loftis 1990, Abrams 1992). Even though researchers generally agree that fire played a role in the establishment of many oak-dominated stands at the turn of the century (Sander and others 1983, Crow 1988, Maslen 1989), there is relatively little research concerning the use of fire in oak ecosystems. Most of the forestry research about fire-oak relationships has dealt with the use of fire to control oaks in pine stands. However, many foresters and ecologists are now recognizing the importance of simulating the natural disturbance regime, which often included frequent fire, to maintain the species composition of certain ecosystems. The purpose of this paper is to (1) describe the role of fire in ecology of oak regeneration, and (2) present tentative guidelines for the silvicultural use of fire to regenerate oak.

THE ROLE OF FIRE IN THE ECOLOGY OF OAK REGENERATION Adaptations of Oak to Fire Fire has traditionally been used in forest management to control plant succession. It is well documented by literally dozens of studies that both dormant- and growing-season fires in pine stands will top-kill small hardwood stems, including oaks. Frequent burns will arrest the development of the hardwood understory, although most species continue to resprout for years. However, it is interesting that oaks have lower mortality rates than competing species in regimes of frequent fire. Waldrop and Lloyd (1991) reported that oak mortality rates after 26 years of biennial summer burning in mature pine stands in the Coastal Plain were still below 50 percent, whereas mortality rates of other woody species ranged from about 60 to 80 percent.

This tenacious ability of small oak rootstocks to resprout repeatedly following frequent top-kill is an important adaptation of oak to frequent fire regimes. This characteristic should enable oak to dominate the advance regeneration pool in areas where fire occurs at frequent intervals. In addition, continued top-killing should result in a more favorable root/shoot ratio and faster growth after release. Other biological adaptations, such as thick bark, resistance to rotting after scarring, and the suitability of fire-created seedbeds for acorn germination (Lorimer 1985) enhance the ability of oaks to survive on sites exposed to frequent fire. Martin (1989) suggests that bark thickness may be the single attribute that best characterizes a species' adaptation to fire. While bark thickness is undoubtedly of great importance to the survival of mature trees in regimes of frequent fire, it would seem that the ability of oak advance regeneration to outlast its competition would be the critical factor insuring that oak is a major component of the next stand (Van Lear 1990).

Functions of Fire in Oak Regeneration Fire has numerous functions which benefit oak regeneration (table 1). Fire removes excessive litter buildup from the forest floor, thereby preparing a favorable seedbed. Areas of thin litter are preferred by squirrels and bluejays for acorn burial (Galford and others 1988). An important ecological finding is that jays collect and disperse only sound nuts (Darley-Hill and Johnson 1981, Deen and Hodges 1990), which implies that if these acorns escape predation they will result in well-established first-year seedlings. Seedlings from freshly germinated acorns are unable to emerge through a heavy litter cover. Germination and first-year survival are best when acorns are buried about 3 cm deep in the mineral soil (Sander and others 1983).

> Although removal of thick litter may expedite the germination process by encouraging the caching of acorns by squirrels and jays, it is important that not all the humus layer be consumed. The humus layer keeps the surface of the soil porous, so that uncached acorns can more easily penetrate the soil, retains moisture, and provides support for the new seedling (Carvell and Tryon 1961). The intensity and severity of a prescribed burn will determine the amount of organic matter lost on a site (Wells 1979).

> Fire helps to control insect predators of acorns and new seedlings. Martin and Mitchell (1981) illustrated how insect populations can be reduced or eliminated directly or indirectly by fire (table 2). Insect pests act as primary invaders, secondary invaders, parasites, or scavengers on or in acorns (Gibson 1972). Many of these insects spend all or part of their lives on the forest floor. Infestations, which can vary from year to year and even from tree to tree in some areas, are a major contributor to the oak regeneration problem (Marquis and others 1976).

Annually about 50 percent of the acorn crop in Ohio is destroyed by the larvae of *Curculio* weevils, acorn moths, and gall wasps. Other insects attack germinating Page 130 of 404

acorns and oak seedlings. However, recent studies indicate that prescribed burning may reduce populations of oak insect pests when conducted under proper conditions and at the appropriate time in the insects' life cycle (Galford and others 1988). A reduction in insect predation would allow more acorns to be scattered and buried by jays and squirrels, thus enhancing the probability of successful germination, and also encourage subsequent seedling establishment. Burning may also reduce rodent habitat, eliminating another source of acorn predation (Hannah 1987).

Function	Reference		
Prepare seedbed and encourage caching	Galford and others 1988, Sander and others 1983		
Discourage acorn and seedling predators	Galford and others 1988, Martin and Mitchell 1981, Hannah 1987		
Open understory and reduce fire-intolerant competitors	Crow 1988, Maslen 1989, Harmon 1984, Martin 1989, Van Lear and Waldrop 1989, Loftis 1990		
Xerify sites	Crow 1988, Van Lear 1990		
Allow oak to dominate advance regeneration pool	Little 1974, Van Lear and Waldrop 1983		
Increase flammable fuels	Komarek 1965, Martin and others 1975		

Table 1-Functional roles of fire in the ecology of oak regeneration

Table 2-Effects of fire on insects¹

	Direct Effects
HEAT > INSECTS SMOKE	In litter, duff On understory vegetation Exposed on trees
	Indirect Effects
ORGANIC MATT > VEGETATION NUTRIENTS	ER > INSECTS
> VEGETATION	INSECT Grass & forbs > HABITAT Shrubs Trees
	> INSECTS SMOKE ORGANIC MATT > VEGETATION NUTRIENTS

¹ Martin and Mitchell (1981).

A regime of frequent burning over long periods of time creates an open stand in pine or hardwood stands. In hardwood stands, long-term burning tends to eliminate small understory stems outright and gradually reduces the midstory and overstory canopy through mortality resulting from fire wounds. Increased light reaching the forest floor in these open stands will maintain the vigor of oak advance regeneration. Loftis (1990) demonstrated that elimination of the subcanopy with herbicides encouraged development of advance regeneration of red oak in mature mixed hardwood stands in the Southern Appalachians. Long-term burning should create a stand structure similar to those created by injecting understory hardwoods with herbicides.

Severe or frequent fires xerify the surface of forest sites by consuming much of the forest floor and perhaps even organic matter in the mineral soil, as well as by exposing the site to greater solar radiation through canopy reduction (Van Lear 1990). Adequate advanced oak regeneration in the Southeast is generally found more often on xeric sites than on mesic sites (Sander 1988). Crow (1988) cited the lower frequency of fire in recent years as a major factor in the failure of oak to regenerate on better sites. Conversion of mesic sites to more xeric conditions by intense fires or by a long regime of low intensity fires could explain in large part the ability of oaks to dominate sites where more mesic species normally occur. Mesic sites may only have burned during cyclic periods of dry weather which have apparently occurred in the Southeast for millennia.

The absence of fire since the turn of the century has allowed species that are intolerant of fire to become established and grow to a size where they, because of thicker bark associated with age, can now resist fire. At greater than 5 cm (2 in.) d.b. h., yellow-poplar becomes almost as fire resistant as oaks (Maslen 1989). Mockernut and pignut hickories, scarlet oak, red maple, and blackgum are examples of such species that are often found on sites where fire has been long absent (Harmon 1984, Martin 1989).

Suppression of fire has allowed shrubby understory species to occupy drier sites where fire was once frequent and oak more dominant. In particular, rhododendron has dramatically increased its areal extent (Van Lear and Waldrop 1989, Martin 1989). Impenetrable thickets of ericaceous species, such as rhododendron, mountain laurel, and huckleberry, now often dominate midstories and understories of hardwood stands in the Southern Appalachians and prevent desirable hardwood regeneration from becoming established (Beck 1989). Fire would top-kill these species and, although they do sprout, new growth is slow and they would likely be relatively unsuccessful competitors of regenerating oaks.

Yellow-poplar produces an abundance of seed almost annually, and although the seed has low viability, many remain viable in the litter and duff layer for several years (Carvell and others 1955, Maslen 1989). Yellow-poplar seed germinate readily following burning (Shearin and others 1972). However, in a regime of frequent fire, small yellow-poplar seedlings would be killed and the reservoir of stored seed in the duff would be gradually depleted. Thus, frequent fires would control to a large degree this major competitor of oaks on high-quality sites.

Frequent fire functions to allow accumulation of oak in the advance regeneration pool. Nearly all hardwood species sprout in a regime of annual winter fire (Thor and Nichols 1974, Langdon 1981, Waldrop and others 1987). Hardwood sprouting is more vigorous following periodic winter burns because of greater carbohydrate reserves (Hodgkins 1958). Thor and Nichols (1974) noted that even with periodic and annual winter burning, oak stems tend to increase at the expense of competing hardwoods. After two periodic winter burns and eight annual winter burns, oak stems comprised 61 and 67 percent of the total stems, compared to 51 percent oak stems on the unburned plots.

Annual summer fires eventually eliminate all hardwood sprouts (Langdon 1981, Waldrop and others 1987). Biennial summer fires also gradually eliminate hardwood sprouts, but, as mentioned earlier, oak succumbs more slowly than many other species (figure 1). Oaks, in the absence of prolific root sprouters, such as sweetgum, would gradually dominate the advance regeneration pool in mature mixed hardwood stands because of the tenacity of their sprouting (Carvell and Tryon 1961, Waldrop and others 1987). Increases in the number of oak sprouts and, more importantly, the number of top-killed oak stems (up to 15-cm ground diameter) with basal sprouts following summer broadcast burns suggest that periodic summer burning would be expected to favor oak even more (Augspurger and others 1987).




Figure 1—Cumulative mortality of hardwood roots over 26 years of biennial prescribed burning (Langdon 1981).

When repeated burning occurs in stands with mixed advanced regeneration, oaks have an advantage over less fire-resistant vegetation, which is killed by fewer fires of lower intensity (Waldrop and others 1987). This loss usually exceeds species gain through invasion, since the frequency of the fires is as important to reduction of fire-susceptible species as the intensity of the fire (De Selm and others 1990). Sander (1988) stated that effective hardwood competition control may require as many as three or more burns at 2- to 3-year intervals.

Studies of effects of single fires on composition of mixed stands have produced varied results. McGee (1979) found that single spring and fall burns in small sapling-sized mixed hardwood stands in northern Alabama had little effect on species composition other than to increase relative dominance of red maple and the number of multiple stem oak clumps. However, a single intense wildfire in a young mixed hardwood stand in West Virginia shifted species composition to a predominately oak stand (Carvell and Maxey 1969).

Frequent fire in oak stands may also increase the production of legumes and grasses, which benefit numerous wildlife species, but which also create a more flammable understory. At the turn of the century, summer fires were quite common in the Southeastern United States as farmers burned the land to facilitate grazing. They had learned from early settlers, who in turn had learned from their Indian predecessors, that growing-season fires best maintained an open forest with a rich herbaceous layer (Komarek 1965). Thor and Nichols (1974) noted an increase in herbaceous vegetation following frequent burning in mixed hardwood stands in Tennessee. Similar findings have been reported in pine forests of the Southeast by numerous researchers. Therefore, a burning regime of frequent fire functions to create and maintain a ground cover that encourages the return of fire, which for the reasons stated above would favor the establishment of oak advance regeneration.

SILVICULTURAL USE OF FIRE IN OAK REGENERATION

While some new oak stands result from stump sprouts, there is little dispute among silviculturists that oak advance regeneration is often critical to the re-establishment of many oak-dominated stands (Clark and Watt 1971, Sander and others 1983, Loftis 1988, Lorimer 1989). However, while many acknowledge that fire may have played a role in creating the present mature oak stands, no guidelines have been developed for using fire to regenerate oak stands.

Based on the history of fire in the southeastern United States, and on biological adaptations of oak and ecological functions of fire discussed earlier, the following *tentative* guidelines are suggested for using fire in oak management. Our hypothesis is that silviculture which mimics the disturbance regime that created present-day stands dominated by mature oak will create future stands dominated by oak. Further research will be necessary to test and fine-tune these suggestions before they can be recommended as silvicultural practices.

To Promote Advance Regeneration It has been suggested (Little 1974, Sander 1988, Van Lear and Waldrop 1989) that an extended period of repeated burns prior to harvest may improve the status of oak in the advance regeneration pool, especially on better sites. Although figure 2 depicts the sequence of suggested actions and likely responses to fire, there is no research that currently documents a series of burning treatments that will

successfully accomplish this goal. Therefore, a burning regime might include a mix of winter and summer fires adjusted to enhance the relative position of oak in the advance regeneration pool.

The famous Santee Fire Plot study showed that annual summer burns for 5 years in a pine stand in the Coastal Plain killed about 40 percent of oak root stocks compared to 55 to 90 percent of other woody competitors (Waldrop and others




Figure 2—Tentative scenario of using prescribed fire to encourage advance reproduction of oak.

1987, Waldrop and Lloyd 1991). Biennial summer burning killed hardwood root stocks more slowly, but the rate of mortality for other woody species was still significantly greater than that of oak species. Annual winter burning, while not as effective as summer burning in altering species composition, still tends to xerify the site by consuming litter and reducing shading of top-killed understory species.

Initial height growth of oak advance regeneration is slow, since most of the early growth goes into the root system (Kelty 1988). Burning can increase the average annual shoot growth of oak seedlings, providing a potential advantage over competing stems (Johnson 1974). Oak advanced regeneration occurs as true seedlings or sprouts; the latter have root systems older than the stems and are often referred to as seedling-sprouts (Sander and others 1976). A large root system is necessary for initiation of shoot growth when environmental conditions become favorable (Crow 1988). Thus, a regime of frequent understory burns, perhaps including both growing-season and winter burns during a period of 5 to 20 years prior to harvest, should promote a favorable root/shoot ratio during oak seedling establishment. The timing of the burns would be dependent on the observed vigor of the oak advance regeneration and its competitors.

Once an adequate number of oak seedling-sprouts are present and numbers of competing species have been sufficiently reduced, fire should be withheld to allow the oak advance regeneration to attain sufficient size to outgrow other species which germinate or sprout after the mature stand is cut. A relatively open stand with few midstory and understory trees would provide adequate light for the oak advance regeneration to develop into stems of sufficient size to outgrow other species after the overstory is removed. Sander and others (1983) recommend that 1,075 advance regeneration oak stems/ha over 1.5 m tall be present before the overstory is removed.

Herbicides may be required to remove midstory trees that have grown too large to be killed by low-intensity fires. Loftis (1988, 1990) has convincingly shown that growth of advance regeneration of northern red oak can be enhanced by herbicidal removal of midstory and understory competitors. Herbicides provide initial selectivity of midstory stems to be eliminated prior to burning. A combination of herbicide treatment and frequent fire may be required to secure oak regeneration and allow it to maintain its vigor in mixed hardwood forests which have not been burned for decades.

Although methods have been developed to predict fire-induced mortality of large trees based on stem size and extent of fire damage (Loomis 1973), research is needed to determine if and how prescribed fire can be used without excessive damage to stems of large valuable crop trees in mature hardwood stands. It should be understood that the prescribed fire regime being suggested here is for use of fire only during the regeneration period. In this case, if rot should develop in damaged trees, it will have a relatively short time to grow and damage butt-log quality.

Foresters have long recognized that wildfire is a major cause of butt rot in hardwoods, but relatively little information is available concerning the relationship between prescribed fires and stem damage. Wendel and Smith (1986) found that a strip-head fire in the spring in an oak-hickory stand in West Virginia caused a

decline in overstory vigor and resulted in death of many trees during the 5 years after burning. However, a low-intensity winter fire in a mixed hardwood stand in the Southern Appalachians resulted in little or no cambium damage to large crop trees (Sanders and others 1987). More research is needed to determine the extent of bole damage from prescribed fire in different seasons and under varying frequencies in mature hardwood stands.

It must be emphasized that a series of burns over an indefinite preharvest period will likely be required to favor oak regeneration. The first burn may be detrimental to oak advance regeneration in that small rootstocks may be killed. However, our interpretation of the literature suggests that, over the long-run, oak will be less adversely affected than competitors and will therefore receive a competitive advantage that will enable them to favorably respond to subsequent release.

Vigorous, abundant sprouting of oaks occurs following broadcast burning of clearcut areas. Burning of clearcut hardwood or mixed pine-hardwood stands promotes better quality oak sprouts by forcing them to develop from the groundline. Over 97 percent of all oak sprouts developing after broadcast burning from for unburned areas (Augspurger and others 1987). Suppressed buds higher on the buds at or below groundline is encouraged by fire, reducing the likelihood of rot being transferred from stumps to new sprouts (Roth and others 1939). Poorly formed tops of small (<15 cm) oak stems killed by fire, reducing the likelihood of rot desirable sprouts, which are more likely to develop into sound timber trees than other types of oak regeneration (Roth and Hepting 1943, Teuke and Van Lear other types of oak regeneration (Roth and Hepting 1943, Teuke and Van Lear other types of oak regeneration (Roth and Hepting 1943, Teuke and Van Lear 1982).

Some research suggests that fire may cause multiple sprouting from top-killed advance regeneration (McGee 1979). In addition, some rootstock may be damaged or even consumed by fire, thereby reducing the regeneration potential of the stand. Many questions remain unanswered regarding the effects of broadcast burning on stem quality of regenerating oak stems. However, observations of dozens of hardwood-pine stands regenerated using the "fell and burn" site preparation technique (Abercrombie and Sims 1986) suggests that broadcast burning under carefully prescribed conditions is favorable to quality oak regeneration in the Southern Appalachians.

Intense fires can sometimes result in the introduction of oak in the succeeding stand. Nowacki (1988) documented cases in northern Wisconsin where clearcutting of old-growth maple-hemlock stands and slash burning resulted in even-aged stands dominated by northern red oak. Lorimer (1989) suggested that these oak stands probably developed from acorns brought into the burned area by birds and animals. Similar observations have been made following an intense wildfire in the mountains of South Carolina.

> To Increase Quality and Numbers of Oak Stems after Clearcutting

CONCLUSIONS Oaks in the Southeast are being replaced by other species on better sites where oaks were once dominant. The fire history of this region, biological adaptations of oak and other species to fire, and ecological functions of fire in oak ecosystems strongly suggest that oak replacement on these better sites is largely the result of a fire regime different from that which existed in the region in previous millennia. Until the past half century, frequent fires apparently allowed oak regeneration to accumulate and develop in the open understory of mature stands at the expense of shade-tolerant, fire-intolerant species. When the overstory of these stands was either completely or partially removed by various agents (wind, insects, wildfire, Indian clearing, harvesting, etc.), conditions were created which allowed advance regeneration dominated by oak to develop into mature stands dominated by oak.

If oaks are to be maintained as a dominant overstory species on good quality sites in the Southeast, foresters will have to either restore fire to some semblance of its historical role as a major environmental factor or develop methods that simulate the effects of fire. It will be essential for foresters, as well as the public, to recognize that fire was a major factor shaping the composition and structure of many forest ecosystems.

LITERATURE CITED Abercrombie, J. A., Jr.; Sims, D. H. 1986. Fell and burn for low-cost site preparation. Forest Farmer. 46:14-17.

Abrams, M. D. 1992. Fire and the development of oak forests. BioScience. 42(5):346-353.

- Augspurger, M. K.; Van Lear, D. H.; Cox, S. K.; Phillips, D. R. 1987. Regeneration of hardwood coppice with and without prescribed fire. In: Phillips, D. R., ed. Proceedings Fourth Biennial Silvicultural Research Conference; 1986 November 4-6; Atlanta, GA. Gen Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service: 82-92.
- Beck, Donald E. 1989. Regenerating cove hardwood stands. In: Smith, H. Clay; Perkey, Arlyn W.;
 Kidd, William E., eds. Guidelines for Regenerating Appalachian Hardwoods. 1988 May 24-26;
 SAF Publications 88-03. Morgantown, WV: West Virginia University Books, Office of Publications: 156-166.
- Carvell, Kenneth L.; Korstain, C. F. 1955. Production and dissemination of yellow-poplar seed. Journal of Forestry. 53:169-170.
- Carvell, K. L.; Maxey, W. R. 1969. Wildfire adversely affects composition of cove hardwood stands. West Virginia Agricultural Experiment Station Bulletin. 2(2):4-5.
- Carvell, K. L.; Tryon, E. H. 1961. The effect of environmental factors on the abundance of oak regeneration beneath mature oak stands. Forest Science. 7:98-105.
- Clark, F. Bryan; Watts, Richard. 1971. Silvicultural methods for regenerating oaks. Oak Symposium Proceedings. August 16-20; Morgantown, WV. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.
- Crow, T. R. 1988. Reproductive mode and mechanisms for self-replacement of northern red oak (Quercus rubra)--a review. Forest Science. 34:19-40.
- Darley-Hill, S.; Johnson, W. C. 1981. Acorn dispersal by the blue jay (Cyanocitta cristata). Oecologia. 50:231-232.

- Deen, Robert T.; Hodges, John D. 1990. Oak regeneration in abandoned fields: Presumed role of the blue jay. P. 84-93. In: Proceedings Sixth Biennial Southern Silvicultural Research Conference. Gen. Tech. Rep. SE-70. Vol. I. Asheville, NC: U.S. Department of Agriculture Forest Service, Southeastern Forest Experiment Station. 504 pp.
- De Selm, H. R.; Clebsch, E. E. C.; Nichols, G. M.; Thor, E. 1973. Response of herbs, shrubs, and tree sprouts in prescribed-burn hardwoods in Tennessee. In: Proceedings Tall Timbers Fire Ecology Conference 13; 1973 March 22-23; Tallahassee, FL: 331-344.
- Galford, Jimmy R.; Peacock, John W.; Wright, Susan L. 1988. Insects and other pests affecting oak regeneration. In: Smith, H. Clay; Perkey, Arlyn W.; William E., eds. Guidelines for Regenerating Appalachian Hardwood Stands; 1989 May 24-26; Morgantown, WV. SAF Publication 88-03. Morgantown, WV: West Virginia University Books, Office of Publications.
- Gibson, Lester P. 1972. Insects that damage white oak acorns. Res. Pap. NE-220. Upper Darby,
 PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.
 7 pp.
- Hannah, Peter R. 1987. Regeneration methods for oaks. Northern Journal of Applied Forestry. (4):97-101.
- Harmon, Mark E. 1984. Survival of trees after low-intensity surface fires in Great Smokey Mountains National Park. Ecology. 65(3):796-802.
- Hodgkins, E. J. 1958. Effects of fire on undergrowth vegetation in upland southern pine forests. Ecology. 39:36-46.
- Johnson, Paul S. 1974. Survival and growth of northern red oak seedlings following a prescribed burn. Res. Note NC-177. St. Paul, MN: U. S. Department of Agriculture, Forest Service, North Central Experiment Station. 3 pp.
- Kelty, Matthew J. 1988. Sources of hardwood regeneration and factors that influence these sources.
 P. 17-30. In: Proceedings Guidelines for Regenerating Appalachian Hardwood Stands; 1988
 May 24-26; Morgantown, WV. SAF Publication 88-03. West Virginia University Books, Office of Publications, Morgantown, WV.
- Komarek, E. V. 1974. Appalachian mountains, mesophytic forest, and grasslands. P. 269-272. In: Kozlowski, T.T.; Ahlgren, C. E., eds. Fire and Ecosystems. New York: Academic Press.
- Langdon, O. G. 1981. Some effects of prescribed fire on understory vegetation in loblolly pine stands. In: Wood, G. W., ed. Prescribed fire and wildlife in Southern forests: proceedings of a symposium; 1981 April 6-8; Myrtle Beach, SC. Georgetown, SC: Clemson University, Belle W. Baruch For. Sci. Inst.: 143-153.
- Little, Silas. 1974. Effects of fire on temperate forests: Northeastern United States. In: Kozlowski, T. T.; Ahlgren, C. E., eds. Fire and Ecosystems. New York: Academic Press.
- Loftis, David L. 1988. Regenerating oaks on high sites, an update. In: Smith, H. Clay; Perkey, Arlyn W.; Kidd, William E., Jr., eds. Guidelines for Regenerating Appalachian Hardwood Stands. 1988 May 24-26; Morgantown, WV. SAF Publication 88-03. West Virginia University Books, Office of Publications, West Virginia: 199-209.
- Loftis, L. David. 1990. A shelterwood method for regenerating red oak in the southern Appalachians. Forest Science. 36:917-929.
- Loomis, Robert M. 1973. Estimating fire-caused mortality and injury in oak-hickory forests. Research Paper NC-94. St. Paul, Minnesota: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 6 pp.

- Lorimer, Craig G. 1989. The oak regeneration problem: New evidence on causes and possible solutions. University of Wisconsin, Deptartment of Forestry, Forest Resource Analyses #8. 31 pp.
- Marquis, D. A.; Ekert, P. L.; Roach, B. A. 1976. Acorn weevils, rodents, and deer all contribute to oak regeneration difficulties in Pennsylvania. Res. Pap. NE-356. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 5 pp.
- Martin, Robert E.; Mitchell, Russel G. 1981. Possible, potential, probable, and proven fire-insect interactions. P. 138-144. In: Proceedings of the 1980 Convention of the Society of American Foresters.
- Martin, William H. 1989. The role and history of fire in the Daniel Boone National Forest. Winchester, KY: U. S. Forest Service, Daniel Boone National Forest.
- Maslen, Paul. 1989. Response of immature oaks to prescribed fire in the North Carolina piedmont.
 P. 259-266. In: Proceedings of the 5th Biennial Southern Silvicultural Research Conference; 1988 November 1-3; Memphis, Tenn. Gen. Tech. Rep. SO-74. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 618 pp.
- McGee, C. E. 1979. Fire and other factors related to oak regeneration. In: Holt, H. A.; Fischer, B. C., eds. Regenerating oaks in upland hardwood forests. Lafayette, IN: John S. Wright Forestry Conference, Purdue University: 75-81
- Nowacki, G. J. 1988. Community and historical analysis of northern red oak along a moisture gradient in north-central Wisconsin. University of Wisconsin: Stevens Point. Thesis.
- Roth, E. R.; Hepting, G. H. 1943. Origin and development of oak stump sprouts as affecting their likelihood to decay. Journal of Forestry. 41:27-36.
- Roth, E. R.; Sleeth, Bailey. 1939. Butt rot in unburned sprout oak stands. Tech. Bull. 684. U.S. Department of Agriculture. 42 pp.
- Sander, Ivan L. 1988. Guidelines for regenerating Appalachian oak stands. P. 189-198. In: Proceedings Guidelines for Regenerating Appalachian Hardwood Stands. 1988 May 24-26; Morgantown, WV. SAF Publication 88-03. Morgantown, WV: West Virginia University Books, Office of Publications.
- Sander, Ivan L.; Johnson, Paul S.; Watt, Richard F. 1976. A guide for evaluating the adequacy of oak advance reproduction. Gen. Tech. Rep. NC-23. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 7 pp.
- Sander, Ivan L.; McGee, Charles E.; Day, Kenneth G.; Willard, Ralph E. 1983. Oak-Hickory. In: Russell M. Burns (comp.) Silvicultural systems for the major forest types of the United States. Agriculture Handbook No. 445. Washington, D.C.: U.S. Department of Agriculture, Forest Service: 116-120.
- Sanders, D. L.; Van Lear, D. H.; Guynn, D. C. 1987. Prescribed burning in mature pinehardwood logging slash in the Southern Appalachians. In: Phillips, Douglas R. (comp.) Proceedings of the 4th biennial silvicultural research conference; 1986 November 4-6; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 93-96.
- Shearin, A. T.; Bruner, Marlin H.; Goebel, N. B. 1972. Prescribed burning stimulates natural regeneration of yellow-poplar. Journal of Forestry. 482-484 pp.
- Teuke, M. J.; Van Lear, D. H. 1982. Prescribed burning and oak advance regeneration in the Southern Appalachians. Res. Pap. 30. Georgia Forestry Commission. 11 pp.

- Thor, E.; Nichols, G. M. 1974. Some effects of fire on litter, soil, and hardwood regeneration. Proc. Tall Timbers Fire Ecology Conference. 13:317-329.
- Van Lear, D. H. 1990. Fire and oak regeneration in the Southern Appalachians. In: Nodvin, Stephan C.; Waldrop, Thomas A., eds. Fire and the Environment: Ecological and Cultural Perspectives. Gen. Tech. Rep. SE-69. U. S. Department of Agriculture, Forest Service: 15-21.
- Van Lear, David H.; Waldrop, Thomas A. 1989. History, uses, and effects of fire in the Appalachians. Gen. Tech. Rep. SE-54. U.S. Department of Agriculture, Forest Service. 20 pp.
- Waldrop, Thomas A.; Van Lear, David H.; Lloyd, F. Thomas; Harms, William R. 1987. Longterm studies of prescribed burning in loblolly pine forests of the Southeastern Coastal Plain. Gen. Tech. Rep. SE-45. U.S. Department of Agriculture, Forest Service. 23 pp.
- Waldrop, Thomas A.; Lloyd, F. Thomas. 1991. Forty years of prescribed burning on the Santee fire plots: Effects on overstory and midstory vegetation. In: Nodvin, Stephan C.; Waldrop, Thomas A., eds. Fire and the Environment: Ecological and Cultural Perspectives. Gen. Tech. Rep. SE-69. U. S. Department of Agriculture, Forest Service: 15-21.
- Wells, Carol G.; Campbell, Ralph E.; Debano, Leonard F. 1979. Effects of fire on soil. Gen. Tech. Rep. WO-7. Washington, D.C.: U.S. Department of Agriculture, Forest Service, Washington Office. 34 pp.
- Wendel, G. W.; Smith, H. Clay. 1986. Effects of a prescribed fire in a Central Appalachian oakhickory stand. NE-RP-594. U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 8 pp.

Flowering and Oak Regeneration

Robert A. Cecich, North Central Forest Experiment Station, USDA Forest Service, Columbia, MO 65211

ABSTRACT Episodic acorn production appears to be the norm for the genus *Quercus*. This year-to-year variation has been associated with the number of pistillate flowers, the supply of pollen, weather, insects, nutrition, and genetics. Acorns are the fruit of the oak tree, developing from the pistillate flower. Depending upon successful pollination and fertilization by pollen produced in the staminate flowers, one of six ovules will survive and grow; the other five will abort. We usually notice the tiny pistillate flower after fertilization when the cupule and pericarp start to enlarge and begin to look like an acorn. This is also about the time when acorn weevils begin to oviposit. It is a long, arduous journey from the emergence of the flowers in the spring to their maturation as acorns. The appearance of abundant pistillate flowers does not guarantee a successful acorn crop. Can we predict an acorn crop prior to a few weeks before seedfall? Not reliably. Can we predict the appearance of a flower crop? Not yet, and not before we understand clearly how biotic and abiotic factors affect flower initiation and development in individual species.

When the subject of "flowering" in oaks arises, we tend to think about acorn production. Although these are two different topics, they are part of the continuum where flowering leads to acorn production. This paper dissects that continuum into component processes such as flower initiation, flower development, and embryology, in the context of addressing the following questions:

- 1. What should a forester know about flowering?
- 2. What factors affect flowering?
- 3. What is the relationship between flowers and acorns?
- 4. Can we predict flower crops?

WHAT A FORESTER SHOULD KNOW ABOUT FLOWERING

Pistillate Flowers

The oak pistillate flower begins its journey to becoming an acorn when the meristematic tissues of the bud receive a signal, the as yet unknown flowering stimulus, that directs axillary primordia in some of the leaves to become an inflorescence stalk and not a vegetative bud (Minina 1954, Turkel and others 1955, Romashov 1957, Merkle and others 1980). These researchers could not differentiate the stalk from the bud until late summer when bud scales began to arch over the vegetative bud primordium. In contrast, the inflorescence primordia remain relatively naked, having only one or two bracts (Merkle and others 1980), and may be somewhat larger than the vegetative bud (Turkel and others 1955). The conclusions of Turkel and others (1955) and Merkle and others (1980) are based on microscopic observation of prepared specimens of *Quercus alba* (table 1).

-	
Latin names	Common names
North Ame	rican
Q. alba L.	White oak
coccinea Muenchh.	Scarlet oak
gambelii Nutt.	Gambel oak
ilicifolia Wangenh.	Bear oak
prinus L.	Chestnut oak
rubra L.	Northern red oal
velutina Lam.	Black oak
European d	and Asian
Q. aegilops L.	
ilex L.	
macrolepis Ky. (Probably a su	bspecies of Q. aegilops)
myrsinaefolia Blume	
petraea (Mattushka) Lieblein (pubescens W.	(Includes Q. sessiliflora)
robur L. (Includes Q. peduncu trojana Webb	ılata)

Table 1-Species of Quercus discussed in this paper

Minina (1954) and Romashov (1957), viewing fresh material of Q. robur under a dissecting microscope, could not identify the inflorescence primordia until late winter or early spring. Bonnet-Masimbert (1978) could not identify the flowers of Q. pedunculata (robur) or Q. sessiliflora (petraea) during the winter.

Pistillate inflorescence development in *Q. alba* resumes in late March (Merkle and others 1980). The axis begins to elongate and several additional bracts are produced in a spiral. In the axil of each bract a pistillate flower begins to differentiate, but, in general only the lower ones became functional. There are usually 2-3 functional flowers, with a range of from 1-5. In *Q. rubra*, pistillate flowers form in the axils of two lower, opposite bracts, and occasionally only one is formed (Sattler 1973).

The ontogeny or sequence of development of the oak flower parts from the floral apex of Q. *rubra* was described in detail by Langdon (1939) and Sattler (1973). Six perianth primordia are initiated, three outer and then three inner. These primordia are elevated by meristematic activity beneath them. Three gynoecial (carpel) primordia appear on the apex opposite the three outer perianth primordia and they grow together laterally to eventually become the stigmas. As growth continues upward, the area beneath becomes the ovary wall. Conrad (1900) observed that carpels are already evident in winter buds of Q. velutina. Thus, there may be much variation between species in the same subgenus.

The young ovary closes as the gynoecial primordia are carried up with the ovary wall. Concurrently, the growth between and at the base of the gynoecial primordia initiates the septa. Three septa are formed and become appressed at their upper, inner margins; they are not joined at their base. Two placentae form initially as slight protrusions along the base and on each side of the septa. Thus, in each locule there are two placentae—one from each septum. When the ovary cavity of Q. myrsinaefolia is closed, the upper portion remains uniloculate, having three separate septal protrusions, but is triloculate at the basal region (Okamoto 1982).

The ovule of Q. rubra develops as a further enlargement of the placental bulge, but the timing in relation to anthesis is unknown (Sattler 1973). Okamoto (1982) found that the ovules of Q. myrsinaefolia are not initiated at anthesis, while Turkel and others (1955) found that ovule development begins at that time. However, the timing of ovule development in the erythrobalanus (red oak) subgenus may differ from the lepidobalanus (white oak) subgenus.

What is an ovule? Botanically speaking, an ovule is a megasporangium; i.e., a structure that bears the megaspore mother cell (MMC) (Davis 1966). The MMC undergoes meiosis or reduction-division, producing four haploid cells, only one of which survives to become the functional megaspore. By a series of mitotic divisions the megaspore gives rise to the megagametophyte or embryo sac, an eight-nucleate structure at the tip of the nucellus. The nucellus is partly covered by the inner and outer integuments. When the integuments have elongated over the end of the nucellus, the "hole" that is formed is called the micropyle. This is the route through which the pollen tube approaches the embryo sac (Benson 1894). Major food reserves of starch and lipids are located almost exclusively within the outer integument, while the inner integument is virtually void of food reserves (Mogensen 1973). Mogensen proposed that the pathway for food materials in the ovule is from the outer integument to the chalaza (basal portion of the ovule) and then through the postament (central core) of the embryo sac.

The study of embryo sac formation in plants has received much attention from botanists. Conrad's (1900) study of *Q. velutina* is considered the initial study of this structure in the genus *Quercus*. That study briefly described the pattern of nuclear division that yields an eight-nucleate embryo sac at the tip of the nucellus. It was 50 years before the next papers on embryo sac development appeared. These included investigations of *Q. macrolepis* (Bagda 1948, 1952), *Q. robur* (Hjelmqvist 1953), *Q. alba* (Turkel and others 1955), *Q. ilex* (Corti 1959), *Q. aegilops* (Scaramuzzi 1960), and *Q. trojana* (Bianco 1961).

Once the embryo sac is formed, fertilization of its egg and central cell via germinating pollen must occur for seed development to continue. Unfortunately, the details of pollen tube growth through the stigma and stylar tissues are not documented for any species of *Quercus*. Some observations suggest that pollen tube growth does not proceed for several weeks after the pollen grains land on the stigmatic surface and that pollen germination waits for the ovule to complete development (Jovanovic and Tukovic 1975). Notwithstanding, Mogensen (1972) provided the only detailed evidence of pollen tube invasion through the micropyle and into the egg apparatus of the embryo sac of *Q. gambelii*. Upon reaching the embryo sac, the branched pollen tube grows along one of the synergids and penetrates it by growing through the filiform apparatus. The pollen tube opens at its tip and releases its contents. Although fertilization occurred, Mogensen (1972) did not observe the isolated male gametes or their union with the egg nucleus or the central cell.

Following fertilization, a free-nuclear endosperm grows before the first division of the zygote occurs (Hjelmqvist 1953, 1957; Brown and Mogensen 1972). In general, as the endosperm becomes cellular, the embryo begins to differentiate (Singh and Mogensen 1976), moving quickly through the heart-shaped stage. Singh and Mogensen (1976) concluded that the endosperm does not have a major function as a food storage tissue, but rather may serve as a translocating tissue to the embryo. During zygote and early embryo stages, lipids may be more important as a nutrient source, while starch is probably utilized at later stages of embryo development (Singh and Mogensen 1975).

Information about embryo and cotyledon growth is very limited. Mogensen (1965) provided the most detailed picture in his comparative study of Q. alba and Q. velutina. One major difference he noted was that the epicotyl apex of Q. alba produced from three to five leaf primordia prior to acorn maturity, while Q. velutina produced none. Stairs (1964) also found no leaf primordia in mature embryos of Q. coccinea. Cecich (unpubl. data) found large concentrations of druse crystals in cotyledons of Q. velutina and Q. rubra, but none in Q. alba. The relative unpalatability of erythrobalanus acorns, considered to be related to increased content of lipids and tannins (Goodrum and others 1971, Short 1976, Smallwood and Peters 1986, Smith and Follmer 1972), may also be related to irritations caused by these large crystals.

It is difficult to provide a generalized calendar of events from ovary development through acorn maturation. Variation among and within species, geographic location, weather conditions, and sampling problems would make the results of that task questionable. Merkle and others (1980) provided an example of this variation when they compared their observations of Q. *alba* ontogeny with those of Turkel and others (1955). Nevertheless, the time of pollen shed is probably the best local index for the beginning of the seed production cycle but, does not include flower initiation and development within the bud. In the lepidobalanus group the pollen tube fertilizes the egg about 4-6 weeks after initiating growth. Variation in the time of embryo growth and maturation of the acorn occurs and, depending upon species, acorns drop over a several-month period (USDA 1974).

Staminate Flowers The first sign of differentiation of the staminate inflorescence primordium ranged from late May (Merkle and others 1980) and early June (Turkel and others 1955) in *Q. alba* to June and July in *Q. robur* and *Q. petraea* (Minina 1954, Romashov 1957, Jovanovic and Tucovic 1975). Similar observations are not available for species in the erythrobalanus subgenus.

The inflorescence, which is inserted in the axil of a bud scale and not a leaf (Minina 1954), is without appendages until late June or early July, when meristematic areas appear on the axis (Merkle and others 1980). These floral apex primordia appear before or coincident with the subtending bract primordia (Turkel and others 1955). However, Sattler (1973) found that the floral apex of Q. rubra is initiated in the axil of the bract, just the opposite of Q. alba.

On the flank of each floral apex, the perianth primordia appear, fusing into a single perianth as they grow. The stamen primordia appear on the apex opposite the

perianth members in mid to late July. By fall, these stamen primordia grow into immature anthers and filaments. The overwintering condition of the slightly-lobed anther is that of a homogeneous parenchymatous mass (Turkel and others 1955).

Resumption of anther development in the spring varies by species and location: early March (Bonnet-Masimbert 1984), late March (Merkle and others 1980), April (Turkel and others 1955), and mid to late April (Conrad 1900). The parenchymatous mass differentiates into the sporogenous mass and the parietal layers. The number of sporogenous cells increases mitotically and eventually become the microspore mother cells which undergo meiosis to become microspores and, finally, pollen grains. Stairs (1964) has provided the only account of meiosis in *Quercus*. During the meiotic process, the parietal layers differentiate into the tapetum and the anther wall. Dehiscence of pollen grains occurs about 6 weeks after resumption begins (Turkel and others 1955). Before leafing out, the staminate inflorescence, bearing numerous staminate flowers, elongates and emerges from the bud scales as the familiar catkin (Vogt 1969).

There are conflicting observations about the dynamics of pollen tube growth after the pollen lands on the stigmatic surface. For instance, pollen germination in Q. *robur* was completed within 24 hours, but fertilization occurred 6-7 weeks later (Jovanovic and Tukovic 1975). Benson (1984) didn't find pollen tubes in Q. *robur* until just before fertilization. In contrast, Allard (1932) observed that: "When pollen reaches the stigma of members of the white oak group, the growth of the pollen tube containing the male cells follows an uninterrupted advance into the tissues of the style until the ovules are fertilized." These two extremes of pollen tube behavior for members of the same subgenus suggest that we should investigate this anomaly to better understand the among-species variation. Allard also found that in the red oak group the pollen tubes cease growth at the base of the style until the following spring when fertilization of the ovules occurs.

Pollen tube growth may also be affected by temperature. Pollen of Q. pubescens requires a higher temperature for germination (38 °C) than Q. robur (20 °C). Low and erratic fertility of Q. pubescens in cold habitats may be attributed to the prevention of pollen germination. This temperature limit may also explain the northern boundary for the species range (Jicinska and Koncalava 1978).

Flower Distribution The distribution of various bud types on a branch has been in *Q. robur* and *Q. petraea* (Minina 1954; Romashov 1957; Bonnet-Masimbert 1978, 1984). Minina and Romashov described five types of buds: simple male, male and vegetative mixed, female and vegetative mixed, complex (male, female, and vegetative), and vegetative only (active or dormant). Bonnet-Masimbert recognized six types: vegetative, vegetative and male, male only (rare, but occur in years of heavy flowering), vegetative and female, vegetative and hermaphroditic (not the same as Minina's complex type). Bonnet-Masimbert's sixth type is the latent bud which Minina puts under the vegetative category. Buds containing uncommitted primordia were considered important in reconstituting a branch system after insect attack.

Many researchers of oak reproductive biology have concluded that flowering is irregular from year to year without mentioning the variation in distribution of bud

types. Bonnet-Masimbert determined that a large acorn crop is related to a large number of male flower buds, manifested through an increase in hermaphroditic buds and, to a lesser extent, to the number of female flowers. When the acorn crop was poor, large numbers of buds in the potentially male zone on a branch did not evolve.

Schlarbaum and Rhea (unpubl. data) evaluated flowering in a 17-year-old Q. rubra seedling seed orchard in Tennessee. They selected a light-, medium-, and heavy-flowering tree and counted all pistillate flowers. The majority of flowers were located in the upper one third of the crown of each tree, and there were no differences in flower numbers among the quadrants in a crown. Similarly, Jovanovic and Tukovic (1975) cited observations by Rempe (1937) and Piatnitsky (1954) that the greatest quantity of pollen was produced in the upper part of the crown. Sharp and Chisman (1961), however, found that pollen catkins were evenly distributed across the crown of Q. alba. I have also observed the latter in Q. alba, Q. rubra, and Q. velutina.

FACTORS AFFECTING FLOWERING

Phase Change and Flowering

One of the difficulties in forest tree breeding and flowering research is the long time between generations. Most oaks take 15 to 25 years to reach minimum seed-bearing age (USDA 1974). Reducing that time would reduce the length of a generation cycle and make it possible to increase genetic gain-per-unit-time by making earlier selections (Lambeth 1980). If earlier flowering can be attained in oak species, it would be possible to generate inbred lines, whose usefulness has been demonstrated in many crops. Inbreeding not only increases heritability and facilitates selection, but it also helps to reduce the "genetic noise" common to physiology research. Although inbreeding has been successfully demonstrated in *Quercus* (Irgens-Moller 1955), Jovanovic and others (1971) did not get successful embryo development in self-pollinated oak flowers.

It is generally accepted that the change from a juvenile to a mature state (Poethig 1990) in forest trees occurs at the time of first flowering (Zimmerman 1972). A juvenile tree is not capable of flowering because it is not able to respond to stimuli that would otherwise induce flowering; while a mature tree may not flower because of the absence of the stimuli or genetic control causing sterility. Thus, the onset of flower production is used as an indicator that the juvenile phase has ended and that the mature phase has begun (Wareing 1959).

How do you get a tree to pass from the juvenile to the mature phase sooner? A common strategy is to grow seedlings in an environment that greatly increases growth rate. Some of the cultural methods used to do this include elevated temperatures, long photoperiods, adequate water, and fertilization. Thus, size *per se* is positively correlated with early flowering.

The inheritance of early flowering in forest trees has been demonstrated in *Betula* verrucosa (Johnsson 1949) and in *Pinus sylvestris* (Teich and Holst 1969). A dominant major gene has been implicated in controlling early flowering in both species. Because early flowering is inherited, selection pressure can be applied to increase the frequency of progenies that flower early. In jack pine (*Pinus*

banksiana Lamb.) the larger trees in a family are more likely to flower first (1 year from seed) (Bolstad and others 1991). Once again, the size of the tree is positively correlated with the attainment of early flowering.

An alternative to waiting for seedling-origin plants to produce flowers is to use sprouts. Wolgast and Stout (1977) used sprouts in clearcuts to determine the earliest age at which acorns of Q. *ilicifolia* could be produced. Pistillate flowers appeared on sprouts at the beginning of the second growing season, indicating that flower primordia were initiated during the first season. Acorns were mature at the end of the third season. Sharik and others (1983) also found that stems of coppice-origin Q. prinus first produced acorns in the third season, compared to 20 years for seedlings. Advance reproduction seedlings and seedling-sprouts of the same age produced no acorns.

Components of the Flowering Process There are three major components to the flowering process in oaks: (1) Initiation, (2) Differentiation of the staminate and pistillate inflorescences and their flower primordia, and (3) Emergence of the flowers, receptivity of the stigmas, and shedding of pollen. Let us examine how various factors may affect these components.

Initiation refers to how chemical, genetic, and abiotic factors interact during a critical time period to cause a cell or meristem to commit itself to become a flower or flower part. This is not the same as differentiation wherein the structural manifestation of the initiation process occurs; e.g., the appearance of the staminate inflorescence in late May (Merkle and others 1980). Most of what we know about flower initiation is based on research with annual plants (Evans 1969, Bernier 1988). However, woody plants behave differently. They have long juvenile periods during which they do not flower, even though the proper environmental stimuli may be present. In addition, annuals tend to have terminal flowers, that is, the shoot apical meristem *per se* becomes a flower. In woody plants, which must grow year after year, the flowers are normally axillary. One way of defining the period of flower initiation in conifers is through the use of plant growth regulators, primarily the gibberellin $A_{4/7}$ mixture (GA_{4/7}) (Owens and Blake 1985). Unfortunately, GAs rarely induce flowering in hardwoods and there is no known method for reliably doing so. This includes the application of mineral fertilizers, which have no apparent direct effect on flowering, although there may be an indirect effect from a correlated increase in crown vigor. Wolgast and Stout (1977) did note a positive fertilizer response in *Q. ilicifolia*. Any speculation about how a factor may be related to flower initiation must be accompanied by a notation of timeliness; i.e., during what specific time period does that factor operate. That notation does not now exist and so the discussion of flower initiation becomes almost fruitless (no pun intended).

Differentiation of reproductive structures many extend from late May (Merkle and others 1980) to the time of pollen shed and female receptivity about 1 year later. So how does one realistically ascribe the success or failure of the development of a flower crop over that length of time to factors such as weather? Except for deep freezes in late spring (Sharp 1958, Sharp and Sprague 1967, Goodrum and others 1971, Wolgast and Trout 1979), does the weather affect differentiation? Sharp and

Chisman (1961) found that *Q. alba* on 10 sites produced good-to-heavy pollen crops each year and that varying numbers of pistillate flowers were produced. Since staminate flowers were produced abundantly each year, did any factor inhibit their development during that year of differentiation? If a pistillate flower crop in a given spring is rated as poor, was it caused by biotic or abiotic factors during development or were the flower primordia ever initiated? Sharp and Sprague (1967) speculated that a warm late April and cool early May were related to early catkin emergence and delayed pollen dispersal, respectively. Under that weather scenario, pollen shed and pistillate flower receptivity were considered to be more closely aligned. They provided no anatomical observations to confirm their speculations.

Genetic control over seed production in oaks has been demonstrated by a number of investigators. However, Farmer (1981) found that in a given year seed production among clones of northern red oak was most highly correlated with the percentage of pistillate flowers that were fertilized, while year-to-year differences were associated with variation in the number of flowers. He believed that fecundity could be increased by selecting high-yielding clones in a grafted orchard. Ledig and others (1971) and Wright (1953) also found much tree-to-tree variation in reproductive ability. Grafting of oak scions selected from mature, flowering individuals can be readily accomplished and, thus, flowers can be made quickly available (Irgens-Moller 1955).

Floral sex ratios in *Q. ilicifolia* changed with position of the tree on a slope (Aizen and Kenigsten 1990). At the top of the slope tall stems had most of the male flowers. While at the bottom of the slope, there was no height relationship but there were fewer male flowers on all trees than there were pistillate flowers. The authors could speculate only that a change in temperature along the gradient influenced the physiological basis for sex allocation. Only stems at the top of the slope had second-year acorns, perhaps related to the increase in pollen availability at the top.

Emergence, Receptivity, and Shedding. Emergence of the staminate inflorescence and shedding of pollen are known to increase or hasten with rising temperatures and to drop with decreasing temperature (Romashov 1957). Rainy weather, associated with decreased temperature, also reduced pollen dispersal. The success of the acorn crop has not always been related to pollen dispersal. Sharp and Chisman (1961) concluded that pollen dispersal occurred when relative humidity dropped and remained below 45 percent for several hours, but they did not mention the success of the acorn crop. Similarly, Jovanovic and Tukovic (1975) cited European literature indicating that pollen grains separated better when relative humidity was lowest.

Wolgast (1972) explored the effect of relative humidity experimentally. In a series of growth chamber experiments using Q. *ilicifolia*, he demonstrated that relative humidity at the time of pollen shed and stigma receptivity can limit the size of an acorn crop. No acorns matured when relative humidity exceeded 61 percent, but about half the flowers matured into acorns when relative humidity was lower. While hot, dry winds in early May caused dessication of pollen catkins, Sharp and

Sprague (1967) found no correlation with relative humidity and acorn yields in field studies and concluded that temperature was a primary factor in acorn crops.

Probably the most important factor controlling the emergence of pistillate flowers and their receptivity is temperature, which directly or indirectly influences flower emergence through branch and leaf elongation. Low air temperatures were associated with a delay in development of pistillate inflorescences of Q. robur near Moscow (Minina 1954). However, Goodrum and others (1971) concluded that the influence of low temperatures on flowering, setting of fruit, and subsequent acorn yield was inconclusive. Sharp (1958) also concluded that low temperatures in the spring did not affect flowering unless there was a freeze sufficient to damage shoots and leaves.

RELATIONSHIP BETWEEN FLOWERS AND ACORNS

Given that staminate and pistillate flowers have been initiated, differentiation completed, and pollen shed on the receptive stigma, what factors then affect how the pollinated pistillate flowers develop into acorns?

Until now I have presented information about oak flowering in a positive context; i.e., how the primordium originates and develops into a flower. But we also need to look at subsequent events in a somewhat negative context; i.e., how various factors may lead to the abortion of flowers or ovules and, thus, reduce the size of a potential seed crop. Several authors have concluded that the size of an acorn crop is not related to the size of a flower crop. The appearance of numerous pistillate flowers in the spring does not guarantee numerous acorns (Sharp 1958, Sharp and Chisman 1961, Wright 1953, Gysel 1956, Cecich and others 1991). What then are the factors or events related to the loss of the flowers? How are these factors and their timing related to the developmental chronology of the pollinated flower as discussed earlier? Information for answering these questions is limited.

Kossuth (1974) concluded that abscissions occurring during the first 11 weeks after anthesis in Q. alba were probably determined at or before receptivity by degeneration of the ovary. Although ovules had just differentiated at the time of receptivity, ovule growth did not appear to influence premature acorn abscission before or after anthesis. She confirmed the observations of Turkel and others (1955) that an abscission layer begins to form during anthesis between the ovary wall and the receptacle (cupule), and considered this to be an abscission layer analagous to that of a leaf petiole. It is my opinion that this abscission layer is simply the beginning of the separation of the seed coat (pericarp) and cupule manifested in a mature acorn.

Kossuth (1974) noted certain developmental thresholds associated with flower abortion. If ovules developed normally through anthesis, the first threshold was passed and development continued. The failure of megasporogenesis in all ovules of a flower seemed to be determined by anthesis, but abscission was not immediate. A second threshold following fertilization was related to the growth of the functional ovule. Most embryo abortion occurred before the cotyledons were onethird their final size. After a series of leaf size and photosynthesis measurements, Kossuth concluded that expanding leaves were associated with acorn retention and that leaves and developing flowers were not competitive sinks for assimilates. By the time of pistillate flower receptivity, the half-grown leaves were net photosynthesizers.

Kossuth's observations of flower survival and abortion agree with other studies, such as Williamson (1966), who found that about 90 percent of the premature flower abscissions occurred by the time of fertilization, Kossuth's second threshold. My observations of flower abortion in Q. alba agree with this. Fertilization in Q. alba in mid-Missouri occurs during late June, as noted by the presence of an endosperm in the embryo sac. By the first week of July a cellular endosperm and embryo are present (unpublished data). In 4 consecutive years, the percentage of flowers aborted by the first week of July was 98, 76, 83, and 90 percent, respectively. The number of maturing acorns per year, as a percent of the initial flowers, ranged from 0 to 6 percent. Clearly, most of the potential acorn crop is lost by the time of fertilization. Late-spring freezes can also kill second-year pistillate flowers in the erythrobalanus group and prevent them from maturing into acorns (Wolgast and Trout 1979). The impact of drought on either differentiation of flowers or on developing acorns is inconclusive (Sharp and Sprague 1967). Fungi probably have no effect on fertilization success, even though they are widely found on stigmas (Kolpak and others 1980).

The literature suggests that the abortive ovule of Quercus develops a normal embryo sac, fertilization occurs, and a zygote or proembryo stage may be reached (Stairs 1964, Mogensen 1965). However, Mogensen (1975) showed that there are several possible pathways for abortion, even within the same ovary. In observing O. gambelii he found that fertilization to the normal embryo sac failed to occur 45 percent of the time. Ovule abortion caused by failure of the zygote or embryo accounted for 28 percent of the ovules. In 26 percent of the ovules examined, an embryo sac failed to develop within the nucellus. Occasionally, embryo sacs were found without any cellular contents, but these could have been preparation artifacts. He also confirmed this classification in Q. alba and Q. velutina. Mogensen (1975) could but speculate on how only one ovule per ovary develops into a seed. Because only ovules with a normal embryo sac have the potential to become a seed, he speculated that the first ovule in an ovary to be fertilized suppresses further development of the remaining ovules by producing a growth regulator. The capability today to culture ovules in vitro, while varying growth regulator and nutritional components, offers some opportunities to examine this hypothesis.

Insects also are a factor in acorn loss. Most literature deals with the impact of weevils (*Curculio* spp. and *Conotrachelus* spp.) on acorn crops (Gibson 1964, Kearby and others 1986). But weevils don't generally oviposit until midsummer after fertilization and embryogenesis has begun in the pistillate flower. Most flower loss occurs by the time embryogenesis has begun (Williamson 1966, Kossuth 1974, Cecich and others 1991), suggesting that weevils are not the major cause of poor acorn crops. For example, based on the 1990 data for *Q. alba*, about 80 percent of the flowers have aborted by mid-July (Cecich and others 1991). Assuming mid-July as the time of weevil oviposition, only 20 percent of the potential crop is present. Even if the weevils destroyed all the remaining young

acorns, they would not destroy 100 percent of the potential acorn crop as has been asserted, only 20 percent. The major loss occurs before fertilization.

There is another group of insects that also deserves attention in respect to its impact on acorn crops. Treehoppers (Membracidae) are sucking insects that, depending on the species, spend most of their life cycle in the crowns of oak trees (Kopp and Yonke 1973 a,b,c, 1974). The older larval stages and young adults feed primarily on meristematic or succulent tissues and flowers. I have observed them in May and June feeding on oak pistillate flowers by inserting their stylets into the stigma of the flower. One week later these flowers were dying, dead, or abscised (Cecich and others 1991). We are presently conducting controlled feeding experiments to better define the potential impact of this insect family on flower abortion. Their possible involvement in flower loss is plausible because the insects' feeding activity coincides with the time of most of the flower abortions in early May to mid-June.

PREDICTING FLOWER CROPS

Can we predict the size or availability of a flower crop? Probably not, at least with today's information. Predicting an acorn crop may be easier than predicting a flower crop because there are visible indicators-the number of pistillate and staminate flowers and inflorescenses (Gysel 1958, Feret and others 1982). Feret and others (1982) concluded that production of Q. alba acorns could be best predicted from the number of peduncles borne per shoot. They found that this single variable accounted for 84 percent of the observed variability in acorn production. In contrast, I am unaware of any physiological or structural predictive indicators of flowering potential in oaks. As discussed earlier, many factors can impinge upon flower initiation and development, because of the length of time a flower primordium is exposed to potentially disruptive factors. Nevertheless, we could make the assumption that, once a flower is initiated, the pathway for its development will be continuous and successful. This "all-or-none" hypothesis can be tested by sampling a population of buds in early fall and microscopically searching for inflorescence primordia. In late winter or early spring, sample branches could be put into bottle culture indoors, forcing the buds to flush so that flowers could be observed in leaf axils. The number of inflorescences could be compared to the fall count and total number of flowers determined. Subsequently, these values would be compared to inflorescence and flower numbers from intact branches on the source tree. A significantly lower flower or inflorescence count at anthesis could indicate that either the process was disrupted or that sampling was inadequate.

An alternative hypothesis, and one that is probably more realistic, is that there are many steps to successful flower emergence in oak, beginning with inflorescence initiation. From research on annual plants, we know that there are many genes that regulate flower development (Bowman and others 1989, Shannon and Ry Meeks-Wagner 1991, Smyth and others 1990). It is not unreasonable to assume that similar genes control the development of various flower components in oak. Testing for the specific genes would require traditional breeding experiments, utilizing one or more individuals that possess a stable flower development mutation, so that the progeny could be evaluated. Unfortunately, a consequence of breeding oaks is the many years of waiting for the progeny to flower; i.e., we must wait for the juvenile-mature phase change to occur. However, current molecular technologies, such as Restriction Fragment Length Polymorphisms (RFLPs), cDNA hybridization, Random Amplified Polymorphic DNA (RAPD), are available that utilize vegetative tissues and may help to identify or locate specific flower development genes. If there were a mutation that leads to cessation of flower development, there would be no viable flower and one could not discriminate that pathway from the disrupted "all-or-none" pathway. This assumes that phase change has occurred. However, a change in gene structure (a mutation) could still be identified with these technologies.

CONCLUSION The intent of this paper was to provide a comprehensive review of the literature and synthesize it into a plausible "story" of oak flowering biology. Easier said than done! Although there is much information about certain topics within the subject area of flowering biology, the information voids are even larger. A measure of the paucity of information about oak flower biology can be shown by restating the four questions put forth in the beginning and seeing how well they were answered.

What Should a Forester Know About Flowering? There is still much to learn about the anatomy and physiology of oak flowers, especially the staminate flowers that produce pollen. The limited information we have from only a few species indicates that flower structure varies among species and subgenera. For *Q. alba*, the most-studied North American species, there is no complete "story" describing the continuum from initiation through seed maturation.

What Factors Affect Flowering? Reviewing this literature was easy; there wasn't much. The long juvenile phase is important to the forest manager who wants to enhance or encourage acorn production in a young stand as soon as possible after harvesting. There is still going to be a long wait. We still don't know when initiation occurs, although the first sign of flower differentiation has been observed in two lepidobalanus species. Genetic control over the amount of flowers and seed production has been demonstrated. Emergence, receptivity, and pollen shed appear to be controlled by weather, especially temperature and relative humidity.

What is the
Relationship Between
Flowers and Acorns?Pistillate flowers don't necessarily produce acorns. In fact, most of the time they
don't; which is why we're concerned about this problem. Developing acorns and
leaves are apparently not competitive sinks for photoassimilates. Most flower
abortion occurs before weevil oviposition; however, weevils can destroy the
remainder of the crop.

Can We Predict Flower Crops? There are no published accounts of how to predict the appearance or number of oak flowers. Because we lack so much information about flower initiation and development within the bud, it is not reasonable to estimate the time required for making reasonable predictions. Before we can interpret how environmental, physiological, and genetic factors influence the production of flowers and acorns, we must develop a solid understanding of the flowering biology of individual species. The fragmented information for oak flowering does not lend itself to developing a clear picture of the flowering process, but it does indicate where the shortcomings exist. And, this provides guidance and opportunities for asking meaningful questions and doing the appropriate research.

LITERATURE CITED Aizen, Marcelo A.; Kenigsten, Alejandra. 1990. Floral sex ratios in scrub oak (Quercus ilicifolia) vary with microtopography and stem height. Canadian Journal of Botany. 68:1364-1368.

- Allard, H.A. 1932. A progeny study of the so-called oak species *Quercus saulii*, with notes on other probable hybrids found in or near the District of Columbia. Bulletin of the Torrey Club. 59(2):267-277.
- Bagda, von Haydar. 1948. Morphologische und biologische untersuchungen uber Valonea Eichen (Quercus macrolepis Ky.) im Haci-Kadin-Tal bei Ankara. Communication de Faculte des Sciences de l'Universite d'Ankara. 1:89-125.
- Bagda, von Haydar. 1952. Untersuchungen uber den weiblichen Gametophyten der Valonea-Eiche (Quercus macrolepics Ky.). Review de Faculte des Sciences de l'Universite d'Istanbul. Series
 B: Sciences des Naturale. 17:77-94.
- Benson, Margaret. 1894. Contributions to the embryology of the amentiferae. Part I. Transactions of the Linnean Society of London. 2nd Series-Botany. 3:409-424.
- Bernier, G. 1988. The control of floral evocation and morphogenesis. Annual Review of Plant Physiology and Plant Molecular Biology. 39:175-219.
- Bianco, Pasqua. 1961. Ricerche sul cicle riproduttivo di specie de genere Quercus della flora Italiana. <u>VI</u>. Contributo alla biologia di Quercus trojana webb in Puglia. Annali della Accademia di Scienze Forestali. 10:59-96 (+ plates).
- Bolstad, S.B.; Kang, H.; Cecich, R.A. [in press]. Promoting and selecting for early flowering in jack pine in a controlled environment. In: DeHayes, D., ed. Proceedings of the First Northern Forest Genetics Association Conference; 1991 July 22-24; Burlington, VT.
- Bonnet-Masimbert, M. 1978. Biologie Florale des chenes pedoncules et sessiles (*Quercus pedunculata* Ehrb. et *Q. sessiliflora* Sal.) In Symposium. sur la Regeneration et le Traitement des Forets Feuillues de Qualite et zone Temperee. Nancy, France: IUFRO. 17-29 pp.
- Bonnet-Masimbert, M. 1984. Biologie florale et cycle de reproduction de quelques arbes forestiers. Douglas, pin sylvestre, chene. In P. Pesson et J. Louveaux, eds. Pollinisation et productions vegetales. Paris, France: INRA. 219-243 pp.
- Bowman, John L.; Smyth, David R.; Meyerowitz, Elliot M. 1989. Genes directing flower development in Arabidopsis. The Plant Cell. 1(1):37-52.
- Brown, Roy C.; Mogensen, H. Lloyd. 1972. Late ovule and early embryo development in *Quercus gambelii*. American Journal of Botany. 59(3):311-316.
- Cecich, R.A. 1991. Seed production in oak. In: Laursen, S.; Philips, S.L., eds. The oak resource in the upper midwest - Implications for management; 1991 June 3-6. Winona, MN: University of Minnesota: 125-131.

- Cecich, R.A.; Brown, G.L.; Piotter, B.K. 1991. Pistillate flower abortion in three species of oak.
 (Abstract). In: McCormick, L., ed. Eighth Central Hardwood Forest Conference. 1991 March 3-6. University Park, PA: Pennsylvania State University. 578 pp.
- Conrad, Abram H. 1900. A contribution to the life history of Quercus. Botanical Gazette. 29(2):408-418 (+ plates).
- Corti, Roberto. 1959. Ricerche sul ciclo riproduttivo di specie del genere Quercus della flora Italiana. <u>IV</u>. Osservazioni sulla embriologia e sul ciclo riproduttivo in Quercus ilex L. Annali della Accademia di Scienze Forestali. 8:19-42 (+ plates).
- Davis, Gwenda L. 1966. Systematic embryology of the angiosperms. New York: John Wiley and Sons, Inc. 528 pp.
- Evans, L.T. (ed.). 1969. The induction of flowering. Ithaca, NY: Cornell University Press. 488 pp.
- Farmer, R.E., Jr. 1981. Variation in seed yield of white oak. Forest Science. 27(2):377-380.
- Feret, P.P.; Kreh, R.E.; Merkle, S.A.; Oderwald, R.G. 1982. Flower abundance, premature acorn abscission, and acorn production in *Quercus alba* L. Botanical Gazette. 143(2):216-218.
- Gibson, Lester P. 1964. Biology and life history of acorn infesting weevils of the genus *Conotrachelus* (Coleoptera:Curculionidae). Annals of the Entomological Society of America. 57:521-526.
- Goodrum, P.D.; Reid, V.H.; Boyd, C.E. 1971. Acorn yields, characteristics, and management criteria of oaks for wildlife. Journal of Wildlife Management. 35(3):520-532.
- Grisez, Ted J. 1975. Flowering and seed production in seven hardwood species. Research Report NE-315. U.S. Department of Agriculture, Forest Service. 8 pp.
- Gysel, Leslie W. 1956. Measurement of acorn crops. Forest Science. 2(1):305-313.
- Gysel, Leslie W. 1958. Prediction of acorn crops. Forest Science. 4(3):239-245.
- Hjelmqvist, H. 1953. The embryo sac development of *Quercus robur* L. Phytomorphology. 3(4):377-384.
- Hjelmqvist, H. 1957. Some notes on the endosperm and embryo development in Fagales and related orders. Botaniska Notiser. 110(2):173-195.
- Irgens-Moller, H. 1955. Forest genetics research: Quercus L. Journal of Economic Botany. 9(1):53-71.
- Jicinska, Dagmar; Koncalova, Marie Nadezda. 1978. Flowering and fertilization process in European Sambucus and Quercus species. In: Bonner, F., ed. Proceedings of a symposium on flowering and seed development in trees. Starkville, MS: 103-111.
- Johnsson, H. 1949. Hereditary precocious flowering in *Betula verrucosa* and *B. pubescens*. Hereditas. 35:112-114.
- Jovanovic, Milutin; Tucovic, Aleksandar. 1975. Genetics of common and sessile oak (Quercus robur L. and Q. Petraea Liebl.). Annales Forestales. 7(2):23-53.
- Jovanovic, M.; Tucovic, A.; Vuletic, D. 1971. Comparative analysis of the processes of macrosporogenesis, macrogrametogenesis, and early embryogenesis in the common oak (*Quercus robur L.*) in relation to the type of pollination. Genetika. 3(1):131-145.

- Kearby, William H.; Christisen, Donald M.; Myers, Steven A. 1986. Insects: their biology and impact on acorn crops in Missouri's upland forests. Missouri Conservation Department Terrestrial Series #16. 47 pp.
- Kolpak, M.X.; Feret, P.P.; Stipes, R.J. 1980. Fungi associated with the pistillate flowers of white oak and their effect on pollen germination. Forest Ecology and Management. 3:69-72.
- Kopp, Dennis D.; Yonke, Thomas R. 1973. The treehoppers of Missouri: Part 1. Subfamilies Centrotinae, Hoplohorioninae, and Membracinae (Homoptera: Membracidae). Journal of the Kansas Entomological Society. 46(1):42-64.
- Kopp, Dennis D.; Yonke, Thomas R. 1973. The treehoppers of Missouri: Part 2. Subfamily Smiliinae; Tribes Acutalini, Ceresini, and Polyglyptini (Homoptera: Membracidae). Journal of the Kansas Entomological Society. 46(2):233-276.
- Kopp, Dennis D.; Yonke, Thomas R. 1973. The treehoppers of Missouri: Part 3. Subfamily Smiliinae; Tribe Smiliini. Journal of the Kansas Entomological Society. 46(3):375-421.
- Kopp, Dennis D.; Yonke, Thomas R. 1974. The treehoppers of Missouri: Part 4. Subfamily Smiliinae; Tribe Telamonini (Homoptera: Membracidae). Journal of the Kansas Entomological Society. 47(1):80-130.
- Kossuth, Susan V. 1974. Premature acorn abscission in white oak (*Quercus alba L.*). New Haven, CT: Yale University. 127 pp. Dissertation.
- Lambeth, Clem C. 1980. Juvenile mature correlations in Pinaceae and implications for early selection. Forest Science. 26(4):571-580.
- Langdon, LaDema Mary. 1939. Ontogenetic and anatomical studies of the flower and fruit of the Fagaceae and Juglandaceae. Botanical Gazette. 101:301-327.
- Ledig, F. Thomas; Beland, James W. <u>IV</u>; Fryer, John H. 1971. Breeding techniques for white oak. In: Proceedings of The Eleventh Conference on Southern Forest Tree Improvement. 275-283 pp.
- Merkle, S.A.; Feret, P.P.; Croxdale, J.G.; Sharik, T.L. 1980. Development of floral primordia in white oak. Forest Science. 26(2):238-250.
- Minina, E.G. 1954. Biological bases of flowering and fruit-bearing in oak. Trudy Instituta Lesa Akademiya, Nauk, SSSR. 17:5-97. [Translated by Indian National Scientific Documentation Centre. Delhi, India.]
- Mogensen, H. Lloyd. 1965. A contribution to the anatomical development of the acorn in *Quercus* L. Iowa State Journal of Science. 40(3):221-255.
- Mogensen, H. Lloyd. 1972. Fine structure and composition of the egg apparatus before and after fertilization in *Quercus gambelii*: the functional ovule. American Journal of Botany. 59(9):931-941.
- Mogensen, H. Lloyd. 1973. Some Histochemical, ultrastructural, and nutritional aspects of the ovule of *Quercus gambelii*. American Journal of Botany. 60(1):48-54.
- Mogensen, H. Lloyd. 1975. Ovule abortion in *Quercus* (Fagaceae). American Journal of Botany. 62(2):160-165.
- Okamoto, Motoharu. 1982. Ontogeny of the female inflorescence and flower of Quercus myrsinaefolia. Bulletin of the Osaka Museum of Natural History. 36:21-28.
- Owens, J.N.; Blake, M.D. 1985. Forest tree seed production. Information Report PI-X-53. Canadian Forestry Service, Agriculture, Canada. 161 pp.

- Poethig, R. Scott. 1990. Phase change and the regulation of shoot morphogenesis in plants. Science. 250:923-930.
- Romashov, N.V. 1957. Laws governing fruiting in oak. Botanicheskii Zhurnal. 42(1):41-56. [Translated by Israel Program for Scientific Translation.]
- Sattler, R. 1973. Organogenesis of flowers. Toronto: University of Toronto Press. 207 pp.
- Scaramuzzi, Franca. 1960. Richerche sul ciclo riproduttivo di specie del genere Quercus della flora Italiana. v. osservazioni sul ciclo riproduttivo e sulla embriologia di Quercus aegilops L. Annali della Accademia di Scienze Forestali. 9:289-322.
- Shannon, Susan; Meeks-Wagner, D. Ry. 1991. A mutation in Arabidopsis TFL1 gene affects inflorescence meristem development. The Plant Cell. 3(9):877-892.
- Sharik, T.L.; Ross, M.S.; Hopper, G.M. 1983. Early fruiting in chestnut oak (Quercus prinus L.). Forest Science. 29(2):221-224.
- Sharp, Ward M. 1958. Evaluating mast yields in the oaks. Pennsylvania Agricultural Experiment Station Bulletin 635. 22 pp.
- Sharp, Ward M.; Chisman, Henry H. 1961. Flowering and fruiting in the white oaks. I. Staminate flowering through pollen dispersal. Ecology. 42(2):365-372.
- Sharp, Ward M.; Sprague, Vance G. 1967. Flowering and fruiting in the white oaks. Pistillate flowering, acorn development, weather, and yields. Ecology. 48(2):243-251.
- Short, Henry L. 1976. Composition and squirrel use of acorns of black and white oak groups. Journal of Wildlife Management. 40(3):479-483.
- Singh, A.P.; Mogensen, H. Lloyd. 1975. Fine structure of the zygote and early embryo in *Quercus gambelii*. American Journal of Botany. 62(2):105-115.
- Singh, A.P.; Mogensen, H. Lloyd. 1976. Fine structure of early endosperm in *Quercus gambelii*. Cytologia. 41(1):55-71.
- Smallwood, Peter D.; Peters, Wm. David. 1986. Grey squirrel food preferences: the effects of tannin and fat concentration. Ecology. 67:168-174.
- Smith, Christopher C.; Follmer, David. 1972. Food preferences of squirrels. Ecology. 53(1):82-91.
- Smyth, David R.; Bowman, John L.; Meyerowitz, Elliot M. 1990. Early flower development in *Arabidopsis*. The Plant Cell. 2:755-767.
- Stairs, G.R. 1964. Microsporogenesis and embryogenesis in *Quercus*. Botanical Gazette. 125(2):115-121.
- Teich, A.H.; Holst, M.J. 1969. Genetic control of cone-clusters and precocious flowering in *Pinus sylvestris*. Canadian Journal of Botany. 47(7):1081-1084.
- Turkel, Harold S.; Rebuck, Allen L.; Grove, Alvin R., Jr. 1955. Floral morphology of white oak. Pennsylvania Agricultural Experiment Station Bulletin 593. 14 pp.
- USDA. 1974. Seeds of woody plants in the United States. Agric. Handb. 450. Washington, DC: U.S. Department of Agriculture. 883 pp.
- Vogt, A.R. 1969. Reproduction of the oak. Ohio Report. 54(2):19-21.

- Wareing, P.F. 1959. Problems of juvenility and flowering in trees. Journal of the Linnean Society of London. Series B. 56(366):282-289.
- Williamson, Malcom J. 1966. Premature abscissions and white oak acorn crops. Forest Science. 12(1):19-21.
- Wolgast, Leonard Joseph. 1972. Mast production in scrub oak (*Quercus ilicifolia*) on the coastal plain in New Jersey. New Brunswick, NJ: Rutgers University. 137 pp. Dissertation.
- Wolgast, Leonard J.; Stout, Benjamin B. 1977. Effects of age, stand density, and fertilizer application on bear oak reproduction. Journal of Wildlife Management. 41(4):685-691.
- Wolgast, Leonard J.; Trout, J.R. 1979. Late spring frost affects yields of bear oak acorns. Journal of Wildlife Management. 43(2):239-240.
- Wright, Jonathan W. 1953. Notes on flowering and fruiting of northeastern trees. Station Paper No. 60. U.S. Department of Agriculture, Forest Service, Northeast Forest Experiment Station. 38 pp.
- Zimmerman, Richard H. 1972. Juvenility and flowering in woody plants: a review. Hortscience. 7(5):447-455.

Acorns and Oak Regeneration

D. E. Beck, Southeastern Forest Experiment Station, USDA Forest Service, Asheville, NC 28806

ABSTRACT Acorn production is extremely variable: among species, trees, locations, and years. From the time of flower initiation through seed maturation, a host of intrinsic and extrinsic factors, including weather, insects, disease, and flower and seed predators, combine to eliminate almost all acorns in most years. Only from bumper crops, which on the average for a given species occur once out of every 4 or 5 years, are there any acorns left over to produce seedlings. In specific locations, such complete failures may extend over periods of 10 years or longer.

> Those acorns that do mature and escape predation are far from sure to produce seedlings. Viability deteriorates quickly due to drying if acorns are not covered by soil or leaf litter. The proportion of sound seed that escapes predation and chances to find conditions favorable for germination and seedling survival is very small in all but truly bumper years.

> Despite the seemingly insurmountable odds, oaks have managed to keep their dominant position in the eastern hardwood forests through the buildup of advance reproduction in the form of seedlings and seedling-sprouts that result from these sporadic bumper crops of acorns. However, from a management standpoint, advance reproduction is not always in place when and where we want it, and our inability to ensure acorn production or even to predict it can severely limit our ability to regenerate a specific stand. Sporadic and unpredictable acorn crops also become major problems for planting and direct seeding due to the limited storage life of acorns.

> Given our current state of knowledge, management of acorn crops is limited to ensuring that the *potential* for acorn production is in place. The most consistent finding in studies of acorn production is that only a minority of oak trees are inherently good producers of acorns. Therefore, at present, acorn crop management is largely a matter of identifying the good producers and maintaining them in a dominant position in the stand.

INTRODUCTION There is little question that following natural disturbances and commercial harvest by a variety of means, many of the new stands will contain less oak than the previous ones. On the better quality sites the decrease in oak composition can be dramatic. The problem is generally attributed to a lack of the well-established

advance reproduction that can compete with other vegetation when released. My charge in this paper is to consider: (1) How much, if any, of the oak regeneration problem is due to lack of acorns? (2) If acorns are a problem, is there anything that can be done about it from a management standpoint?

THE PROBLEM Natural Regeneration Natural Regeneration Very few studies have dealt directly with the relationship between acorn production and seedling establishment. In one such study Marquis and others (1976) reported that in an area with abundant advance reproduction (5,250 seedlings per acre) the number of viable seed trapped was about five times greater than in an area with poor advance reproduction (less than 100 seedlings per acre). They also found losses to rodents to be very high in the low-reproduction area and relatively low in the high-reproduction area. After depredations, almost 60 times as many seed could be expected to germinate in the area with good advance reproduction as in the low-regeneration area.

> In other areas with poor reproduction the evidence pointing to lack of acorns as a factor is more speculative. Most of the information on the subject is observational and anecdotal based on the very erratic and uncertain nature of acorn production. Acorn crops are extremely variable from tree to tree, from location to location, and from year to year (Beck and Olson 1968, Beck 1977, Christisen 1955, Christisen and Korschgen 1955, Downs and McQuilken 1944, Gysel 1956, Minckler and Janes 1965, Tryon and Carvell 1962a, 1962b). In a given geographic area, acorn crops may vary year-to-year from nearly zero to over 250,000 acorns per acre. Good acorn crops may occur at long intervals-on average at intervals of 4 or 5 years. For specific stands more than 10 years may pass without good crops (Beck 1977). In the Mississippi Delta, Johnson (1979) found 100,000 Nutall oak seedlings per acre established in a good seed year. However, over the next 15 years seedlings were established in only 2 years and then in only small quantities. This variability is related to a broad array of poorly understood factors. But, sporadic acorn production-coupled with loss to predators, short storage life, seedling establishment problems, and low survival of established seedlings-no doubt contributes to scarcity of reproduction.

- Artificial Regeneration If artificial regeneration programs demand acorns and seedlings on a large scale and on a continuous, predictable basis, then sporadic seed production will become a major problem. Given the short storage life of acorns, long intervals with very low or nonexistent seed production will be highly disruptive to major planting and seeding programs.
 - **THE PROCESS** The acorn production process—from initiation and development of flowers, through pollination, fertilization, and acorn development to germination and seedling establishment—is affected by a host of biotic and abiotic factors (figure 1). Acorns develop from fertilized flowers with all species of oak producing both male (staminate) and female (pistillate) flowers on the same tree. The white oak group (Lepidobalanus subgenus) requires 1 year for the maturation of fruit while the red oaks (Erythrobalanus subgenus) require 2 years. At each stage there are multiple factors, both intrinsic and extrinsic, which affect the process and determine whether

or not mature, viable acorns will be available for regeneration. Are there critical points in the process where losses occur that are amenable to management?

The early ontogony of acorn initiation has been extensively studied and is well understood (Conrad 1900, Hartig 1851, Hjelmqvist 1953, Langdon 1939, Merkle and others 1980, Mogensen 1975, Stairs 1964, Turkel and others 1955). The most complete information for any one species on the reproductive cycle exists for whiteoak (Q. *alba* L.) (Feret and others 1982, Merkle and others 1980, Mogensen 1965, Sharp and Chisman 1961, Sharp and Sprague 1967, Stairs 1964, Turkel and others 1955). Even here there are many gaps and much disagreement as to what may or may not be the causes of the huge variability in acorn crops among trees, places, and years. And for most of the other species, the reproductive biology is very poorly understood.




Figure 1-Acorn production-process and impacts.

Flower Formation There is a great deal of equivocation as to whether or not flowering, or, more properly the lack of it, limits acorn production. Romashov (1957) and Sharp and Sprague (1967) stated that year-to-year acorn crop variability was not a function of variability in flower production as flowers were observed annually. Olson and Boyce (1971), in a summary of current knowledge on acorn production, concluded that flowers are formed every year in great abundance by practically every tree of flowering size, and age, and therefore, year-to-year variability in size of acorn crops is not due to lack of flowering capability, nor to cyclic formation of flowers. On the other hand, Feret and others (1982) found acorn production was significantly related to spring flower abundance for white oak in the Virginia Piedmont. Farmer (1981) and Williamson (1966) have reported great differences in flower production from year to year for white oak. The evidence is not conclusive that failure to either induce or differentiate flowers is a cause of "off" year production, but it remains a real possibility.

Page 161 of 404

Anthesis, Pollination, Fertilization Whether or not flowers are consistently developed, it is certain that the oak species frequently produce mature fruits from only a small fraction of their female flowers (Cecich 1991, 1992; Feret and others 1982; Kossuth 1974; Williamson 1966). It is not unusual for 75 to nearly 100 percent of flowers to abscise prematurely; the heaviest losses occurring early in the process prior to fertilization—the period that covers pollination and ovule development.

Numerous climatic factors have been implicated in pollen dispersal, emergence of pistillate flowers, receptivity, etc. Temperature, relative humidity, precipitation, wind, and late freezes have all been implicated at various times and localities (Goodrum and others 1971, Sharp 1958, Sharp and Chisman 1961, Sharp and Sprague 1967, Wolgast and Trout 1979). Wolgast (1972) demonstrated experimentally that high relative humidity at the time of flowering is detrimental to fruit set in scrub oak. Other conclusions about the effects of weather are based on observation and simultaneous occurrence of events, and not cause and effect relationships. Thus, the influence of weather variables on setting of fruit and subsequent acorn yield is mostly inconclusive. Cecich (1991,1992) has hypothesized that a complex of sucking insects (tree hoppers, Membracidae) may play an important role in pistillate flower abortion, but the theory is yet to be proved.

Stephensen (1981) did an exhaustive synthesis of the literature on the proximate factors that limit fruit and seed production between anthesis and dispersal for a large number of taxonomically and ecologically diverse species that regularly abscise a large portion of their flowers and immature fruits. He concluded that, in some cases, lack of pollination, fruit and seed predation, or adverse weather may hold fruit production below the upper limit. But in most cases, resources limit fruit production. Pollinated flowers and juvenile fruits abscise until fruit and seed number match the available resources. The level of resources available for fruit development in a given reproductive episode is influenced by a host of intrinsic as well as extrinsic factors. The causes of premature abscission may well occur prior to and not in conjunction with the event.

- **Developing Acorns** Reasons for the premature drop of developing acorns is largely unknown. However, in a Missouri study the primary moth invader, *M. latiferreanus*, and cynipid gall wasps infested 17 percent of immature acorns collected in a 4-year study (Kearby and others 1986).
 - Mature Acorns Mature acorns incur even heavier losses from insects. Weevils of the genus *Curculio* are the major culprit (Barrett 1931, Beck 1977, Beck and Olson 1968, Collins 1961, Downs and McQuilken 1944, Kearby and others 1986). However, *Conotrachelus* weevils, cynipid gall wasps, and both primary and secondary Lepidoptera have all been implicated (Kearby and others 1986). Infestation rates vary considerably from year to year, with nearly 100 percent in some years. In several long-term studies, infestation by insects averaged around 50 percent—a very significant impact (Beck 1977, Burns and others 1954, Christisen 1955, Downs and McQuilken 1944, Kearby and others 1986).

Mature acorns that escape destruction by insects face yet another hurdle. Nearly 200 species of forest wildlife consume acorns. Some, such as the darge og ame

animals, consume large quantities as individuals and in aggregate. But the sciurids (chipmunks, squirrels, etc.) probably have the largest impact (Johnson and others 1989). In looking at cumulative mast needs by forest wildlife in the Southern Appalachians, they said that acorn crops of <200 lbs/acre usually are totally consumed. And, over a period of years the demand will always exceed the supply. In a 12-year study in the Southern Appalachians at one location, production of sound acorns exceeded 200 lbs/acre only four times (Beck 1977). Average production was 186 lbs/acre and that number was influenced heavily by one bumper year with 800 lbs produced. Even in poor mast years some acorns escape predators and produce seedlings. But it takes a bumper-crop year for any appreciable number to do so.

Even those acorns which mature and escape destruction by insects and animals are far from certain to produce seedlings. Sound, undamaged acorns have a germinative capacity between 75 and 95 percent (USDA 1974). But for germination to occur, the moisture content of acorns must not drop below 30 to 50 percent for white oaks and 20 to 30 percent for red oaks (Korstian 1927). Those acorns not covered by litter or soil are very susceptible to drying. Therefore, a fairly good percentage of those acorns which mature and escape predation probably fail to produce seedlings.

- **POTENTIAL IMPACTS** Taking an overall view, it is evident that numerous intrinsic and extrinsic factors influence acorn production throughout the process from initiation of flowers to germination of the acorns. There is no point at which we can identify a particular factor(s) and say this factor is the cause of the problem. It is quite conceivable that a combination of factors may reduce acorn yield at a particular time or place; or at others, a single factor could be limiting, but not always the same one. Stephensen (1981) suggests, for example, that premature abscission of flowers could be due to different causes on different branches of the same tree. There is no simple or easy explanation, or fix, for the problem of sporadic, unpredictable acorn crops.
 - **Extrinsic Factors** Various weather factors can adversely affect production. Some, such as unfavorable relative humidity at time of pollen dispersal, can limit production. Others, such as late freezes, may completely wipe out a crop at a given location. In either case there isn't much we can do about it. We do need to understand the effect of climatic factors, if for no other reason than to sort out some of the "noise" so we can better understand other factors which limit acorn production and may be more amenable to management. Understanding weather effects may also help in short-term prediction of acorn crops.

The possibility for control of the other major extrinsic factor with direct impact—insects—has been demonstrated (Dorsey and others 1962, Dorsey 1967). They were able to reduce weevil damage to white and red oak acorns using systemic insecticides. However, control of insects in a forest setting is probably not economically feasible. In seed production areas or seed orchards, experience with the pine species suggests we can control damage if there is economic demand for seed. But we need a great deal more information than we have now. Highly effective control of insect damage is unlikely, however, to eliminate poor acorn crops. But insect control does have the potential to raise the average level.

Intrinsic Factors A great deal of attention has been focused on the extrinsic factors—mainly weather, insects, and wildlife—which directly limit acorn production. In the long run, however, whether or not acorns are produced probably depends more on intrinsic factors which determine physiological condition of the tree that allows it to initiate and develop flowers, set fruit, and supply the resources to carry the fruit to maturity.

Age/Size. One of the more obvious factors is tree age and/or size. There are records of acorn production from very young trees of sprout origin (Wood 1934) and from a few short-lived species such as scrub oak (Wolgast 1972, Smith 1929, Little and others 1958). However, most species do not begin production until the trees are around 20 to 25 years old (USDA 1974, Reid and Goodrum 1957). They also suggested that crowded trees appear to begin production later than trees with more growing space. Some workers suggest that substantial yields of acorns cannot be expected earlier than age 40 (Goodrum and others 1971). Yields apparently increase with age and or at least with size up to a point, level off for an extended period, and then decrease (Downs and McQuilken 1944). But the yield-age relationship is anything but clear.

Crown Position. Crown size and dominance has been shown to be a major determinant of acorn production (Carvell and Korstian 1955, Christisen and Korschgen 1955, Matthews 1963, Reid and Goodrum 1957). Drake (1991) found that only 9 percent of suppressed trees and 38 percent of intermediates produced seed, and those that did, produced only small quantities. Although there was much variation among trees in terms of quantity, 75 percent of codominants and 96 percent of dominants produced at least some seed.

Inherent Capability. The most consistent finding in acorn production studies across all species and under the most diverse environmental conditions is that some trees are consistently good producers and others consistently poor producers, and the differences are not related to obvious characteristics of the trees (Burns and others 1954, Christisen 1955, Christisen and Korschgen 1955, Collins 1961, Cypert 1951, Cypert and Webster 1948, Downs 1949, Downs and McQuilken 1944, Feret and others 1982, Goodrum and others 1971, Reid and Goodrum 1957, Sharp 1958, Smith 1929, Tryon and Carvell 1962, Wood 1934). These production tendencies are most often attributed to inherent capabilities. Wolgast (1972, 1977) lends support to this thesis with his studies of scrub oak. Good, medium, and poor producers were transplanted to a common site but retained their production tendencies. Farmer (1981) found strong clonal control over acorn yield in a grafted white oak seed orchard. In a good seed year clonal differences accounted for over 50 of the variance in percentage of shoots flowering, number of flowers and inflorescences per shoot, percentage of flowers maturing to acorns, and acorn yield.

MANAGEMENT IMPLICATIONS Of the major extrinsic factors (weather, insects, and wildlife) which limit acorns available for regeneration, none is very amenable to management in the general forest setting. Weather variables have the potential to reduce or completely eliminate an acorn crop in a given time and place. And we need a better grasp of the impacts of weather variables if we expect to understand the overall process. But we are not likely to exert any direct control in the near future under any circumstances.

Insects are potentially very damaging to the acorn crop at many stages of the production process. Numerous insects feed on foliage of oaks. Their impact on acorn production has not been assessed. But such insects as aphids have been shown to have large impacts on nut production in species such as pecans by depleting available energy. It has been hypothesized that premature abscission of flowers may be related to leaf hoppers. It has been observed repeatedly that weevils destroy 50 percent of mature acorns on an average. Experience with seed orchards and seed production areas for the southern pines demonstrates the possibilities for control of insects when an economic demand exists. However, the technical base does not exist for the oak species even if the demand were there.

It is technically possible to reduce animal depredations by reducing or excluding animal populations. But in a forest setting it is probably not ecologically desirable or politically possible to reduce populations. In fact mast production specifically for wildlife may often be a management objective. It is probably possible to reduce animal depredations in seed orchards; but even there, reducing losses to wildlife will not be a major way to increase acorns available for regeneration.

At the present time the best chance for acorn production enhancement is in applying what we know about tree-to-tree variation. We should recognize and manage those trees capable of producing large acorn crops on a consistent basis. The large inherent differences among trees in productive capability have been observed and reported numerous times over the past 50 years. But this knowledge has not been applied due to the difficulty in identifying acorn producers. Given the many other factors that may impact production in a given year, identification of consistent producers is not always easy to do. Maintaining the good acorn producers in the stand in a dominant position is currently the only practical way to enhance acorn production.

LITERATURE CITED	Beck, Donald E.; Olson, David F., Jr. 1968. Seed Production in Southern Appalachian oak stands.
	Research Note SE-91. Asheville, NC: U.S. Department of Agriculture, Forest Service,
	Southeastern Forest Experiment Station. 7 pp.

- Beck, Donald E. 1977. Twelve-years acorn yield in southern Appalachian oaks. Research Note SE-244. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 8 pp.
- Burns, Paul Y.; Christisen, Donald M. 1954. Acorn production in the Missouri Ozarks. Research Note Bulletin 611. Agricultural Experiment Station, University of Missouri, College of Agriculture.
- Cecich, Robert A. 1991. Seed production in oak. In: Laursen, Steven B.; DeBoe, Joyce F. Conference proceedings; The oak resource in the Upper Midwest implications for management; 1991 June 3-6; publication number NR-BU-5663-5. St. Paul, MN: University of Minnesota: 125-131.
- Cecich, Robert A. 1992. Flowering and Oak Regeneration. In: Christisen, D.M. 1955. Yield of seed by oaks in the Missouri Ozarks. Journal of Forestry. 53:439-441.
- Christisen, D.M.; Korschgen, L.J. 1955. Acorn Yields and Wildlife Usage in Missouri. Trans. 20th North American Wildlife Conference: 337-357.

- Collins, J.O. 1961. Ten-year acorn mast production study in Louisiana. Louisiana Wildlife Fish Commission; P-R Rep., Proj. W-29-R-8. 33 pp.
- Conrad, A.H. 1900. A contribution to the life history of Quercus. Botanical Gazette. 29:408-418.
- Cypert, Eugene. 1951. Suggestions for the management of oak forests for mast production. Mimeo. paper presented at Southeastern Association Game and Fish Commissioners' Conference, Edgewater Park, Miss.
- Cypert, Eugene; Webster, B.S. 1948. Yield and use by wildlife of acorns of water and willow oaks. Journal of Wildlife Management. 12:227-231.
- Dorsey, C.K.; Tryon, E.H.; Carvell, K.L. 1962. Insect damage to acorns in West Virginia and control studies using granular systemic insecticides. Journal of Economic Entomology. 55:885-888.
- Dorsey, C.K. 1967. Experiment to control acorn weevils with systemic insecticides. Forest Science. 13:390-396.
- Downs, Albert A.; McQuilkin, William E. 1944. Seed production of southern Appalachian oaks. Journal of Forestry. 42:913-920.
- Drake, William E. 1991. Evaluation of an approach to improve acorn production during thinning. In: McCormack, Larry H.; Gottschalk, Kurt W., eds. Proceedings, 8th central hardwood forest conference; 1991 March 4-6; Gen. Tech. Rep. NE-148. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.
- Farmer, R.E., Jr. 1981. Variation in seed yield of white oak. Forest Science. 27(2):377-380.
- Feret, P.P.; Kreh, R.E.; Merkle, S.A.; Oderwald, R.G. 1982. Flower abundance, premature acorn abscission, and acorn production in Quercus alba L. Botanical Gazette. 143(2):216-218.
- Goodrum, P.H.; Reid, V.H.; Boyd, C.E. 1971. Acorn yields, characteristics, and management criteria of oaks for wildlife. Journal of Wildlife Management. 35:520-532.
- Hjelmqvist, H. 1953. The embryo sac development of *Quercus robur* L. Phytomorphology 3(4):377-384.
- Kearby, William H.; Christisen, Donald M.; Myers, Steven A. 1986. Insects: Their Biology and Impact on Acorn Crops in Missouri's Upland Forests. Terrestrial Series #16. Jefferson City, Missouri: Missouri Department of Conversation. 48 pp.
- Korstian, Clarence F. 1927. Factors Controlling Germination and Early Survival in Oaks. Yale University School of Forestry Bulletin 19. 115 pp.
- Kossuth, S.V. 1974. Premature acorn abscission in white oak (Quercus alba L.). Yale University. 127 pp. Dissertation.
- Langdon, L.M. 1939. Ontogenetic and anatomical studies of the flowers and fruit of Fagacea and Juglandaceae. Botanical Gazette. 101:301-327.
- Marquis, David A.; Eckert, Philip L.; Roach, Benjamin A. 1976. Acorn weevils, rodents, and deer all contribute to oak-regeneration difficulties in Pennsylvania. Research Paper NE-356. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 5 pp.
- Matthews, J.D. 1963. Factors affecting the production of seed by forest trees. Forestry Abstracts. 24:1-13.
- Merkle, S.A.; Feret, P.P.; Croxdale, J.G.; Sharik, T.L. 1980. Development of floral primordia in white oak. Forest Science. 26:238-250.

- Mogensen, H.L. 1965. A contribution to the anatomical development of the acorn in Quercus L. Iowa State Journal of Science. 40:221-255.
- Mogensen, H.L. 1975. A contribution to the anatomical development of the acorn in Quercus L. Iowa State Journal of Science. 40:221-225.
- Olson, David F.; Boyce, Stephen G. 1971. Factors affecting acorn production and germination and early growth of seedling and seedling sprouts. In: Oak symposium proceedings; [Date of conference unknown]; [Location of conference unknown]. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 44-48.
- Reid, Vincent H.; Goodrum, Phil D. 1957. Factors influencing the yield and wildlife use of acorns. Proc. 6th Annual For. Symposium, Louisiana State University: 46-79.
- Romashov, N. V. 1957. Laws governing fruiting in oak. Botanicheskii Zhurnal. 42(1):41-56.
- Sharp, Ward M.; Sprague, Vance G. 1967. Flowering and fruiting in the white oaks. Postillate flowers acorn development, weather, and yields. Ecology. 48(2):243-251.
- Sharp, Ward M.; Chisman, Henry H. 1961. Flowering and fruiting in the white oaks. I. Staminate flowering through pollen dispersal. Ecology. 42(2):365-372.
- Sharp, W.M. 1958. Evaluating mast yields in the oaks. Bulletin 635 University Park, PA: Pennsylvania State University, Agriculture Experiment Station. 22 pp.
- Smith, J. R. 1929. Tree Crops: A Permanent Agriculture. New York: Harcourt, Brace and Co. 333 pp.
- Stairs, G.R. 1964. Microsporogenesis and embryogenesis in Quercus. Botanical Gazette. 125:115-121.
- Stephenson, A.G. 1981. Flower and fruit abortion: proximate causes and ultimate functions. Ann. Rev. Ecol. Syst. 12:253-279.
- Tyron, E.H.; Carvell, K.L. 1962. Acorn production and damage. Bulletin 466T. West Virginia University, Agricultural Experiment Station, College of Agriculture, Forestry and Home Economics. 18 pp.
- Turkel, H.S.; Rebuck, A.L.; Grove, A.R., Jr. 1955. Floral morphology of white oak. Pennsylvania Agriculture Experiment Station Bulletin 593. 15 pp.
- USDA. 1974. Seeds of woody plants in the United States. Agriculture Handbook 450. Washington, DC: U.S. Department of Agriculture. 883 pp.
- Williamson, M.J. 1966. Premature abscissions in white oak acorn crops. Forest Science. 12:19-21.
- Wolgast, Leonard J. 1972. Mast production in scrub oak (Quercus Ilicifolia) on the coastal plain in New Jersey. Thesis.
- Wolgast, Leonard J.; Trout, J.R. 1979. Late spring frost affects yields of bear oak acorns. Journal of Wildlife Management. 43(2):239-240.
- Wood, O.M. 1934. A Brief Record of Seed Productivity for Chestnut Oak in Southern New Jersey. Journal of Forestry. 32:1014-1016.

Insects and Diseases Affecting Oak Regeneration Success

Steven W. Oak, Forest Pest Management, Southern Region, USDA Forest Service, Asheville, NC 28802

ABSTRACT Flowers, acorns, seedlings, sprouts, and mature oaks are damaged and killed by myriad insects and diseases, but relatively few have been shown to prevent successful oak regeneration. The best known are acorn-infesting insects (*Curculio* weevils, the filbertworm, and gall wasps) and root diseases of nursery-grown seedlings caused by *Phytophthora* spp. and *Cylindrocladium* spp. Preventing insect infestation of mature acorns is not yet possible but hot water treatments are used to kill weevil larvae already inside acorns. Management of nursery root diseases is accomplished through the integration of sanitation, soil management practices, and soil fumigation. Emerging pest management issues in oak regeneration include flower pest identification and assessment, control technologies for seed production areas, alternatives to methyl bromide-chloropicrin soil fumigants in nurseries, and the effects of oak decline on natural regeneration opportunities in upland oak forests.

The wide distribution and diversity of oaks in the U.S. are matched by the vast array of diseases and insects affecting them. However, relatively few pests have been conclusively shown to significantly affect the ability to regenerate oak. These are acorn-infesting insects and nursery root diseases. This paper focuses on the most damaging pests in these two groups and presents proven options for controlling them. Other pest management issues are discussed that may emerge with increased emphasis on artificial regeneration and with large-scale ecological impacts resulting from the interactions of oak decline and insect defoliation in the Southeast.

ACORN-INFESTING INSECTS Acorn-infesting insects are the most studied group of pests affecting oak regeneration. There are several types of insects involved but the most notorious among them are several *Curculio* weevil species. Virtually all oak species are attacked by one or more of 22 different weevils recorded by Williams (1989).

Adult weevils are themselves not damaging. However, larvae hatching from eggs laid in late summer in tiny niches beneath the shell may consume most of the nut tissue within a few weeks. When larvae mature, they bore out through the shell and migrate underground to pupate. The rate of infestation is variable but has exceeded 90 percent in some northern red oak collections (Gibson 1982). Single acorns are commonly hosts to three or more larvae; embryos in infested acorns that escape damage may germinate but seedlings grow slower than those from uninfested acorns (Oliver and Chapin 1984).

Infestation rates of the filbertworm (*Melissopus latiferreanus*) are much lower than for acorn weevils (Gibson 1982), but they have been responsible for large losses, particularly in low production years (Drooz 1985). Damage is caused by larval feeding and is usually lethal to infested acorns. Adult moths lay single eggs on leaf surfaces near acorn clusters in mid-summer. After hatching, the larva crawls to an acorn, bores through the shell, and begins feeding. The mature larva exits the acorn after 3 to 5 weeks, spins a cocoon in the top few inches of soil, and pupates in winter.

Non-stinging gall wasps are another group of primary acorn pests. Like *Curculio* spp. and the filbertworm, they infest and can kill intact acorns. There are many species that form unusual galls on various oak tissues, but *Callyrhytis operator* and *C. fructuosa* are the most common in acorns. Gibson (1982) found them to be less ubiquitous than either weevils or filbertworms in northern red oak acorn collections. However, the infestation rate of *C. fructuosa* was second only to weevils in two collections made in consecutive years from a Tennessee seed orchard (L. Barber and author, unpublished data¹). Acorns infested by *C. fructuosa* appear normal on the outside but are filled with up to 2 dozen larvae encased in small stony galls. *C. operator*, on the other hand, forms a gall in the side of the acorn shell. Gall wasps have complicated life cycles, and the same species may induce galls on different plant parts at different times of the year.

Some insects and pathogenic fungi and bacteria can invade acorns damaged by other agents and thereby increase losses. The best known insects with this mode of action are *Conotrachelus* weevils and the acorn moth, *Valentinia glandulella*. Neither can breech intact acorns, but the acorn moth has been known to attack otherwise healthy, germinating acorns (Galford 1986). Examples of pathogens following acorn damage are a bacterium, *Erwinia quercina*, and a fungus, *Fusarium solani*. *E. quercina* causes a disease in California live oak (*Q. agrifolia*), called drippy nut, after it gains access to acorns through oviposition punctures of gall wasps (Hildebrand and Schroth 1967). The name of the disease describes the main symptom, which is the byproduct of anaerobic fermentation and results in acorn rot and premature abscission. *F. solani* was isolated from weeviled acorns collected in Mississippi (Vozzo 1983) and Tennessee (author, unpublished data²). The fungus can cause damping-off and root disease in seedlings, but it's impact in these cases is unknown since saprophytic forms exist and pathogenicity was not confirmed experimentally.

Control of Acorn-Infesting Insects

Hot water treatment is the best proven method for controlling weevil larvae in infested acorns. The key elements are water temperature and duration of treatment, but critical thresholds apparently vary with tree species. Crocker and Morgan (1983) prescribed 30-45 minutes at 43 °C followed by quenching in cold water for live oak acorns without causing undue loss of viability. At the Fusiform Rust Resistance Screening Center in Asheville, N.C., northern red oak acorns are treated for 40 minutes at 49 °C \pm 1 °C and air dried. Floating acorns are discarded and the rest stored in plastic bags after dusting with fungicide (Knighten and others 1988).

^{1,2} Data on file, Southern Region, Forest Pest Management, Asheville, N.C.

Germination after this treatment has been estimated between 80 and 90 percent after 6 months cold storage and 20 percent after 4.5 years (J. Knighten, personal communication). Other acorn-infesting insects are presumed to be killed along with weevil larvae. Research is needed to identify the critical parameters for acorns of different species and to determine if and when heat treatment is necessary. Bonner and Vozzo (1987) and Crocker and Morgan (1983) have suggested that, for some seedlots, the benefits of heat treatment in weevil larvae control might be outweighed by detrimental effects on germination and early growth.

Treatments for preventing infestation do not exist. Some systemic insecticide treatments reduced weeviling and increased the percentage of sound acorns in northern red oak and white oak (Dorsey and others 1962, Dorsey 1967) without depressing germination (Tryon and others 1968). However, all of the tested chemicals have high mammalian toxicity (LD_{so} less than 15 mg/kg). One of the successful chemicals (Bidrin[•]= dicrotophos formulated as a water soluble injectable) is EPA-registered for use on oak.

NURSERY ROOT DISEASES Tree seedlings grown in large numbers in confined areas under the lush growing conditions typical of nurseries are subject to damage from a whole suite of pests not normally found in forested settings. Defoliating insects, foliage disease, shoot

not normally found in forested settings. Defoliating insects, foliage disease, shoot cankers, and root diseases can cause dramatic symptoms in nurseries. However, the most widespread and damaging of oak nursery pests are root diseases caused by fungi in the genera *Phytophthora* and *Cylindrocladium*. They have tough resting spore stages that can survive very harsh conditions and persist in soil for long periods in the absence of suitable hosts. P. cinnamomi is a virulent pathogen of many oak species in nurseries, most notably northern red oak (Crandell and others 1945). It is a member of a group of fungi known as water molds that are favored by periodically wet soil conditions and have a swimming spore stage that attacks feeder roots. Once established in the root system, the fungus spreads to larger laterals and the tap root. Cylindrocladium spp. have no mobile spore stage but they form densely compacted masses of fungal tissue in soil and colonized plant parts called sclerotia that resist all but the most aggressive of control measures. Several different Cylindrocladium spp. have caused root rot in cherrybark (Smyly and others 1977) and shumard oaks (Affeltranger and Burns 1983) in bareroot nurseries and in northern red oak grown in containers (Oak and Triplett 1985).

Root-diseased seedlings may display a range of above-ground symptoms including pre- and post-emergence damping-off, stunting, top dieback, foliage yellowing, and premature defoliation. Below ground, infected root systems are sparse, stunted, and usually blackened. Obviously diseased seedlings are easily recognized and culled, but those with less distinct symptoms may be outplanted and die later or grow slower than healthy seedlings.

Control of Nursery Root Diseases Nursery root disease losses have been most effectively avoided where sanitation, soil management, and chemical measures have been combined into an integrated pest management system (Cordell and others 1989). Sanitation starts with excluding the pathogens that can be transported to uninfested areas in contaminated soil by thoroughly cleaning equipment that has been used in infested areas. If

pathogens become established, then a vigilant inspection schedule, culling of affected stock, and in extreme cases, guarantine, can reduce losses. Soil management practices that have an important influence on root disease probability include crop rotation, choice of cover crops, managing water, and in containerized nurseries, choice of growing medium. Continuous seedling cropping leads to increased populations of pathogens, and some cover crops are also hosts for seedling pathogens. For instance, legume crops increase inoculum levels of Cylindrocladium spp., but grass cover crops do not (Soloman and others 1987). Inadequate drainage or over-watering can simultaneously increase inoculum of *Phytophthora* spp. and increase seedling susceptibility by depressing seedling vigor and inhibiting normal root development. Commercially prepared "artificial" growing media are preferred in container systems over mixtures that include field soil, due to the risk of introducing pathogens and the difficulty in eradicating them once introduced (Oak and Triplett 1985).

Root pathogens are among the most difficult nursery pests to control chemically. Several soil fumigants are EPA-registered but the most effective are formulations containing 67 percent methyl bromide and 33 percent chloropicrin (Cordell and others 1989). Proper soil preparation, soil moisture, temperature, and deep placement of the fumigant are essential for effective treatment. Fungicide drenches applied to the soil (e.g. metalaxyl for *Phytophthora* spp. and benomyl for *Cylindrocladium* spp.) are registered for nursery sites and are widely prescribed, but the research basis for recommending them to control root diseases in oak seedlings is lacking.

OTHER PESTS Other diseases and insects can cause dramatic damage with consequences for oak regeneration success but they are usually limited in time or geographic distribution. Spring oviposition of cicadas (*Magicicada* spp.) results in dieback of shoots that bear flowers and developing acorns. There are many species with life cycles ranging from 4 to 17 years, but damage is most dramatic when overlapping broods of periodical cicadas emerge simultaneously (Borrer and others 1976). Root feeding by nymphs reduced growth but not acorn yield of bear oak in New York (Karban 1985).

The success of natural oak regeneration is usually determined not by pests, but by the competitiveness of seedlings, seedling sprouts, and stump sprouts relative to the propagules of other species that might displace it. However, cankers on stump sprouts caused by *Botryodiplodia gallae* have resulted in repeated regeneration failures in mixed oak stands on poor sites after clearcutting in a National Forest in Michigan (Croghan and Robbins 1986). Cankers caused by *Botryodiplodia* spp., as well as other relatively weak pathogens, are common on twigs and branches of stressed trees in both forested and ornamental settings, but rarely have they been seen as the primary cause of death of trees. Except for avoiding the use of clearcutting, management guidelines do not yet exist. Silvicultural treatments to encourage regeneration from seedlings or seedling sprouts are being explored.

EMERGING PEST MANAGEMENT ISSUES

It seems likely that the use of direct seeding and planting will increase as means for compensating for oak reproduction shortages in many settings. Presently, seed requirements are met by wild collections, but small seed orchards are reaching seed-bearing age. The value of the crop in these special areas is high enough that costly pest management strategies unthinkable in forest settings become feasible. While prevention of acorn weevil infestation is the most urgent pest control priority given current knowledge, there is a large gap in knowledge concerning the role of other pests in seed production. Therefore, the highest pest management priority in seed production areas is pest identification and assessment. Efforts in this area have already begun as a component of research into oak flowering and acorn production biology. Monitoring in Missouri and Tennessee has revealed that mature acorns can represent as little as 3 or 4 percent of the initial flower crop (L. Barber, R. Cecich, unpublished data).³ Treehoppers (Homoptera: Membracidae) have been found killing oak flowers in Missouri during the period when losses are most pronounced (R. Cecich and others 1991), but the relative importance of insects, weather, and other potentially limiting factors is unknown.

Once pests are identified, control technologies will be needed. Parallels in pine seed orchard pest management include new pesticide formulations; new application techniques for existing pesticides; behavioral chemicals for insect population monitoring, insecticide application timing, or mating disruption; and orchard cultural practices. A filbertworm sex pheromone has already been identified and is used in commercial filbert nut orchards for population monitoring and for timing insecticide treatments (Aliniazee 1983).

Soil fumigation is the only management response available when nursery root diseases are found. EPA review of the most effective methyl bromide-chloropicrin formulations underscores the need for alternatives in the event that label registrations are withdrawn. Research is needed in soil population monitoring, the delineation of inoculum levels associated with damage under different nursery conditions, and alternative control measures including biocontrol.

Most references to oak regeneration problems are for high-quality sites, but oak decline has the potential to limit natural regeneration opportunities in mature oak stands growing on drier sites. There is no doubt that large areas are being affected by oak decline. Over 1 million acres of oak forest type were affected in the northern Piedmont and Appalachian Mountains of Virginia in 1986 (Oak and others 1991). Preliminary estimates for the Piedmont and mountains of North Carolina were about .9 million acres in 1990 (R. Sheffield, unpublished data).⁴ The regeneration impacts are less clear, however. Acorn production potential is reduced as a result of crown dieback and tree mortality (Oak and others 1989). Advanced physiological age, root disease, and carbohydrate physiology prevent vigorous sprouting of overstory trees in decline areas. Whether these effects result in an inability to adequately regenerate oak depends on the abundance and competitiveness of oak propagules relative to other species that might displace it. Monitoring of a declining scarlet oak/black oak stand in Missouri over 5 years showed overstory mortality increasing from 36 to 62 percent and shifts in understory composition towards white and post oaks, red maple, and shortleaf pine

³ Data on file, Southern Region, Forest Pest Management, Asheville, N.C., and North Central Forest Experiment Station, Columbia Mo.

⁴ Data on file, Southeastern Forest Experiment Station, Asheville, NC.

(Johnson and Law 1989). Oaks will probably be a smaller component in the next stand and oak species diversity will be less. Similar conclusions were reached after 8 years of monitoring in gypsy moth defoliated areas of the Pocono Mountains in Pennsylvania (Gansner and others 1983), where the greatest oak losses were in

areas suffering repeated, heavy defoliations. Pre-defoliation decline conditions were not reported. On the George Washington National Forest in Virginia, the combined effects of pre-existing oak decline and gypsy moth defoliation have resulted in 22,500 acres of severe mortality accumulated between 1987-1991. Research is needed into the interactions of oak decline, insect defoliation, overstory competition, understory composition, and regeneration timing as well as options for dealing with stands suffering severe mortality from decline and defoliation.

LITERATURE CITED Affeltranger, Charles E.; Burns, Edmond B. 1983. Root rots on shumard oak in a central Louisiana nursery. Tree Planters Notes. 34(3):38-39.

- Aliniazee, M.T. 1983. Monitoring the filbertworm, *Melissopus latiferreanus* (Lepidoptera: Olethreutidae), with sex attractant traps: Effects of trap design and placement on moth catches. Environmental Entomology. 12(1):141-146.
- Bonner, F.T.; Vozzo, J.A. 1987. Seed biology and technology of *Quercus*. Gen. Tech. Rep. SO-66. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 21 pp.
- Borrer, Donald J.; Delong, Dwight M.; Triplehorn, Charles A. 1976. An introduction to the study of insects. 4th ed. Atlanta, GA: Holt, Rinehart and Winston. 852 pp.
- Cecich, R.A.; Brown, G.L.; Piotter, B.K. 1991. Pistillate flower abortion in three species of oak [Abstract]. In: Proceedings of the eighth central hardwoods forest conference; 1991 March 3-6; University Park, PA.
- Cordell, Charles E.; Anderson, Robert L.; Hoffard, William H. [and others], tech. coord. 1989. Forest nursery pests. Agric. Handb. 680. Washington, DC: U.S. Department of Agriculture, Forest Service: 184 pp.
- Crandell, Bowen, S.; Garvatt, G.F.; Ryan, Margaret Milburn. 1945. Root disease of Castaneu species and some coniferous and broadleaf nursery stocks, caused by *Phytophthora cinnamomi*. Phytopathology. 35:162-180.
- Crocker, Robert L.; Morgan, David L. 1983. Control of weevil (*Curculio* sp.) larvae in acorns of live oaks (*Quercus virginiana*) by heat. HortScience. 18(1):106-107.
- Croghan, Catherine F.; Robbins, Kathryn. 1986. Cankers caused by Botyrodiplodia gallae associated with oak sprout mortality in Michigan. Plant Disease. 70(1):76-77.
- Dorsey, C.K. 1967. Experiments to control acorn weevils with systemic insecticides. Forest Service. 13(4):390-396.
- Dorsey, C.K.; Tryon, E.H.; Carvell, K.L. 1962. Insect damage to acorns in West Virginia and control studies using granular systemic insecticides. Journal of Economic Entomology. 55 pp.
- Drooz, Arnold T., ed. 1985. Eastern forest insects. Misc. Publ. 1426. Washington, DC: U.S. Department of Agriculture, Forest Service. 608 pp.
- Galford, Jimmy R. 1986. Primary infestation of sprouting chestnut, red, and white oak acorns by Valentinia glandella (Lepidaptem: Blastrobasidae). Entomological News. 97(3):109-112. Page 173 of 404

- Gansner, David A.; Herrik, Owen W.; Debold, Paul S.; Achiavatti, Robert E. 1983. Changes in forest condition associated with gypsy moth. Journal of Forestry. 81(3):155-157.
- Gibson, Lester P. 1982. Insects that damage northern red oak acorns. Res. Pap. NE-220. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 6 pp.
- Hildebrand, D.C.; Schroth, M.N. 1967. A new species of *Erwinia* causing the drippy-nut disease of live oaks. Phytopathology. 57(3):250-253.
- Johnson, Leo J.; Law, Jay R. 1989. A five year record of change for a declining scarlet oak stand in the Missouri Ozarks. In: Proceedings of the seventh central hardwood forest conference; 1989 March 5-6; Carbondale, IL: 103-107.
- Knighten, John L.; Young, Carol H.; McCartney, Thomas C.; Anderson, Robert L. 1988. Resistance Screening Center procedures manual: a step-by-step guide used in the operational screening of southern pines for resistance to fusiform rust. [Unnumbered report]. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Region, Forest Pest Management. 62 pp.
- Oak, Steven W.; Huber, Cindy M.; Sheffield, Ray M. 1991. Incidence and impact of oak decline in western Virginia, 1986. Resour. Bull. SE-123. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 16 pp.
- Oak, Steven W.; Starkey, Dale A.; Dabney, Joseph M. 1989. Oak decline alters habitat in southern upland forests. In: Proceedings of the Southeastern association of fish and wildlife management agencies; 1988 November 7-10; Hilton Head, SC.
- Oak, Steven, W.; Triplett, Joey T. 1985. Cylindrocladium crotalariae kills container grown northern red oak. Tree Planters Notes. 36(2)6-9.
- Oliver, A.D.; Chapin, J.B. 1984. Curculio fulvus (Coleoptera: Curculionidae) and its effects on acorns of live oaks, Quercus virginiana Miller. Environmental Entomology. 13(6):1507-1510.
- Smyly, W.B.; Filer, T.H., Jr. 1977. Cylindrocladium scoparium associated with root rot and mortality of cherrybark oak seedlings. Plant Disease Reporter. 61(7):577-579.
- Soloman, J.D.; McCracken, F.I.; Anderson, R.L. [and others]. 1987. Oak pests: a guide to major insects, diseases, air pollution, and chemical injury. Prot. Rep. R8-PR7. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 69 pp.
- Starkey, Dale A.; Oak, Steven W.; Ryan, George W. [and others]. 1989. Evaluation of oak decline areas in the South. Prot. Rep. R8-PR17. U.S. Department of Agriculture, Forest Service, Southern Region. 36 pp.
- Tryon, E.H.; Carvell, Kenneth L.; Dorsey, C.K. 1968. Germination of acorns from oaks treated with systemic insecticides. Forest Science. 14(4):427-428.
- Vozzo, J.A. 1984. Insects and fungi associated with acorns of *Quercus* sp. In: Proceedings, cone and seed insects working party conference; 1983 July 31-August 6; Athens, GA. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 40-43.
- Williams, Charles E. 1989. Checklist of North American nut-infesting insects and host plant. Journal of Entomological Science. 24(4):550-562.

Sources of Oak Reproduction

Paul S. Johnson, North Central Forest Experiment Station, USDA Forest Service, Columbia, MO 65211

There are three juvenile growth forms of oak that silviculturists commonly term ABSTRACT "reproduction": (1) seedlings, (2) seedling sprouts, and (3) stump sprouts. A seedling is a form that has not experienced shoot dieback. Seedlings become seedling sprouts by surviving shoot dieback and resprouting one or more times. Seedling sprouts may persist and accumulate for decades in xeric forests and sometimes in mesic and hydric forests that are recurrently burned, grazed, or otherwise disturbed. Stump sprouts arise from the stumps of cut overstory trees after a thinning or final harvest and sometimes from the bases of fire-killed trees. When present before a silvicultural event such as clearcutting or shelterwood removal, all three growth forms are collectively called *advance reproduction*. The number, size, and spatial distribution of oak advance reproduction and the capacity of the parent stand (overstory) to produce oak stump sprouts can be used to assess a stand's oak regeneration potential, i.e., its capacity to occupy and dominate growing space at a specified time in the new stand. Because the population dynamics of oak reproduction vary greatly among species and different kinds of oak forests, this variation should be recognized in assessing stand regeneration potential.

REGENERATION TACTICS AND STRATEGIES OF OAKS

Oak forests can regenerate from seed and vegetatively from sprouts. Seeding, various modes of sprouting, and vegetative multiplication represent different tactics that oaks use in their regeneration strategy. Although all of the North American oak species rely to some extent on both seeding and sprouting, they differ greatly in their dependence on one tactic versus the other. For a given species, regeneration strategy may vary among regions and disturbance regimes. The large number of oak species that occur in the Eastern United States and the wide range of sites they occupy suggest that regeneration strategies are likely to vary greatly among the oaks.

Associated with these regeneration tactics are three commonly recognized growth forms that are collectively termed oak "reproduction"¹: (1) seedlings, (2) seedling sprouts, and (3) stump sprouts. Seedlings originate directly from seed

¹*Reproduction* herein refers to seedlings and juvenile sprouts of tree species. *Reproduce* refers to the creation of new individual plants, either from seed or by certain vegetative processes of plant multiplication (e.g., the spreading and subsequent separation of individual plants in lignotuberous oak species of the arid Southwest). *Regeneration* refers to the ecological *process* of renewing and sustaining populations of reproduction. These definitions may differ from those adopted by others (cf. Harper 1977).

and have not experienced shoot dieback. The seedling state typically lasts only one or a few years, depending on environment. Seedlings become seedling sprouts after dying back and resprouting one or more times. They may persist for decades, depending on species and environment. Stump sprouts arise from the stumps of cut overstory trees and sometimes from the bases of trees whose tops have been killed by fire or other injuries. When present before a silvicultural event such as clearcutting or shelterwood removal, all three growth forms are collectively termed *advance reproduction*. All living oaks from seedlings to mature trees thus can contribute to the *regeneration potential* of a stand. Regeneration potential is the capacity of a species to occupy and dominate growing space at a specified time in the new stand.

In the relatively droughty oak forests of the Missouri Ozarks, which are often dominated by some combination of black oak (Q. velutina Lam.), white oak (Q. alba L.), scarlet oak (Q. coccinea Muenchh.), and post oak (Q. stellata Wang.), the total density of oak advance reproduction typically ranges from 1,000 to 2,000 seedlings and seedling sprouts per acre. After a good acorn crop, 150 to 300 new oak seedlings per acre may become established (Sander 1979). But success in regenerating these forests largely depends on the relatively few oak seedling sprouts (e.g., about 400 per acre) with large root systems capable of supporting rapid shoot growth after overstory removal (Sander and others 1984). The regeneration of oaks in the Missouri Ozarks is thus largely dependent on sprouting, and with few exceptions these forests are seldom successionally displaced by other species.

Oaks of the arid Southwest may regenerate almost exclusively by sprouting. For example, Gambel oak (Q. gambelii Nutt.) produces three distinct root-like structures: lignotubers, rhizomes, and true roots (Tiedemann and others 1987). Lignotubers are burl-like structures with adventitious buds, which are the primary source of new shoots when tree crowns are killed. Several lignotubers may be connected by rhizomes, which have fewer buds. Rhizomes facilitate the development of wide-spreading clones (Muller 1951) that quickly develop from the seedling state (Christensen 1955). Later, the physical separation of individual lignotubers resulting from the death of connected lignotubers within a clone results in a form of plant multiplication that is important to the species' regeneration. Gambel oak thus regenerates primarily by sprouting from lignotubers and rhizomes, whereas seedlings are usually only a minor source of reproduction (Harper and others 1985). Muller (1951) concluded that rhizomes in oaks are associated with arid and semiarid climates where environments are especially unfavorable for germination and seedling survival. In addition to Gambel oak, other rhizomatous oaks in the United States include live oak (Q. virginiana Mill.) and several shrubby oaks native to the arid Southwest.

Northern red oak (*Q. rubra* L.), which occupies the middle ground between wet and dry extremes throughout its wide range in the eastern deciduous forest, is relatively flexible in its regeneration strategy. This flexibility is evidenced by large numbers of new red oak seedlings after bumper acorn crops (Johnson 1974), potentially rapid seedling shoot growth (Farmer 1975), ability to regenerate from seedlings established after final harvest (Johnson and others 1989), and capacity to sprout from large parent trees of advanced age (P. S. Johnson 1975, Wendel 1975). But unlike the xeric oak forests of the Missouri Ozarks and elsewhere, northern red oak forests are frequently displaced successionally by other species (Crow 1988, Johnson 1976, Loftis 1990a, Lorimer 1989, Nowacki and others 1990) Although sprouting is an important attribute of all oaks, there is growing evidence that seeding may be an important regeneration tactic of some bottomland oaks. For example, recent studies have shown that water oak (O. nigra L.) seedlings established after final harvest can be important contributors to the stocking of the new stand (Golden and Loewenstein 1991, Loewenstein 1992). But bottomland oaks also can sprout prolifically, which favors their development in stands disturbed by fire, grazing, and other factors (Aust and others 1985). These oaks, neither *obligate* seeders or sprouters, are flexible in their regeneration strategy. The relative importance of seeding as a regeneration tactic is also sometimes revealed by the number of seedlings that occur after a bumper acorn crop. For example, Nuttall oak (Q. nuttallii Palmer) seedlings exceeded 100,000 per acre in a southern bottomland forest after a bumper acorn crop (R. L. Johnson 1975). If only 0.1 percent of those seedlings (100 per acre) were competitively successful and well distributed, they would capture much of the available growing space within 20 years of overstory removal. The inherently rapid growth of Nuttall oak reproduction (Johnson 1981) and the productive sites they grow on may further reinforce this species' seeding strategy.

In bottomland oak forests, competing non-oak reproduction such as green ash (*Fraxinus pennsylvanica* Marsh.) and sweetgum (*Liquidambar styraciflua* L.) nevertheless often outgrow and suppress oak reproduction after overstory removal (Johnson and Krinard 1983, Aust and others 1985). Moreover, periods of high oak advance reproduction density in hydric forests are often followed by prolonged periods of low seedling density because of low seedling survival rates and infrequent acorn crops (R. L. Johnson 1975). There are thus frequent and prolonged periods with little or no oak advance reproduction. Consequently, bottomland oak forests are often successionally displaced by other species.

These anomalies in regeneration strategy among the oaks emphasize the difficulty of generalizing the regeneration problem across oak species and regions. Different oaks have different ways of solving their regeneration problems, and some species are more flexible than others (figure 1). In turn, flexibility in regeneration strategy is shaped by each species' environment and genetics.

SEEDLINGS AND SEEDLING SPROUTS

The unpredictable nature of flowering in oaks results in the irregular occurrence of acorn crops and thus new seedlings (Cecich 1991). On the average, most species produce a good acorn crop once every 3 or 4 years (Olson 1974). Numerous biotic and abiotic factors influence acorn viability, germination, initial seedling establishment, and survival. For example, dry weather, droughty soils, and freezing temperatures can reduce acorn viability and germination (Korstian 1927). Acorn crops also are frequently destroyed by unpredictable but frequent infestations of acorn weevils (Christisen and Kearby 1984). Most of the remaining acorns may be consumed by rodents, deer, birds, and other animals (Marquis and others 1976, Sork and others 1983). So, even after a bumper acorn crop, few acorns may be available for seedling production. Among the few remaining viable acorns, many fall into microsites unsuitable for germination and seedling establishment. Significant numbers of new oak seedlings thus occur as unpredictable population waves associated with bumper acorn for a patchy spatial distribution. The relatively infrequent occurrence of large seedling populations originating from one acorn crop (cohort) usually coincides with a bumper acorn crop combined with other fortuitous events, such as weather, that favor the preservation of acorn viability through fall and spring germination periods and low populations of acorn consumers.




Figure 1—Conceptualized dependence on seeding versus sprouting as a regeneration tactic for six oak species. [Diameters of circles are proportionate to species' flexibility in selecting the alternative tactic.]

Several studies have shown that survival of northern red oak seedlings originating from a single cohort is influenced by overstory density and other stand characteristics (figure 2). In general, reducing overstory density increases seedling survival and growth (Beck 1970; Crow 1992; Loftis 1988, 1990a). A dense layer of lower story trees, shrubs, or ground cover also can reduce seedling survival and growth (Beck 1970, Loftis 1990a, Scholz 1955). Other factors that can reduce oak seedling survival include animal browsing, insect defoliation, droughty soils, inadequate light, and frost (Crow 1992, Gottschalk 1988, Hanson and others 1987, Korstian 1927, McGee 1988).

During their first 9 years, numbers of northern red oak seedlings from a single cohort in North Carolina declined exponentially beneath the parent stand (figure 2A). Unfortunately, there are few detailed reports of oak seedling survival of similar duration. Short-term studies nevertheless point out the great variation in survival rate among seedlings of the same species growing in various regions representing different stand densities and different light and competition environments. They also establish the range over which we might reasonably expect the survival rates of oak advance reproduction to recur. For example $_{0,0}$ after

5 years, survival of individual cohorts of northern red oak seedlings ranged from about 0.16 to 0.86, depending on overstory density or understory competition (figure 2).




Figure 2—Survival curves for northern red oak seedlings originating from single acorn crops (cohorts) expressed as a negative exponential annual survival rate (K) from age 0 (germination) or age 1 year. [Survival probability $(P) = K^{\gamma}$, where y = years since initial population census, i.e., years since P = 1. (A) Under thinned and unthinned mesic forests in North Carolina (adapted from Loftis 1988); (B) In four overstory/understory release treatments in mesic forests in North Carolina (adapted from Beck 1970); (C) In clearcut, thinned, and unthinned stands treated with an herbicide in dry-mesic forests in northeastern Wisconsin (adapted from Crow 1992); and (D) Competing with three classes of ground cover under fully stocked mesic forests in southwestern Wisconsin (adapted from Scholz 1955)].

But the acorn producing capacity of a stand, and thus the rate of seedling input into oak forests, changes with time. Large trees usually produce more acorns than smaller trees because, other factors being equal, acorn production increases with crown area. In turn, crown area is correlated with bole diameter (Goodrum and others 1971). However, in some species there is a threshold diameter above and below which acorn production decreases (Downs 1944). Large, senescent oaks are poor acorn producers (Huntley 1983). The production of acorns per unit of crown area is also greater in open-grown trees than in forest-grown trees of the same size (Gysel 1956, Sharp 1958). Moreover, some trees are better acorn producers than others even when tree size and environmental factors are the same (Sharp and Sprague 1967). For example, in mature white oaks in Pennsylvania, only 30 percent produced any acorns even in good seed years (Sharp 1958), and an even smaller percent produced a good crop in those years (Sharp and Sprague 1967). So, although variation in stand structure and age can account for some of the variation in both the temporal and spatial variation in oak seedling establishment, inherent variation in acorn production among trees introduces an essentially random element into predicting seed and seedling inputs into oak forests.

Despite the seemingly complex problem of predicting the establishment of oak seedlings, more than half the variation in the density of black oak and white oak advance reproduction in xeric forests in northern Lower Michigan was explained by relatively simple measures of overstory density and structure (Johnson 1992). For both species, 55 percent of the variation in reproduction density was explained by total overstory basal area and the basal area of "large" trees presumed to be the primary seed producers. Large trees were defined as those at least 14 in. d.b.h. for black oak and those at least 12 in. d.b.h. for white oak (figure 3). The related models also showed that, per unit basal area of large trees, white oak was more efficient at producing seedlings than black oak. Moreover, high densities of black oak reproduction were favored under low density stands, whereas the reverse was true for white oak. Other studies have shown that topographic factors, stand history, and site quality also influence oak reproduction density (Arend and Scholz 1969, Carvell and Tryon 1961, Nowacki and others 1990, Ross and others 1986, Walters 1990).

Because the seedling stage is usually brief, seedling sprouts are the predominant form of oak reproduction in many, if not most, oak forests. Seedlings can sprout from dormant buds anywhere along the stem between the root collar and the terminal bud cluster. Dieback and resprouting seem to be important processes in the life of oak reproduction. Although recurrent shoot dieback is common to most hardwoods, it is especially prominent and ecologically important in the xerophytic oaks, which are morphologically and physiologically adapted to survival in environments subjected to repeated fire and drought (Abrams 1990, Grimm 1984, Wuenscher and Kozlowski 1971).

The natural environment of seedling sprouts of the xerophytic oaks imposes stresses that periodically decrease shoot mass and leaf area through shoot dieback. Surviving seedling sprouts thus develop increasingly greater root:shoot ratios as roots grow incrementally larger and shoots recurrently die back. In turn, high root:shoot ratio and large root mass enable oak reproduction to opportunistically respond to favorable environmental conditions by facilitating two or more long flushes of shoot growth (multiple flushing) during one growing season (Dickson 1991, Johnson 1979). Conditions that are usually favorable for the growth of a




Figure 3—Estimated density of black oak and white oak advance reproduction in relation to total stand basal area and the basal area of "large" trees of the same species (QBA) in xeric oak forests in northern Lower Michigan. [The trees comprising QBA are the presumed primary acorn producers. For white oak, QBA includes trees ≥ 12 in. d.b.h., and for black oak, trees ≥ 14 in. d.b.h. The models explain 55 percent of the variation in reproduction density of both species (from Johnson 1992).]

reproduction are created by timber harvesting and forces that destroy the overstory including fire, windthrow, insects, and disease.² Successional replacement of oak by oaks thus heavily depends on conditions that favor the long-term accumulation of oak reproduction with high root:shoot ratios and large root mass. Lacking those characteristics, oaks are usually at a competitive disadvantage. This is especially true of the reproduction of the xeromorphic upland species, which grow slowly even under optimal conditions until they develop the requisite root mass and root:shoot ratio. Shoot dieback thus may be an important aspect of the evolutionary development and adaptive strategy of oaks.

The accumulation of oak reproduction under a parent stand is one of the most important aspects of the regeneration ecology of oaks. Oak silviculturists call this "advance reproduction" because, in the even-aged management of oaks, it is present in advance of final harvest. Its presence and development largely determine the importance of oaks after natural or human-caused events that destroy or remove the parent stand. Oaks opportunistically capitalize on this accumulation process because it facilitates the capture of growing space when the overstory is destroyed or removed. This capacity largely depends on the characteristics of the competing vegetation and the accumulated population of oak seedling sprouts with large roots.

² An exception occurs when the removal of the protective overstory canopy precedes a spring frost that kills the new spring shoot growth of oak reproduction (Johnson 1979, McGee 1988).

The size distribution and age distribution of this population, in turn, depend on the balance of birth, death, and growth rates of reproduction intrinsic to each type of oak forest. Nevertheless, it is the number, size, and spatial distribution of all three classes of reproduction that largely express the total oak regeneration potential of a stand With the possible exception of some bottomland oak forests largely depends on perpetuating as species well adapted to capturing small canopy gaps (Crow 1988, Ehrenfeld 1980), they have the potential to do so if advance reproduction of sufficient size is For example, 10-ft.-tall advance reproduction of northern red oak captured canopy gaps as small as 1/25-acre in a mixed sugar maple-oak stand in preestablished reproduction (propagules) from one generation to the next. Whether the overstory is removed in small gaps or over large areas, the resulting spatial units of reproduction are even aged. Although oaks generally are not recognized southwestern Wisconsin (Lorimer 1983). dominated (Sander and others 1984). oak sustaining present. forests,

density by killing trees with thin, fire-sensitive bark, and (3) By killing the tops of Recurrent fire promotes the accumulation of oak reproduction in at least three oak reproduction, which increases the root:shoot ratio of those that survive by resprouting. The first two factors increase light on the forest floor, which in turn Although poorly adapted to surviving under shade, oak reproduction is well adapted These buds often remain an inch or more below the soil surface where they are (1) By eliminating or reducing the number of fire-sensitive understory competitors, including shrubs and shade-tolerant trees, (2) By reducing overstory reproduction. to surviving fire because of the concentration of dormant buds near the root collar. creates conditions more conducive to the accumulation of oak protected from lethally high temperatures (Korstian 1927). ways:

Many xeric oak forests appear to be relatively stable and show little evidence of Such forests occur on the drier sites throughout the central and eastern region including the Ohio Valley (Minckler and Woerheide 1965), the Appalachians (Ross succession to more shade tolerant or mesophytic species. Probably the largest However, even in that region, the successful development of advance oak North American forest of this type occurs in the Ozark Plateau in Missouri. reproduction varies with topographic features such as slope position and aspect, which collectively control light, heat, and soil moisture (Sander and others 1984). and others 1986), and the Lake States (Johnson 1966). Oak reproduction often may be absent or scarce on mesic or hydric sites in the absence of disturbance (Carvell and Tryon 1961, R. L. Johnson 1975, Will-Wolf 100,000 per acre in bottomland forests (R. L. Johnson 1975). When oak seedlings survival rates, most of the seedlings from a single cohort may die before the next good acorn crop occurs. The rapid rate of seedling disappearance results largely from the shade-intolerance of oak reproduction and the low light levels on the forest floor (Hanson and others 1987). In the absence of disturbance, these forests typically possess high overstory basal areas and multiple subcanopy layers (Braun Nevertheless, oak reproduction densities in these forests may at times do occur, they may represent only one or two acorn crops. Because of low Such vertical stratification occurs in mesic and hydric oak exceed 50,000 seedlings per acre in mesic forests (Tryon and Carvell 1958) and forests throughout the deciduous forest region. 1967, Loftis 1990b). 1991).

Microsites where oak advance reproduction becomes initially established in large numbers may not be where it ultimately survives to form populations with large root systems (Harrison and Werner 1984, Johnson 1966). Although initial establishment is favored by cool moist sites such as those on northeast-facing slopes, long-term survival is favored by the more southerly and neutral southeast and northwest aspects where overstory density and vertical stratification of crowns tend to be lower than on the more mesic northeast aspects (Sander and others 1984, Walters 1990, Carvell and Tryon 1961).

There is a general relation between site quality and regeneration success: the better the site the more difficult it is to regenerate oaks (Arend and Scholz 1969, Loftis 1990b, Lorimer 1989, Trimble 1973). Obtaining the accumulation of oak reproduction necessary for successful regeneration on highly productive sites requires recurrent disturbance. Historically, fire was associated with the origin of oak stands (Abrams 1992, McGee 1979). Thus, one way to sustain oak-dominated forests on productive sites may be prescribed burning. Based on a review of research on prescribed burning in eastern hardwoods and southern mixed pinehardwood stands, Van Lear and Waldrop (1988) concluded that fire, if correctly used, can be effective in regenerating oaks. However, fire also can kill or damage oaks and thus reduce the economic value of stands (Loomis 1973, Rouse 1986).

STUMP SPROUTS AND RELATED GROWTH FORMS

Stump sprouts originate from dormant buds at or near the base of the stump of a harvested overstory tree. In the silviculture of central hardwoods, overstory trees are defined as those 2 in. d.b.h. and larger (Roach and Gingrich 1968). However, that definition has not been universally adopted. The biological distinction between a stump sprout and a seedling sprout is nevertheless arbitrary because all oaks, from small seedlings to large standing trees, have some potential to produce basal sprouts when the parent stem is cut. When wind, fire, or other factors destroy an oak stand, sprouts also may develop from the bases of trees that have broken off or from standing trees with dead tops.

For several species of oaks, the percentage of stumps expected to produce sprouts after timber harvesting can be estimated from tree diameter and tree age (Johnson 1977). In general, the frequency of sprouting decreases with increasing tree diameter, age, and site quality (figure 4). But other factors, such as season of cutting and shading, also can affect stump sprouting in hardwoods. For some species of oaks, there is evidence that stumps sprout more frequently when trees are cut or killed during the dormant season than during the growing season (Clark and Liming 1953, Kays and others 1988). However, some of the live oaks of the Western United States sprout prolifically regardless of season of cutting (Longhurst Although sugar maple stumps exposed to full light sprouted more 1956). frequently than shaded stumps (Church 1960), similar responses of oak stumps to shading have not been reported. McGee and Bivens (1984) observed that numbers of stems in white oak sprout clumps were about the same for stumps that had been released from directly overtopping trees and stumps that were not released. Regardless of treatment, nearly 100 percent of the stumps of trees between 2 and 8 in. d.b.h. sprouted. Larger trees or those older than 60 years produced few or no sprouts.




Figure 4—Estimated sprouting probabilities for (A) black oak stumps and (B) white oak stumps in the Missouri Ozarks in relation to parent tree diameter and age (years) when site index is 65 ft. (base age 50) (from Johnson 1977).

Stump sprouts originating from pole-size and larger parent trees are, in effect, mature root systems connected to juvenile shoots. This root:shoot combination results in rapid height growth. During their first decade, open-grown stump sprouts can produce four or more flushes of shoot growth per year totaling 3 ft. or more even under droughty conditions (Cobb and others 1985, Johnson 1979, Reich and others 1980). The large root mass of stump sprouts and their large carbohydrate storage and absorptive capacity, together with other factors, facilitate multiple flushing in oaks. In contrast, multiple flushes are not produced in mature oaks, shaded seedlings and seedling sprouts, and small seedlings and seedling sprouts under water stress (Borchert 1976, Buech 1976, Cook 1941, Johnston 1941, Kienholz 1941, Longman and Coutts 1974).

Frequency of flushing and total shoot elongation in oaks usually decline as stems increase in size and age and as root systems approach their maximum size. The number of flushes in scarlet oak stump sprouts decreased from an average of about two per growing season the first year to one by the fourth growing season (Cobb and others 1985). Thus, by the fifth year, the pattern of shoot growth approached that of the single flush of a mature tree. The progression from multiple to single flushes may be attributable to a declining root:shoot ratio that results in increasingly longer periods for roots and shoot to restore "functional balance" after shoot elongation and leaf expansion (Borchert 1975).

The number and spatial distribution of sprouts around the stump also influence sprout growth. The importance of sprout distribution around the stump may be related to the pattern of vascular connections that develop between sprouts and the parent tree root system, with each stem helping to sustain a portion of the root system (Kharitonovich 1937, Kramer and Kozlowski 1979, Roth and Sleeth 1939, Wilson 1968).

In the Missouri Ozarks, the number of sprouts per stump was positively correlated with the early height growth of five oak species (Johnson 1977). The same relation also was observed for various oak species in other regions (P. S. Johnson 1975, Ross and others 1986, Schwarz 1907). Collectively, the observed spatial distribution and clump density effects suggest that numerous well-distributed sprouts maintain the parent tree root system and thus an efficient root-shoot feedback system that promotes rapid early height growth.

However, the apparent benefits of a balanced distribution of stems and high clump density are short lived. In Wisconsin, rapid growth of the dominant stem in unthinned clumps of 4- to 23-year-old northern red oak stump sprouts was associated with high clump density (P. S. Johnson 1975). Similarly, northern red oak clumps thinned to one stem as early as age 4 subsequently grew faster than stems in unthinned clumps (Johnson and Rogers 1984). This would seem to indicate that competition between stems in the same clump begins very early. However, 12-year-old northern red oak stump sprouts in Appalachian forests did not respond to clump thinning (Lamson 1988). Although northern red oak commonly initiates several dozen sprouts, stem crowding soon induces rapid stem mortality so that by the end of the first decade only four or fewer stems per clump typically remain. But this natural clump thinning process varies greatly among and within species (cf. P. S. Johnson 1975, Roth and Hepting 1969, Schwarz 1907).

The diameter of the parent tree and the correlated size of the root system also affect the growth of oak stump sprouts. For five species of oaks in the Missouri Ozarks, the correlation between stump diameter and height growth of the dominant stem within a sprout clump was consistently negative for 5-year-old sprouts of all species (Johnson 1977). However, the opposite was true of oak sprouts in Virginia (Ross and others 1986). Such discrepancies might be explained by the range of stump diameters observed in any given study. For example, data from a study of black oak and white oak sprouting that included trees ranging from less than 1 in. to more than 12 in. in basal diameter showed that the most rapid height growth occurred in clumps originating from 6-in. stumps (Johnson 1979). Sprouts from stumps larger or smaller than that grew less. The height growth of oak reproduction thus changes continuously, but not unidirectionally, from small advance reproduction to large-diameter overstory trees (figure 5).

Because of the large variation in number of stems per clump within a stump diameter class, we might expect roots and shoots of many young sprout clumps to be physiologically imbalanced. Accordingly, among the imbalanced clumps, rapid changes in clump structure would be expected as the clumps move toward functional balance. This view is supported by the initially large but rapidly decreasing variation in shoot growth among dominant shoots within black oak and white oak sprout clumps during the first 4 years (Johnson 1979). The large amount of unexplained variation in the relation between shoot growth and stump diameter also reflects rapidly changing root-shoot relations. Other factors that may be significant sources of variation in the shoot growth of oak sprouts include site quality, genetic variation, competition, parent tree age, and season of cutting. The significance of these factors may vary among oak species and regions.




Figure 5—Relation between height growth of black oak and white oak reproduction and parent stem basal diameter during the first 4 years after clearcutting in the Missouri Ozarks (adapted from Johnson 1979).

ASSESSING REGENERATION POTENTIAL

Oak stands can be regenerated by completely removing the overstory on small to large areas using clearcutting, shelterwood, or group selection methods (Hannah 1987, Sander and Clark 1971) and possibly by using other methods that substantially reduce overstory density. Those regeneration methods are used because oak seedlings and sprouts cannot grow into the overstory as long as they are heavily shaded. But all the oak seedlings and sprouts present at the time of a regeneration cut do not contribute to the future stand. A clear picture of the future stand is thus hidden, or encoded, in certain characteristics of the present stand. The most important of these characteristics are the number, size, and spatial distribution of the oak advance reproduction; the composition and structure of the overstory; and site quality (Loftis 1990a, Sander and others 1984).

If the parent oak stand is to be replaced by another generation of oaks, enough oak reproduction of sufficient size must be present at the time of final harvest (Carvell and Tryon 1961, Sander and Clark 1971, Trimble 1973). But even when the essential characteristics of the oak reproduction are known, it is difficult to assess the contribution of oaks to the regenerated stand because of the unpredictable growth of reproduction. Much of this uncertainty is related to the difficulty of obtaining an accurate measure of root size. Growth of oak reproduction largely depends on root size, which may be much larger and older than the above-ground stem. Recurrent stem dieback and resprouting of oak reproduction beneath an Page 186 of 404 overstory may go on for decades before the reproduction expresses its growth potential. This potential is finally expressed when the overstory is harvested or when it is destroyed by natural events such as fire, windthrow, and disease. Although some silviculturists can predict general regeneration trends after timber harvesting based on experience and observation, more quantitative methods are needed to accurately assess the *probable* contribution of oak reproduction to future stand stocking.

Unfortunately, root size is impossible to measure in practical application. The next best measure of the growth potential of reproduction is probably its diameter just above the ground. However, the relation between shoot growth and basal stem diameter, although usually statistically significant, is relatively weak (Dey 1991, Johnson 1979, Sander 1971). To overcome this problem, some predictive regeneration models employ probabilistic methods to assess the importance of oaks in the future stand (Dey 1991, Loftis 1990a, Sander and others 1984). Other regeneration models use other methods (Waldrop and others 1986, Marquis and others 1984). Regardless of their derivation, regeneration models are useful silvicultural tools for assessing stand regeneration potential. Using such models usually requires an inventory of the advance reproduction and the overstory. The practical application of predictive regeneration models thus does not differ greatly from the application of growth and yield models.

Because site quality and competition vary greatly among different kinds of oak forests, ecological classification is a silvicultural tool that can aid in assessing stand regeneration potential. The objective of ecological classification is to define landscape units (ecosystems) *within* which there is great ecological similarity and *among* which there is dissimilarity (Rowe and Sheard 1981). In principle, all stands representing a given unit respond similarly to natural forces and processes and thus to management practices. This does not mean that stands of similar overstory composition and structure are necessarily ecologically equivalent units. In ecological classification, factors such as climate, physiography, parent materials, soils, and lesser vegetation, together with the composition and structure of the overstory, are used to define ecological units. This feature distinguishes ecological classification from traditional forest cover-type descriptions (e.g., Eyre 1980). Defined units thus can be used to distinguish among oak-dominated ecosystems that are superficially similar but ecologically quite different.

Both predictive regeneration models and experience-based assessments of regeneration potential can be developed in conjunction with ecological classification. In either case, increased accuracy of predictions should result because similarities in site, competition, and other factors are intrinsic to each defined ecological unit. For example, from our general understanding of oak regeneration processes, we would expect the rate of accumulation and size distribution of oak reproduction to vary substantially among different kinds of oak forests. Ecological forest site classification and its development and application have been discussed elsewhere (Barnes and others 1982, Cleland and others 1985, Kotar 1991, Rowe 1981). There are classification systems for several areas of the eastern hardwood region (e.g., Albert and others 1986, Hix 1988, Host and Pregitzer 1991, Kotar and others 1988, McNab 1987, Miller 1981, Smalley 1986).

Whether assessments of regeneration potential are based on quantitative models or on experience and general observation, the regeneration progential of all sources of oak reproduction should be considered. But the relative importance of the various sources of reproduction and their population dynamics can be expected to vary among oak species and across the landscapes where they occur. An understanding of the biology and population dynamics of all sources of oak reproduction is therefore prerequisite to a practical understanding of the regeneration ecology of oaks and thus the silvicultural methods required to sustain oak forests.

LITERATURE CITED Abrams, Marc D. 1990. Adaptations and responses to drought in *Quercus* species of North America. Tree Physiology. 7(1-4):227-238.

- Abrams, Marc D. 1992. Fire and the development of oak forests. BioScience. 42(5):346-353.
- Albert, Dennis A.; Denton, Shirley R.; Barnes, Burton V. 1986. Regional landscape ecosystems of Michigan. Ann Arbor, MI: University of Michigan, School of Natural Resources. 32 pp.
- Arend, John L.; Scholz, Harold F. 1969. Oak forests of the Lake States and their management. Res. Pap. NC-31. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 36 pp.
- Aust, W.M.; Hodges, J.D.; Johnson, R.L. 1985. The origin, growth and development of natural, pure, even-aged stands of bottomland oak. In: Shoulders, E., ed. Proceedings, 3rd biennial southern silvicultural research conference; 1984 November 7-8; Atlanta, GA. Gen. Tech. Rep. SO-54. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 163-170.
- Barnes, Burton V.; Pregitzer, Kurt S.; Spies, Thomas A; [and others]. 1982. Ecological forest site classification. Journal of Forestry. 80(8):493-498.
- Beck, D.E. 1970. Effect of competition on survival and height growth of red oak seedlings. Res. Pap. SE-56. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 7 pp.
- Borchert, Rolf. 1975. Endogenous shoot growth rhythms and indeterminate shoot growth in oak. Physiologia Plantarum. 35(2):152-157.
- Borchert, Rolf. 1976. Size and shoot growth patterns in broadleaved trees. In: Fralish, James.; Weaver, George T.; Schlesinger, R.C., eds. Proceedings, Central hardwood forest conference 1; 1976 October 17-19; Carbondale, IL: Southern Illinois University: 221-230.
- Braun, E. Lucy. 1967. Deciduous forests of eastern North America. New York: Hafner. 596 pp.
- Buech, Richard R. 1976. Tree shoot elongation in northern Wisconsin and relationships with temperature and precipitation. Canadian Journal of Forest Research. 6(4):487-498.
- Carvell, K.L.; Tryon, E.H. 1961. The effect of environmental factors on the abundance of oak regeneration beneath mature oak stands. Forest Science. 7(2):98-105.
- Cecich, Robert A. 1991. Seed production in oak. In: Laursen, Steven B.; DeBoe, Joyce F., eds. Proceedings, Oak resource in the Upper Midwest, implications for management; 1991 June 3-6; Winona, MN. Publ. NR-BU-5663-S. St. Paul, MN: University of Minnesota Extension Service: 125-131.
- Christensen, Earl M. 1955. Ecological notes on the mountain brush in Utah. Proceedings, Utah Academy of Science, Arts, and Letters. 32:107-111.

- Christisen, Donald M.; Kearby, William H. 1984. Mast measurement and production in Missouri (with special reference to acorns). Terrestrial Series no. 13. Jefferson City, MO: Missouri Department of Conservation. 34 pp.
- Church, Thomas W., Jr. 1960. Factors affecting the development and survival of sugar maple sprouts. In: Proceedings, Society of American Foresters meeting, Forestry's expanding horizons; 1960 November 13-16; Washington, DC. Washington, DC: Society of American Foresters: 32-35.
- Clark, F. Bryan; Liming, Franklin G. 1953. Sprouting of blackjack oak in the Missouri Ozarks. Tech. Pap. 137. Columbus, OH: U.S. Department of Agriculture, Forest Service, Central States Forest Experiment Station. 22 pp.
- Cleland, David T.; Hart, James B.; Pregitzer, Kurt S.; [and others]. 1985. Classifying oak ecosystems for management. In: Johnson, James E., ed. Proceedings of the challenges in oak management and utilization; 1985 March 28-29; Madison, WI: University of Wisconsin Cooperative Extension Service: 120-134.
- Cobb, S.W.; Miller, A.E.; Zahner, R. 1985. Recurrent shoot flushes in scarlet oak stump sprouts. Forest Science. 31(3):725-730.
- Cook, David B. 1941. The period of growth in some northeastern trees. Journal of Forestry. 39(11):956-959.
- Crow, T.R. 1988. Reproductive mode and mechanisms for self-replacement of northern red oak (Quercus rubra)-a review. Forest Science. 34(1):19-40.
- Crow, T.R. 1992. Population dynamics and growth patterns for a cohort of northern red oak (*Quercus rubra*) seedlings. Oecologia. 91(2):192-200.
- Dey, Daniel C. 1991. A comprehensive Ozark regenerator. Columbia, MO: University of Missouri. 283 pp. Dissertation.
- Dickson, Richard E. 1991. Episodic growth and carbon physiology in northern red oak. In: Laursen, Steven B.; DeBoe, Joyce F., eds. Proceedings of the oak resource in the Upper Midwest, implications for management; 1991 June 3-6; Winona, MN. Publ. NR-BU-5663-S. St. Paul, MN: University of Minnesota Extension Service: 117-124.
- Downs, Albert A. 1944. Estimating acorn crops for wild life in the southern Appalachians. Journal of Wildlife Management. 8(4):339-340.
- Ehrenfeld, John G. 1980. Understory response to canopy gaps of varying size in a mature oak forest. Bulletin of the Torrey Botanical Club. 107(1):29-41.
- Eyre, F.H., ed. 1980. Forest cover types of the United States and Canada. Washington, DC: Society of American Foresters. 148 pp.
- Farmer, R.E., Jr. 1975. Growth and assimilation rate of juvenile northern red oak: effects of light and temperature. Forest Science. 21(4):373-381.
- Golden, Michael S.; Loewenstein, Edward F. 1991. Regeneration of tree species 7 years after clearcutting in a river bottom in central Alabama. In: Coleman, Sandra S.; Neary, Daniel G. comp. and eds. Proceedings of the 6th biennial southern silvicultural research conference; 1990. October 30-November 1; Memphis, TN. Gen. Tech Rep. SE-70 (vol. 1). Asheville, NC: U.S Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 76-83.
- Goodrum, P.D.; Reid, V.H.; Boyd, C.E. 1971. Acorn yields, characteristics, and managemen criteria of oaks for wildlife. Journal of Wildlife Management. 35(3):520-532.
- Gottschalk, Kurt W. 1988. Gypsy moth and regenerating Appalachian hardwood stands. In: Smith H. Clay; Perkey, Arlyn W.; Kidd, William E., Jr., eds. Proceedings of the guidelines fo Page 189 of 404

regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV. Publ. 88-03. Bethesda, MD: Society of American Foresters: 241-254.

- Grimm, Eric C. 1984. Fire and other factors controlling the Big Woods vegetation of Minnesota in the mid-nineteenth century. Ecological Monographs. 54(3):291-311.
- Gysel, Leslie W. 1956. Measurement of acorn crops. Forest Science. 2(1):305-313.
- Hannah, Peter R. 1987. Regeneration methods for oaks. Northern Journal of Applied Forestry. 4(2):97-101.
- Hanson, P.J.; Isebrands, J.G.; Dickson, R.E. 1987. Carbon budgets of *Quercus rubra* L. seedlings at selected stages of growth: influence of light. In: Hay, Ronald L.; Woods, Frank W.; DeSelm, Hal, eds. Proceedings of the central hardwood forest conference 6; 1987 February 24-26; Knoxville, TN: University of Tennessee: 269-276.
- Harper, John L. 1977. Population biology of plants. New York: Academic Press. 892 pp.
- Harper, Kimball T.; Wagstaff, Fred J.; Kunzler, Lynn M. 1985. Biology and management of the Gambel oak vegetative type: a literature review. Gen. Tech. Rep. INT-179. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. 31 pp.
- Harrison, Janet S.; Werner, Patricia A. 1984. Colonization of oak seedlings into a heterogeneous successional habitat. Canadian Journal of Botany. 62(3):559-563.
- Hix, David M. 1988. Multifactor classification and analysis of upland hardwood forest ecosystems of the Kickapoo River watershed, southwestern Wisconsin. Canadian Journal of Forest Research. 18(11):1405-1415.
- Host, George E.; Pregitzer, Kurt S. 1991. Ecological species groups for upland forest ecosystems of northwestern Lower Michigan. Forest Ecology and Management. 43:87-102.
- Huntley, Jimmy C. 1983. Squirrel den tree management; reducing incompatibility with timber production in upland hardwoods. In: Jones, Earle P., Jr., ed. Proceedings of the 2nd biennial southern silvicultural research conference; 1982 November 4-5; Atlanta, GA. Gen. Tech. Rep. SE-24. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 488-495.
- Johnson, Paul S. 1966. Vegetational developments after logging in southern Michigan xeric oak forests. East Lansing, MI: Michigan State University. 204 pp. Dissertation.
- Johnson, Paul S. 1974. Survival and growth of northern red oak seedlings following a prescribed burn. Res. Note NC-177. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 3 pp.
- Johnson, Paul S. 1975. Growth and structural development of red oak sprout clumps. Forest Science. 21(4):413-418.
- Johnson, Paul S. 1976. Modal development of regeneration in clearcut red oak stands in the Driftless Area. In: Fralish, James S.; Weaver, George T.; Schlesinger, Richard C., eds. Proceedings of the central hardwood forest conference 1; 1976 October 17-19; Carbondale, IL: Southern Illinois University: 455-475.
- Johnson, Paul S. 1977. Predicting oak stump sprouting and sprout development in the Missouri Ozarks. Res. Pap. NC-149. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 9 pp.
- Johnson, Paul S. 1979. Shoot elongation of black oak and white oak sprouts. Canadian Journal of Forest Research. 9(4):489-494.

- Johnson, Paul S. 1992. Oak overstory/reproduction relations in two xeric ecosystems in Michigan. Forest Ecology and Management. 48(3):233-248.
- Johnson, Paul S.; Rogers, Robert. 1984. Predicting 25th-year diameters of thinned stump sprouts of northern red oak. Journal of Forestry. 82(10):616-619.
- Johnson, Paul S.; Jacobs, Rodney D.; Martin A. Jeff; [and others]. 1989. Regenerating northern red oak: three successful case histories. Northern Journal of Applied Forestry. 6(4):174-178.
- Johnson, Robert L. 1975. Natural regeneration and development of Nuttall oak and associated species. Res. Pap. SO-104. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 12 pp.
- Johnson, Robert L. 1981. Oak seeding-it can work. Southern Journal Applied Forestry. 5(1):28-33.
- Johnson, Robert L.; Krinard, R.M. 1983. Regeneration in small and large sawtimber sweetgum-red oak stands following selection and seed tree harvest: 23-year results. Southern Journal of Applied Forestry. 7(4):176-184.
- Johnston, John P. 1941. Height-growth periods of oak and pine reproduction in the Missouri Ozarks. Journal of Forestry. 39(1):67-68.
- Kays, Jonathon S.; Smith, David Wm.; Zedaker, Shephard M.; [and others]. 1988. Factors affecting natural regeneration of Piedmont hardwoods. Southern Journal of Applied Forestry. 12(2):98-102.
- Kharitonovich, F.M. 1937. Poroslevoe lesovozobnovlenie. [Reforestation by coppice shoots.] In: Sorokin, I.F., ed.; Polezoschitnye lesnya polosy [Field-protecting forest belts]; Moscow: 175-239.
- Kienholz, R. 1941. Seasonal course of height growth in some hardwoods in Connecticut. Ecology. 22(3):249-258.
- Korstian, Clarence F. 1927. Factors controlling germination and early survival in oaks. School of Forestry Bulletin 19. New Haven, CT: Yale University. 115 pp.
- Kotar, John. 1991. Importance of ecological classification in oak management. In: Laursen, Steven B.; DeBoe, Joyce F., eds. Proceedings of the oak resource in the Upper Midwest, implications for management; 1991 June 3-6; Winona, MN. Publ. NR-BU-5663.S. St. Paul, MN: University of Minnesota Extension Service: 132-140.
- Kotar, John; Kovach, Joseph A.; Locey, Craig T. 1988. Field guide to forest habitat types of northern Wisconsin. Madison, WI: University of Wisconsin and Wisconsin Department of Natural Resources.
- Kramer, Paul J.; Kozlowski, Theodore T. 1979. Physiology of woody plants. New York: Academic Press. 811 pp.
- Lamson, Neil I. 1988. Role of stump sprouts in regenerating Appalachian hardwood stands. In: Smith, H. Clay; Perkey, Arlyn W.; Kidd, William E., Jr., eds. Proceedings of the guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV. Publ. 88-03. Bethesda, MD: Society of American Foresters: 31-37.
- Loewenstein, Edward Francis. 1992. Relationships between stem origin and growth characteristics of three bottomland hardwood species on a seven-year-old clearcut area in the Coastal Plain of Alabama. Auburn, AL: Auburn University. 107 pp. Thesis.
- Loftis, David L. 1988. Regenerating oaks on high-quality sites, an update. In: Smith, H. Clay; Perkey, Arlyn W.; Kidd, William E., Jr., eds. Proceedings of the guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV. Publ. 88-03. Bethesda, MD: Society of American Foresters: 199-209.

- Loftis, David L. 1990a. Predicting post-harvest performance of advance red oak reproduction in the southern Appalachians. Forest Science. 36(4):908-916.
- Loftis, David L. 1990b. A shelterwood method for regenerating red oak in the southern Appalachians. Forest Science. 36(4):917-929.
- Longhurst, William M. 1956. Stump sprouting of oaks in response to seasonal cutting. Journal of Range Management. 9(4):194-196.
- Longman, K.A.; Coutts, M.P. 1974. Physiology of the oak tree. In: Morris, M.G.; Perring, F.H., eds. Berkshire, UK: The British oak: 194-221.
- Loomis, Robert M. 1973. Estimating fire-caused mortality and injury in oak-hickory forests. Res. Pap. NC-94. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 6 pp.
- Lorimer, Craig G. 1983. Eighty-year development of northern red oak after partial cutting in a mixed-species Wisconsin forest. Forest Science. 29(2):371-383.
- Lorimer, Craig G. 1989. The oak regeneration problem: new evidence on causes and possible solutions. Madison: University of Wisconsin, School of Natural Resources, Department of Forestry, Forest Resource Analyses No. 8. 31 pp.
- Marquis, David A; Eckert, Philip L.; Roach, Benjamin A. 1976. Acorn weevils, rodents, and deer all contribute to oak-regeneration difficulties in Pennsylvania. Res. Pap. NE-356. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 5 pp.
- Marquis, David A.; Ernst, Richard L.; Stout, Susan L. 1984. Prescribing silvicultural treatments in hardwood stands of the Alleghenies. Gen. Tech. Rep. NE-96. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 90 pp.
- McGee, Charles E. 1979. Fire and other factors related to oak regeneration. In: Holt, Harvey A.; Fischer, Burnell C., eds. Proceedings of the regenerating oaks in upland hardwood forests. The 1979 John S. Wright forestry conference; 1979 November 22-23; West Lafayette, IN: Purdue Research Foundation, Purdue University: 75-80.
- McGee, Charles E. 1988. Spring weather, canopy removal, and early budbreak threaten oak seedings. The Journal of the Elisha Mitchell Scientific Society. 104(3):108-115.
- McGee, Charles E.; Bivens, Donald L. 1984. A billion overtopped white oak-assets or liabilities? Southern Journal of Applied Forestry. 8(4):216-220.
- McNab, W. Henry. 1987. Rationale for a multifactor forest site classification system for the southern Appalachians. In: Hay, Ronald L.; Woods, Frank W.; DeSelm, Hal, eds. Proceedings of the central hardwood forest conference 6; 1987 February 24-26; Knoxville, TN: University of Tennessee: 283-294.
- Miller, Mack R. 1981. Ecological land classification terrestrial subsystem, a basic inventory system for planning and management on the Mark Twain National Forest. Rolla, MO: U.S. Department of Agriculture, Forest Service, Mark Twain National Forest. 56 pp.
- Minckler, Leon S.; Woerheide, John D. 1965. Reproduction of hardwoods 10 years after cutting as affected by site and opening size. Journal of Forestry. 63(2):103-107.
- Muller, Cornelius H. 1951. The significance of vegetative reproduction in *Quercus*. Madroño. 11(3):129-137.
- Nowacki, Gregory J.; Abrams, Marc D.; Lorimer, Craig G. 1990. Composition, structure, and historical development of northern red oak stands along an edaphic gradient in north-central Wisconsin. Forest Science. 36(2):276-292. Page 192 of 404

- Olson, David F., Jr. 1974. *Quercus* L. Oak. In: Schopmeyer, C.S., tech. coord. Seeds of woody plants in the United States. Agric. Handb. 450. Washington, DC: U.S. Department of Agriculture, Forest Service: 692-703.
- Reich, P.B.; Teskey, R.O.; Johnson, P.S.; [and others]. 1980. Periodic root and shoot growth in oak. Forest Science. 26(4):590-598.
- Roach, Benjamin A.; Gingrich, Samuel F. 1968. Even-aged management for upland central hardwoods. Agric. Handb. 355. Washington, DC: U.S. Department of Agriculture, Forest Service. 39 pp.
- Ross, Michael S.; Sharik, Terry L.; Smith, David Wm. 1986. Oak regeneration after clear felling in southwest Virginia. Forest Science. 32(1):157-169.
- Roth, Elmer R.; Hepting, George H. 1969. Prediction of butt rot in newly regenerated sprout oak stands. Journal of Forestry. 67(10):756-760.
- Roth, Elmer R.; Sleeth, Bailey. 1939. Butt rot in unburned sprout oak stands. Tech. Bull. 684. Washington, DC: U.S. Department of Agriculture. 43 pp.
- Rouse, Cary. 1986. Fire effects in northeastern forests: oak. Gen. Tech. Rep. NC-105. St. Paul: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 6 pp.
- Rowe, J. Stan; Sheard, John W. 1981. Ecological land classification: a survey approach. Environmental Management. 5(5):451-464.
- Sander, Ivan L. 1971. Height growth of new oak sprouts depends on size of advance reproduction. Journal of Forestry. 69(11):809-811.
- Sander, Ivan L. 1979. Regenerating oaks. In: Proceedings of the national silviculture workshop; 1979 September 17-21; Charleston, SC. Washington, DC: U.S. Department of Agriculture, Forest Service: 212-221.
- Sander, Ivan L.; Clark, F. Bryan. 1971. Reproduction of upland hardwood forests in the Central States. Agric. Handb. 405. Washington, DC: U.S. Department of Agriculture, Forest Service. 25 pp.
- Sander, Ivan L.; Johnson, Paul S.; Rogers, Robert. 1984. Evaluating oak advance reproduction in the Missouri Ozarks. Res. Pap. NC-251. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 11 pp.
- Scholz, Harold F. 1955. Effect of scarification on the initial establishment of northern red oak reproduction. Tech. Note 425. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Lakes States Forest Experiment Station. 2 pp.
- Schwarz, G. Frederick. 1907. The sprout forests of the Housatonic Valley of Connecticut. Forestry Quarterly. 5(2):121-153.
- Sharp, Ward M. 1958. Evaluating mast yields in the oaks. Bulletin 635. University Park: Pennsylvania State University, College of Agriculture, Agricultural Experiment Station. 22 pp.
- Sharp, Ward M.; Sprague, Vance G. 1967. Flowering and fruiting in the white oaks. Pistillate flowering, acorn development, weather, and yields. Ecology. 48(2):243-251.
- Smalley, Glendon W. 1986. Classification and evaluation of forest sites on the northern Cumberland Plateau. Gen. Tech. Rep. SO-60. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 74 pp.
- Sork, Victoria L.; Stacey, Peter; Averett, John E. 1983. Utilization of red oak acorns in non-bumper crop year. Oecologia. 59(1):49-53.

- Tiedemann, A.R.; Clary, W.P.; Barbour, R.J. 1987. Underground systems of Gambel oak (Quercus gambelii) in central Utah. American Journal of Botany. 74(7):1065-1071.
- Trimble, George R., Jr. 1973. The regeneration of Central Appalachian hardwoods with emphasis on the effects of site quality and harvesting practice. Res. Pap. NE-282. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 14 pp.
- Tryon, E.H.; Carvell, K.L. 1958. Regeneration under oak stands. Bulletin 424T. Morgantown: West Virginia University, Agricultural Experiment Station. 22 pp.
- Van Lear, David H.; Waldrop, Thomas A. 1988. Effects of fire on natural regeneration in the Appalachian Mountains. In: Smith, H. Clay; Perkey, Arlyn W.; Kidd, William E., Jr., eds. Proceedings of the guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV. Publ. 88-03. Bethesda, MD: Society of American Foresters: 56-70.
- Waldrop, T.A.; Buckner, E.R.; Shugart, H.H., Jr.; [and others]. 1986. FORCAT: a single tree model of stand development following clearcutting on the Cumberland Plateau. Forest Science. 32(2):297-317.
- Walters, Russell S. 1990. Site quality, fire, and herbicide effects on establishment, growth, and development of regeneration three years after partial cutting of oak stands. Syracuse: State University of New York. 197 pp. Dissertation.
- Wendel, G.W. 1975. Stump sprout growth and quality of several Appalachian hardwood species after clearcutting. Res. Pap. NE-329. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 9 pp.
- Will-Wolf, Susan. 1991. Role of fire in maintaining oaks in mesic oak maple forests. In: Laursen, Steven B.; DeBoe, Joyce F., eds. Proceedings of the oak resource in the Upper Midwest, implications for management; 1991 June 3-6; Winona, MN. Publ. NR-BU-5663-S. St. Paul, MN: University of Minnesota Extension Service: 27-33.
- Wilson, Brayton F. 1968. Red maple stump sprout development the first year. Harvard Forest Paper 18. Petersham, MA: Harvard University. 10 pp.
- Wuenscher, James E.; Kozlowski, Theodore T. 1971. Relationship of gas-exchange resistance to tree-seedling ecology. Ecology. 52(6):1016-1023.

Page 195 of 404

Prediciting & Modeling Oak Regeneration

Predicting Oak Regeneration—State of the Art

David L. Loftis, Southeastern Forest Experiment Station, USDA Forest Service, Asheville, NC 28806

ABSTRACT Early efforts to predict species composition of regeneration (including oaks) resulting from harvest cuts attempted to link presumed species successional status and/or shade tolerance rating to the level of disturbance or residual stand stocking. While this traditional view of forest succession seems applicable in some situations, there are so many exceptions that its applicability as a model for prediction is questionable. Implicit in more recent efforts to predict oak regeneration following harvest cuts is a contemporary set of succession concepts that deal with population dynamics, initial floristic composition, vital species attributes, or life history characteristics. These models recognize that oaks are generally regarded as advance-growth-dependent species and predict the expected oak component in regeneration as a function of the amount and size distribution of advance growth—stump sprout potential and advance reproduction—in an existing mature stand.

INTRODUCTION In this brief introduction to the next two papers, I would like to focus on a few forest succession concepts that are relevant to regeneration in general, and to prediction in particular. If, for a particular stand, one has a goal of regenerating a new oak stand or maintaining an oak component in the new stand, having the ability to predict the consequences of a range of regeneration alternatives is critical in choosing an appropriate silvicultural treatment.

RELAY FLORISTICS VS. INITIAL FLORISTIC COMPOSITION

The first two concepts are posed as alternative views of vegetation development (Egler 1954). Relay floristics (figure 1) presents a very familiar pattern of development, one that many people would associate with Clements (1916). Well-defined seral stages, beginning with pioneer species, successively occupy a site, culminating in a relatively stable community of climax species. We also usually tend to think of pioneer species as shade intolerant and of climax species as shade tolerant.

In using this concept to predict the outcome of the application of various regeneration methods, we have associated species we classify as pioneer and shade intolerant with clearcutting, climax and shade tolerant species with selection methods, and species that are intermediate in their successional status and shade tolerance with shelterwood. And one frequently hears that "cutting sets succession back to an earlier stage."





Figure 1—Relay floristics model.

However, reconciling this neat scheme with observed outcomes of regeneration methods is problematic, particularly in hardwood forests. For example, clearcutting a stand of oaks on a relatively xeric site frequently results in a new stand dominated by oaks, but oaks are not usually considered pioneer species. Regeneration cuts of any kind in some hardwood forests result in excellent representation of species we regard as climax and shade tolerant in the new stand. Further, the application of different regeneration methods in some forest types frequently results in the same species composition. In short, categorization of species by presumed successional status lends little to our ability to predict species composition resulting from regeneration cutting.

As an alternative to relay floristics, Egler proposed a concept he called initial floristic composition (figure 2). This concept suggests that species composition of vegetation following disturbance is determined by the propagules that exist on the site at the time of the disturbance and those which arrive early in the process of stand development. Changing species dominance over time results from differential growth and development of species present. Interestingly, it was Clements (1916) who first made this observation. According to Clements, logging does not initiate a successional sequence because propagules of a variety of species are left after logging. Rather, it creates a dysclimax.

For forest trees, then, species composition following disturbance is determined by the initial load of propagules—stump- and root-sprouts and advance reproduction of the various species that occur in the stand, and by new seedlings that become established soon after the disturbance from buried seed or from seed blown or carried into the stand. The essential feature of the initial floristic composition Page 198 of 404 concept that makes it useful for prediction modeling is that the individuals of species that form the dominant tree canopy 50 to 100 years after disturbance come from the initial load of propagules. A corollary to this concept is that variations in species composition among stands 50 to 100 years after similar disturbances results from variations among stands in the initial loads of propagules.

A number of investigators have recognized the general applicability of the initial floristic composition concept to hardwood stand development following disturbance (e.g. Oliver 1981, Leopold and others 1985, Shugart 1984, Drury and Nisbet 1973). From the standpoint of developing prediction models, we can model the post-disturbance development of existing advance growth—both advance reproduction and stump or root-sprout potential—and we can deal in probabilistic sense with input and development of new seedlings.





VITAL ATTRIBUTES OF SPECIES

Noble and Slatyer (1980) suggested that species composition of stands that develop after disturbance could be predicted from three vital attributes of species:

1. The method of arrival or persistence of a species at a site during or after a disturbance.

- 2. The ability to grow to maturity in the developing stand.
- 3. The time needed for an individual of a species to reach critical life stages.

Because of time constraints, I will discuss only the the first two of these vital attributes.

Several authors have addressed the first attribute by examining the primary reproduction source that is characteristic for various species (Beck 1988, Kelty 1988, Johnson 1989). Oaks, for example, are considered to be highly dependent on advance growth; i.e., they persist through a disturbance as advance reproduction and stump sprouts. Yellow-poplar is an example of a species that usually becomes established after disturbance as new seedlings from seed stored in the forest floor.

The second attribute, the ability to grow to maturity in the developing stand, has been, and is being, addressed more fully for oaks than for any group of hardwood species of which I am aware. In general, the fairly strong relationship between preharvest size of advance growth and its post-harvest development are being used to predict the amount of oak one would expect in the next stand.

- **FINAL NOTE** Initial floristic composition, vital attributes of species, and concepts from population biology are the ecological bases for the oak regeneration potential concept (P. Johnson, this Proceedings) that provides the framework for developing models to predict how much oak we might expect in a new stand following a regeneration cut. Some models already exist, but prediction models are lacking for many ecosystems where oaks are important and for regeneration methods other than clearcutting. But the development and application of these prediction models will help us choose among alternatives. For managers the application of these prediction models has one very clear imperative: a commitment to collect adequate data about existing stand conditions.
- LITERATURE CITED Beck, Donald E. 1988. Regenerating cove hardwood stands. In: H. Clay Smith, Arlyn W. Perkey, and William E. Kidd, Jr., eds. Guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV: West Virginia University Books: 56-166.

Clements, F. E. 1916. Plant succession. Carnegie Inst. Wash., Pub. 242.

- Drury, William H.; Nisbet, Ian C.T. 1973. Succession. Journal of the Arnold Arboretum. 54(3):331-368.
- Egler, F.E. 1954. Vegetation science concepts. I. Initial floristic composition-a factor in old-field vegetation development. Vegetatio. 4:412-417.
- Johnson, Paul S. 1989. Principles of natural regeneration. In: Clark, F. Bryan; Hutchinson, Jay G., eds. Central Hardwood Notes. St. Paul, MN: North Central Forest Experiment Station:3.01-1-5.
- Kelty, Matthew J. 1988. Sources of regeneration and factors that influence these sources. In: Smith, H. Clay; Perkey, Arlyn W.; Kidd, William E., Jr., eds. Guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV. Morgantown, WV: West Virginia University:17-30.
- Leopold, D.J.; Parker, G.R.; Swank, W.T. 1985. Forest development after successive clearcuts in the southern Appalachians. Forest Ecology and Management. 13:83-120.
- Noble, I.R.; Slatyer, R.O. 1980. The use of vital attributes to predict successional changes in plant communities subject to recurrent disturbances. Vegetatio. 43:5-21.
- Oliver, C.D. 1981. Forest Development in North America following major disturbances. Forest Ecology and Management. 3:153-168.

Shugart, H.H. 1984. A theory of forest dynamics. New York: Springer-Verlag. 278 pp. Page 200 of 404

Predicting Quantity and Quality of Reproduction in the Uplands

Dan Dey, Ontario Forest Research Institute, Sault Ste. Marie, Ont. P6A 5N5

The structure of regeneration models is determined by the sources of reproduction ABSTRACT and by the biotic and abiotic factors that affect population dynamics of the ecosystem being modeled. Sources of reproduction include new seedlings, advance reproduction, and sprouts from the cut stumps of overstory trees. The microenvironment (e.g., light, temperature, and soil moisture) influences the development of reproduction. Aspect, slope position, elevation, and site index are commonly used to express the integrated effect of micro-environment on reproduction. In some upland oak ecosystems, wildlife may adversely affect stand regeneration potential. Interference species such as ferns, fire cherry, red maple, and flowering dogwood may dominate the site following harvest, thereby reducing the regeneration potential. The importance of these factors in influencing regeneration potential varies among regions. Thus, regeneration models must be developed for specific regions. ACORn is an example of a regional stand-level regeneration model developed for even-aged forests in the Missouri Ozarks. It can be used to predict future stand characteristics by species and diameter classes, which in turn, can be used as input for growth and yield simulators.

INTRODUCTION Growth and yield models can simulate the development of even-age stands that are about 20 years of age and older. In the Central States, growth models such as TWIGS (Miner and others 1988), STEMS (Belcher and others 1982) and OAKSIM (Hilt, 1985) can be used to project stand changes resulting from real or hypothetical management practices. However, none of these models can predict stand structure or composition of the new stand. During the first two decades after final harvest, rapidly changing tree growth and competition relations determine the character of the mature stand.

The oak reproduction that develops after clearcutting depends on the oak regeneration potential of the parent stand. Regeneration potential refers to the capacity of a species or species-group to occupy and dominate growing space at a specified time in the new stand. For oaks, it can be determined from preharvest characteristics of the parent stand, including the advance reproduction and the overstory. To quantify the regeneration potential of a stand, the various sources of reproduction thus must be considered.

Natural reproduction after final harvest may come from new seedlings, advance reproduction, and sprouts from stumps of overstory trees (Beck 1980). New oak seedlings usually grow slowly and are not a major source of reproduction in most ecosystems (Sander and Clark 1971, Sander 1972, McQuilkin 1975). Viable acorns remain in the forest floor for 6 months or less because they are rapidly consumed by wildlife, damaged by insects and pathogens, and dessicate or freeze during the winter.

Advance reproduction and stump sprouts usually are the major sources of reproduction. The competitiveness of upland oak reproduction usually depends on the development before final harvest of a large root:shoot ratio. Oak advance reproduction can develop large root:shoot ratios through repeated shoot dieback and resprouting. Shoot dieback may result from water stress, weather damage (i.e., frost and freeze), insects, pathogens, fire, and browsing. Nevertheless, oak reproduction may accumulate and develop in the understory for decades (Merz and Boyce 1956, Tryon and others 1980). Once the overstory is removed, larger stems of advance reproduction can sprout vigorously and thus capture much of the newly available growing space (Carvell 1967; Sander 1971, 1972, 1979; McQuilkin 1975; Beck and Hooper 1986; Ross and others 1986; Johnson and Sander 1988; Loftis 1988). Thus, it is the number, size, and spatial distribution of oak advance reproduction that largely determines the oak regeneration potential of a stand.

Sprouts from cut stumps of overstory trees are usually the fastest growing form of oak reproduction (Spaeth 1928, Kuenzel 1935, Sander and Clark 1971, Smith 1979, Zahner and Myers 1984). The large food reserves and absorptive capacity of the parent root system support rapid shoot growth of stump sprouts. Stump sprouts make up variable portions of the total oak reproduction and often compensate for deficiencies in oak advance reproduction (Johnson and Sander 1988).

REGENERATION MODELS There are several hardwood regeneration models for predicting stand development after final harvest. Predictions are based on preharvest inventories of advance reproduction, the overstory (trees larger than 1.6 in. d.b.h.), and site factors. These models thus provide land managers the opportunity to evaluate alternative prescriptions before actual harvest. Methods for evaluating the natural regeneration potential of upland oak forests have been developed by Sander and others (1976, 1984), Johnson (1977), Johnson and Sander (1988), Lowell and others (1987), Waldrop and others (1986), Loftis (1988, 1990) and Marquis and Ernst (1988). Models for evaluating the potential contribution of planted trees to future stocking also have been developed for northern red oak (*Quercus rubra* L.) (Johnson 1988) and other hardwoods (Johnson 1984, Johnson and Rogers 1985).

Marquis and Ernst (1988) developed an expert system for Allegheny hardwood forests called SILVAH. Its application produces prescriptions based on management objectives, present stand conditions, and projections of tree growth and regeneration potential (Marquis and others 1992). To evaluate even-aged stands, critical stocking values are used to predict regeneration success. In applying the system, advance reproduction is inventoried on plots 6 ft. in radius. Whether a plot is stocked depends on species, numbers of stems, and their size. For example, a plot is considered stocked with oak if there are at least 25 stems less than 4.5 ft. tall, or if there is one stem greater than 4.5 ft $_{Ptall_{202}Oaks_4}$ in the

overstory also are inventoried. The number of reproduction plots expected to be stocked with oak stump sprouts at stand age 20 are estimated using models developed by Sander and others (1984). During the preharvest inventory, limiting site factors (poor drainage and stone content), amount of interference species competition, and intensity of deer browsing also are noted. Because these factors reduce the regeneration potential of a stand, the number of plots deemed stocked with desirable reproduction are accordingly reduced. The number of plots stocked with desirable tree species thus determine whether or not the regeneration potential is adequate. In general, when 70 percent of the plots are stocked with desirable advance reproduction, successful regeneration is expected to occur.

Loftis (1990) developed a regeneration model for mixed hardwood forests of Southern Appalachia. The model projects eighth-year post-harvest heights and crown classes of northern red oak advance reproduction based on preharvest measurements of advance reproduction and site quality. Heights of red oak reproduction 8 years after final overstory removal are estimated from preharvest height and basal diameter of advance reproduction and from site index. Preharvest basal diameters and site index are then used to predict the probability that a given stem of oak advance reproduction grows to dominant or codominant crown classes in the new stand at age 8. This model thus can be used to evaluate the contribution of red oak reproduction to the future stand.

Sander and others (1984) developed a regeneration model for the Missouri Ozarks. It predicts the probability of individual oaks surviving and attaining a specific future height that places them in codominant or dominant crown classes. From preharvest inventories of the overstory and advance reproduction, the model predicts the adequacy of future oak stocking. A stand is deemed to have adequate regeneration potential if projected stocking of dominant and codominant oaks at stand age 20 is at C-level or greater based on Gingrich's (1967) stocking relations. However, the model is unable to predict the diameter distribution of oaks in the new stand, and non-oaks are not considered. Although species such as blackgum (*Nyssa sylvatica* Marsh.), sassafras (*Sassafras albidum* (Nutt.) Nees, hickory (*Carya* spp.), and flowering dogwood (*Cornus florida* L.) seldom occur as dominant trees in Ozark forests once they reach 20 years of age, they do affect the development of stands during the first two decades after final harvest because of their high density and rapid growth. These species are also important to wildlife and biodiversity.

In contrast to some other deciduous forests of the Eastern United States, the forests of the Missouri Ozarks regenerate primarily from sprouts of harvested overstory trees and advance reproduction. Therefore, the composition and size structure of the future stand is largely determined by the species composition and size structure of the preharvest advance reproduction and overstory. By observing these features together with measurements of site factors such as slope, aspect, and site index, model users can predict the composition and structure of the new stand.

A COMPREHENSIVE OZARK REGENERATOR Dey (1991) developed a regeneration model called <u>A</u> Comprehensive <u>O</u>zark <u>Regen</u>erator (ACORn) to simulate the regeneration of even-aged stands in the Missouri Ozarks. It was developed from measurements of individual stems of reproduction before and after clearcutting. ACORn compresses two_4modules that

simulate the development of the two primary sources of reproduction in that ecosystem: stump sprouts and advance reproduction. One module simulates the development of advance reproduction and the other the development of stump sprouts. Each module, in turn, contains models for estimating future heights, diameters, and survival of individual stems by species. The probability of survival of individual stems of both advance reproduction and stump sprouts, from preharvest to a specified future stand age, is estimated from initial tree size and site factors. Species-specific models estimate future tree heights and diameters from similar predictors. The predictive models, in turn, facilitate the generation of future diameter and height distributions of surviving reproduction including stump sprouts and advance reproduction by species. Projections for hickory, flowering dogwood, blackgum, and sassafras also can be obtained.

In application, preharvest inventories of the overstory and advance reproduction provide input to survival and growth models that generate diameter distributions 21 years after clearcutting. At that age, mean stand diameter averages 3 in. d.b.h. The diameter distributions then can be used to summarize stand characteristics, such as basal area per acre, stems per acre, and percent stocking by species. From this stand summary, the adequacy of future stocking can be assessed and used to develop appropriate silvicultural prescriptions. The resulting diameter distributions also can be used as input into growth and yield simulators such as TWIGS.

PREDICTING REGENERATION QUALITY

Quality of reproduction is commonly defined in terms of the acceptability of growing stock. Before adequacy of regeneration potential can be assessed, it is necessary to define acceptable growing stock, which requires a consideration of species and tree characteristics. Commonly used tree characteristics are height, diameter, and crown class. Although a qualitative characteristic, crown class is a concept with which most foresters are familiar. It is a widely used descriptor of trees because it facilitates visualization of a tree's social status more easily than diameter or height measurements, *per se*. For example, the definition of acceptable growing stock may be limited to certain species and trees that occupy only codominant and dominant crown classes. ACORn solves that problem by projecting future diameter and survival of individual trees by diameter and crown classes to stand age 21.

Some foresters may consider only codominant and dominant trees as acceptable growing stock. Others may consider trees that are intermediate or larger as acceptable. To integrate this decision into the regeneration model, a threshold tree diameter is defined that classifies trees into one of the two user-defined crown class categories: acceptable or unacceptable. When acceptable growing stock includes only codominant and dominant trees, tree diameters equal to or greater than 3.8 in. are classified as codominant or dominant at stand age 21. When acceptable growing stock is defined as trees that are intermediate or larger, trees with diameters less than 2.6 in. are classified as suppressed, and those with larger diameters are classed as acceptable growing stock. In this way, ACORn incorporates crown class to project twenty-first-year diameter distributions.

Regeneration Surveys Data from preharvest inventories of the overstory and advance reproduction provide the necessary input for ACORn projections of future stand composition and structure. To inventory a stand, the overstory (trees 1.6 in. d.b.h. and larger) is Page 204 of 404

sampled separately from the advance reproduction (trees less than 1.6 in. d.b.h.). Because ACORn is an individual tree model, it projects the survival and growth of single trees and then expands this to a per-acre basis in the form of stand tables. The model thus can accommodate a variety of sampling designs.

Fixed-area or variable-radius plots can be used to inventory the overstory. Sander and others (1984) recommended that the overstory be sampled on 1/20-acre plots in their regeneration guide. Small fixed-area plots were recommended for inventory of advance reproduction. The model of Sander and others (1984) requires that advance reproduction be inventoried on 1/735-acre plots. The choice of sample plot size and number should be done on a stand-by-stand basis. There are numerous sampling techniques that can be used to design efficient and effective inventories (e.g., Freese 1962).

In general, the more variation there is in species composition, tree size, and the spatial distribution of trees, the more sample plots that are needed. It is usually best to sample a large number of small plots where stand variation is great than it is to sample fewer large plots. A common rule-of-thumb is to inventory at least 30 plots regardless of stand size. The upper limit to the number of plots depends on (1) constraints such as budget, crew size, and timeframe; (2) intended use of simulation, i.e., forest planning or stand prescription development; (3) amount of stand variation; and (4) desired level of precision. Plots should be distributed randomly or systematically throughout the stand.

ACORn requires an inventory of the overstory including species, d.b.h., site index (black oak, base age 50), and stand age. When the overstory is even-aged, an average stand age can be substituted for individual tree age. For advance reproduction, the required inventory includes data on species, basal diameters, and heights of reproduction, slope position (upper, middle, or lower), and aspect. A computer program is available to facilitate application of the model.

SUMMARY The structure of regeneration models is determined by the sources of reproduction and by the biotic and abiotic factors that affect population dynamics of the ecosystem being modeled. Sources of reproduction include new seedlings, advance reproduction, and sprouts from the cut stumps of overstory trees. The microenvironment (e.g., light, temperature, and soil moisture) influences the development of reproduction. Aspect, slope position, elevation, and site index are commonly used to express the integrated effect of micro-environment on reproduction. In some upland oak ecosystems, wildlife such as deer may adversely affect stand regeneration potential. Interference species such as ferns, fire cherry, red maple, and flowering dogwood may dominate the site following harvest, thereby reducing the regeneration potential. The importance of these factors in influencing regeneration potential varies among regions. Thus, regeneration models should be developed for specific regions.

> Hardwood regeneration models are available today for individual species or forest types in the United States. Several examples have been discussed in this paper. SILVAH incorporates predictions of regeneration success of northern hardwoods in selection of appropriate silvicultural prescriptions. A regeneration model also is available for northern red oak in the Southern Appalachians. ACORn generates Page 205 of 404

future diameter and height distributions of reproduction (stump sprouts and advance reproduction) for the Missouri Ozarks. The resulting distributions simulate, by species and reproductive origin, the regeneration process from preharvest to the end of the regeneration period (stand age 21). Characteristics of the new stand then are summarized in the output so they also can be used as input into growth and yield simulators such as TWIGS. Resource managers thus can predict, before harvest, future stand development following final harvest. Preharvest inventory of the parent stand and site conditions is a necessary first step to planning appropriate silvicultural activities.

LITERATURE CITED Beck, Donald E. 1980. Natural regeneration of southern hardwoods. In: McElwee, R.L.; Bell, N.H., eds. Regenerating the Southern forests. Tenth Forestry and Wildlife Forum; 1980 April 17-18; Blacksburg, VA: Virginia Polytechnic Institute and State University: 132-141.

- Beck, Donald E.; Hooper, Ralph M. 1986. Development of a southern Appalachian hardwood stand after clearcutting. Southern Journal of Applied Forestry. 10:168-172.
- Belcher, David W.; Holdaway, Margaret R.; Brand, Gary J. 1982. A description of STEMS the stand and tree evaluation and modeling system. Gen. Tech. Rep. NC-79. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 18 pp.
- Carvell, Kenneth L. 1967. The response of understory oak seedlings to release after partial cuttings. Bull. 553. Morgantown, WV: West Virginia University, Agriculture Experiment Station. 20 pp.
- Dey, Daniel C. 1991. A comprehensive Ozark regenerator. Columbia, MO: University of Missouri. 283 pp. Dissertation.
- Freese, Frank. 1962. Elementary forest sampling. Agric. Handb. 232. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 91 pp.
- Gingrich, Samuel F. 1967. Measuring and evaluating stocking and stand density in upland hardwood forests in the Central States. Forest Science. 33(1):38-53.
- Hilt, Donald E. 1985. User's guide to OAKSIM an individual-tree growth and yield simulator for managed, even-aged, upland oak stands. Gen. Tech. Rep. NE-104. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 22 pp.
- Johnson, Paul S. 1977. Predicting oak stump sprouting and sprout development in the Missouri Ozarks. Res. Pap. NC-149. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 11 pp.
- Johnson, Paul S. 1984. Responses of planted northern red oak to three overstory treatments. Canadian Journal of Forest Research. 14:536-542.
- Johnson, Paul S. 1988. Undercutting alters root morphology and improves field performance of northern red oak. In: Worrall, John; Loo-Dinkens, Judy; Lester, Donald P., comp. and eds. Proceedings of the Tenth North American Forest Biology Workshop; 1988 July 20-22; Vancouver, BC: University of British Columbia: 316-323
- Johnson, Paul S.; Rogers, Robert. 1985. A method for estimating the contribution of planted hardwoods to future stocking. Forest Science. 31(4):883-891.
- Johnson, Paul S.; Sander, Ivan L. 1988. Quantifying regeneration potentials of <u>Quercus</u> forests in the Missouri Ozarks. In: Ek, A.R.; Shifley, S.R.; Burk, T.E., eds. Forest growth modelling and prediction, Proceedings, mensuration, growth and yield, subject group S4.01,IUFRO; 1987 August 24-28; Minneapolis, MN. Gen. Tech. Rep. NC-120. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 377-385.

- Kuenzel, John G. 1935. The influence of degree of cutting on the survival and early growth of chestnut oak seedlings. Station Note 23. U.S. Department of Agriculture, Forest Service, Central States Forest Experiment Station. 3 pp.
- Loftis, David L. 1988. Regenerating red oak in the southern Appalachians: predictive models and practical applications. North Carolina State University. 65 pp. Dissertation.
- Loftis, David L. 1990. Predicting post-harvest performance of advance red oak reproduction in the southern Appalachians. Forest Science. 36(4):908-916.
- Lowell, K.E.; Mitchell, R.J.; Johnson, P.S. [and others]. 1987. Predicting growth and "success" of coppice-regenerated oak stems. Forest Science. 33(3):740-749.
- Marquis, David A.; Ernst, Richard L. 1988. User's guide to SILVAH a stand analysis, prescription and management simulator program for hardwood stands of the Alleghenies. Warren, PA: U.S. Department of Agriculture, Forest Service, Silviculture Research Unit, Northeastern Forest Experiment Station. 161 pp.
- Marquis, David A.; Ernst, Richard L.; Stout, Susan L. 1992. Prescribing silvicultural treatments in hardwood stands of the Alleghenies (revised). Gen. Tech. Rep. NE-96. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 101 pp.
- McQuilken, Robert A. 1975. Growth of four types of white oak reproduction after clearcutting in the Missouri Ozarks. Res. Pap. NC-116. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 5 pp.

Merz, Robert W.; Boyce, Stephen G. 1956. Age of oak "seedlings". Journal of Forestry. 54:774-775.

- Miner, Cynthia L.; Walters, Nancy R.; Belli, Monique L. 1988. A guide to the TWIGS program for the North Central United States. Gen. Tech. Rep. NC-125. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 105 pp.
- Ross, Michael S.; Sharik, Terry L.; Smith, David W. 1986. Oak regeneration after clearfelling in southwest Virginia. Forest Science. 32(1):157-169.
- Sander, Ivan L. 1971. Height growth of new oak sprouts depends on size of advance reproduction. Journal of Forestry. 69:809-811.
- Sander, Ivan L. 1972. Size of oak advance reproduction:key to growth following harvest cutting. Res. Pap. NC-79. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 6 pp.
- Sander, Ivan L. 1979. Regenerating oaks with the shelterwood system. In: Holt, H.A.; Fischer, B.C., eds. Proceedings, regenerating oaks in upland hardwood forests. The 1979 J.S. Wright Forestry Conference; 1979 February 22-23; West Lafayette, IN: Purdue University: 54-60.
- Sander, Ivan L.; Clark, F. Bryan. 1971. Reproduction of upland hardwood forests in the Central States. Agric. Handb. 405. Washington, DC: U.S. Department of Agriculture. 25 pp.
- Sander, Ivan L.; Johnson, Paul S.; Rogers, Robert. 1984. Evaluating oak advance reproduction in the Missouri Ozarks. Res. Pap. NC-251. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 16 pp.
- Sander, Ivan L.; Johnson, Paul S.; Watt, Richard F. 1976. A guide for evaluating the adequacy of oak advance reproduction. Gen. Tech. Rep. NC-23. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 7 pp.
- Smith, H. Clay. 1979. Natural regeneration and intensive cultural practices in central Appalachian hardwood stands using clearcutting and selection cutting practices. In: Holt, H.A.; Fishcer, B.C., eds. Proceedings, regenerating oaks in upland hardwood forests. The 1979 J.S. Wright Forestry Conference; 1979 February 22-23; West Lafayette, IN: Purdue University: 30-41. Page 207 of 404

- Spaeth, J. Nelson. 1928. Twenty years growth of a sprout hardwood forest in New York: a study of the effects of intermediate and reproduction cuttings. Bull. 465. Ithaca, NY: Cornell University, Agriculture Experiment Station. 49 pp.
- Tryon, Earl H.; Powell, Douglas S.; Harner, E. James. 1980. Predicting root age of advance growth beneath hardwood canopies. Canadian Journal of Forest Research. 10:264-268.
- Waldrop, T.A.; Buckner, E.R.; Shugart, Jr, H.H.; McGee, C.E. 1986. FORCAT: a single tree model of stand development following clearcutting on the Cumberland Plateau. Forest Science. 32(2):297-317.
- Zahner, Robert; Myers, Richard K. 1984. Productivity of young piedmont oak stands of sprout origin. Southern Journal of Applied Forestry. 8(2):102-108.

Prediction of Oak Regeneration in Bottomland Forests

Robert L. Johnson, USDA Forest Service, Retired, Leland, MS 38756 Robert T. Deen, School of Forest Resources, Mississippi State University, Mississippi State, MS 39762

ABSTRACT

IN 1980 Johnson presented a preliminary method to evaluate oak regeneration potential in southern bottomland hardwood forests. The method evaluates regeneration potential by assigning points based on number and size (height class) of advance regeneration and sprout potential of stumps from severed trees. The method appears promising for evaluating regeneration potential but has not been tested under operational conditions. Research is currently underway to test the method and make refinements if necessary. Current data indicate that 78 percent of plots evaluated as adequately stocked before harvest were still adequately stocked with desirable regeneration following harvest. Preliminary results showing seedling mortality, seedling growth by size (height and root collar), and logging losses indicate that improvements to the method can be made through modifications of point assignment.

INTRODUCTION Through continuing and widespread research efforts, there now exist predictive models to evaluate the potential for oak regeneration success prior to harvest of upland hardwood stands (Dey 1991, Loftis 1990, Marquis and Bjorkbom 1982, Sander and others 1976, Waldrop and others 1986). The foundation underlying the use of these evaluation techniques is that, to successfully regenerate oaks, there must exist adequate numbers and/or size of advance oak regeneration prior to harvest.

The above mentioned predictive models (Dey 1991, Loftis 1990, Marquis and Bjorkbom 1982, Sander and others 1976, Waldrop and others 1986) were designed for use within specific physiographic regions and application elsewhere should be approached with caution. A more complete description of these regeneration techniques will be given by their respective authors elsewhere in these Proceedings.

The harvest and regeneration of bottomland hardwoods has been a fascinating subject for many years. Hardwood regeneration in the bottoms is often easy, but almost as often, the successful regeneration of bottomland hardwoods, especially oaks, can be very difficult. Forest land managers have for many years needed to predict when, where, and how their hardwood regeneration efforts would be successful. In 1980, a hardwood regeneration prediction model for bottomland hardwoods was devised (Johnson 1980). The purpose of this paper is to examine how the Johnson method can be used to predict oak regeneration, and based upon preliminary testing, how the procedure can be improved.

The regeneration prediction method when used within the constraints discussed later has been found to be both useful and reliable. However, we should emphasize that the expected results for any given situation are based upon a combination of empirical results and the senior author's experience. The authors believe that with on-the-ground experience many users can adapt the procedure to more closely fit their needs and conditions.

For bottomland oaks, our target species in this paper (table 1), we know, for example, that acorn production varies considerably among species, years, and phenotypes. Good acorn crops may be 3 to 5 years apart. Acorns are often utilized by wildlife after seedfall and a good seed crop may not produce a good seedling crop. In some years and on some sites acorns of bottomland oaks may be protected from birds and animals when they drop into or are soon covered by water that stays in the forest until early in the growing season. Duration of the water cover is important. If the water leaves before germination, acorns will be exposed and eaten; if the water stays into the summer, acorns will rot. Water that recedes during the germination period usually results in several days of moist seedbed that allows germinating acorns to establish seedlings. Without surface water and given that radicles cannot penetrate a dry, hard surface soil, acorns that roll into soil cracks or that are buried by animals or birds (Deen and Hodges 1991) have the best chance to establish a new seedling.

Oak seedling populations may be large in numbers beneath the crowns of mature oak trees; e.g., Johnson (1975) found up to 100,000+ nuttall oak (*Quercus michauxii* Nutt.) seedlings per acre in the Mississippi Delta. Red oak seedlings in full shade will normally grow approximately a foot in height during the first year, and most die within 3 years. Those that receive 2 hours of direct sunlight daily may survive beyond 3-5 years, but grow only an inch or so in height annually.

Common name of species	Number of stems	Percentage of total		
Willow oak	1,311	45.30		
Water oak	799	27.60		
Green ash	197	6.80		
Cherrybark oak	193	6.70		
Overcup oak	168	5.80		
White oak	108	3.70		
Diamondleaf oak	47	1.60		
Swamp Chestnut oak	31	1.10		
Shumard oak	20	.70		
Southern Red oak	12	.40		
Nuttall oak	5	.20		
Northern Red oak	1	.03		
Total	2,892	99.93		

Table 1—Ranking of desirable species; i.e., oaks and ash, based on their percentage of the total number of stems initially sampled over all study plots in east central and south Mississippi

Oaks that die back and resprout almost always start in small openings where there is enough sunlight to allow for 5-10 ft. of height growth before the overstory closes and direct sunlight is shut off. These larger trees have a well-developed root system that will support rapid shoot growth when the overstory is removed.

Once released, mortality of oak seedlings is most likely to occur the first year. Flooding is a likely cause of mortality, but exposure to frost may result in dieback or mortality (McGee 1988). Thereafter, at least through year 10, three of four oaks should survive even with overtopping competition. With only side competition, oak regeneration will continue growing upward and have a good chance to eventually become dominant.

Height growth of released seedlings ≤ 1 ft. tall is slow for the first 5 years, whereas taller advanced reproduction and stump sprouts grow rapidly from the start. For bottomland oaks, stumps up to 12 in. in diameter have a good chance of producing sprouts that are acceptable regeneration; larger stumps usually do not produce sprouts, or if they do, the sprouts are likely to die. Because of the length of time necessary to produce quality oak lumber, oak stems will grow beyond a threshold-diameter capable of producing many, if any, stump sprouts. Also, the total number of sawlog-sized oak stems per acre in a mature, bottomland hardwood stand would be low enough to discount these stumps as providing adequate stocking of oak sprouts for regeneration purposes. Sawlog-sized oak, then, can be mostly discounted as an adequate source of sprouts from which to restock the newly developing stand (Johnson 1975).

THE PREDICTION MODEL

The method proposed by Johnson (1980) is based on the silvical characteristics of the species as discussed above. It is a numerical evaluation of regeneration potential which emphasizes the size (height class) and numbers of advance regeneration and the contribution of stump sprouts to regeneration potential (form 1). Weights and points given various regeneration components are based on the senior author's 35 plus years of research experience. The following procedures are used in the evaluation process:

1. The sample plot is a circular 1/100th acre. This plot size will accommodate large trees (because stump and root sprouts are considered as reproduction) as well as seedlings and saplings. The selected size also provides an easy blow-up factor.

2. One point for each tree ≤ 1.0 ft. tall. Mortality is high among such trees because they are only a year or two old and are not well established. In fact, all trees ≤ 1.0 ft. tall may die if overstory removal is delayed for more than a year. However, a few small trees may be older, well established, and likely to survive several years with or without overstory release. The one point per tree is an attempt to balance the survival potential of young and older, established trees. Young or old, small trees are not very competitive.

3. Two points for each tree 1.1 to 2.9 ft. tall. Trees of this size are older than 1 year and have a well-developed root system that will aid survival and provide for height growth. Trees in this size class are more competitive than trees ≤ 1.0 ft. tall, but less competitive than those ≥ 3.0 ft. tall.

	Date	
	Plot number	
Form 1	Location	
REPRODUCTION INVENTORY		

Circular 1/100-acre plot (11.8' radius)

Points per tree and number of trees by species and size class

	Height (feet)			DBH (inches)				
	<1.0	1.0-2.9	3.0+	2-5	6-10	11-15	16-20	To
Species	Pts. No.	Pts.No.	Pts. No.	Pts.No.	Pts.No.	Pts.No.	Pts. No.	Pt
Ash	1	2	3	3	3	2	1	-
Red oaks								
	1	2	3	3	2	1	0	_
	1	2	3	3	2	1	0	
	1	2	3	3	2	1	0	
White oaks								-
	1	2	3	3	2	1	0	
	1	2	3	3	2	1	0	-
Hickory	1	2	3	3	3	2	1	
Sweetgum	1	2	3	3	3	2	1	
Blackgum	1	2	3.	3	3	2	1	
Elm	1	2	3	3	2	1	0	-
Yellow-poplar	1	2	3	3	2	1	0	
Maple	1	2	3	3	2	2	0	
Sugarberry	1	2	3	3	2	1	0	
Persimmon	1	2	3	3	2	1	0	
Other								
	1	2	3	3	2	1	0	
	1	2	3	3	2	1	0	
Stocked - 12 po:	ints or mo	re		Yellow-po within 10	plar or s	ycamore s	eed tree	
Total points th				within 10	0 feet	-		+6
Pointeo un				Seedbed b	are			+2
				Seedbed w	eedv			-2

4. Three points for each tree ≥ 3.0 ft. tall but less than 5.5 in. d.b.h. Trees are able to grow into this size class while in small openings but not in full shade. Thus, the trees are several years old, have large root systems, are almost certain to survive an overstory harvest even if their tops are broken off, are good stump sprouters, and should have rapid post-harvest height growth. Trees in this class make good competitors.

5. Two points for each tree 5.6 to 10.5 in. d.b.h. A high percentage of stumps from trees in this size class will produce competitive sprouts that survive and grow very well. Usually no more than three stumps of this size will occur in a 1/100thacre plot.

6. One point for each tree 10.6 to 15.5 in. d.b.h. Not many of these relatively large stumps will produce acceptable, competitive sprouts. No more than one or two trees of this size class will occur in a 1/100th-acre plot.

7. Twelve points per plot is considered high enough to ensure adequate regeneration following harvest. The points could represent 12 trees per plot or 1,200 trees per acre that are a foot or less in height, 4 trees per plot (400 per acre) 3.0 ft. tall to 5.4 in. d.b.h., or from any combination of trees equalling 12 points. Although natural stands may start with 20,000 plus trees per acre, they will thin to fewer than 400 crop trees per acre at minimum commercial size (6 in. d.b.h.). The 12 points is a judgment call, believed conservative, and should be considered only as a general guide.

A possible limitation of the guide is that it does not recognize seedlings which germinate after harvest. Oak seedlings may become established after harvest and become an important part of the regenerated stand, but, this model does not provide for such a happening. If advance regeneration is lacking and if there are no oak stumps of sprout-producing size, this model predicts zero oaks in the regeneration stand. Also, the total number of stocked plots needed for successful regeneration of a given tract is still questionable. It appears that at least 60 percent of the total plots must be stocked (\geq 12 points) and well distributed over the tract to ensure successful regeneration.

Testing the Prediction Model

Research is currently underway to test and evaluate the regeneration prediction method described herein.

Study areas 10 acres and larger were located mainly in the floodplain of minor streambottoms. Permanent 1/100th-acre circular plots were systematically located within each study area.

Woody vegetation was placed into one of three size classes and identified as either desirable (oaks and ash) or undesirable (all other woody species).

To develop a priority rating for predicting regeneration potential of oaks and ash, desirable stems placed in seedling or sapling categories were identified and tagged. Stems could be identified from year-to-year, enabling development patterns to be detected and related to preharvest size.

Desirable and undesirable vegetation was identified by species and placed in the appropriate height class. In addition, permanent trees were selected and measured for root collar diameter. All woody vegetation was designated as to origin class; i.e., sprout or seedlings. Desirable and permanent sample trees were noted as to occurrence of top die-back, insect or disease damage, damage due to harvesting operation, and subjectively rated as to competitive status, i.e., free-to-grow, medium competition from surrounding vegetation, or overtopped.

Regeneration was evaluated for each plot according to Johnson's (1980) technique (form 1) before harvest and after each remeasurement to determine the adequacy

of the technique and, if needed, make adjustments to the guide to more accurately predict regeneration success.

Following harvest, a resurvey of the plots was conducted to quantify logging damage, if any, to permanent sample trees and to the site that would adversely affect subsequent vegetative composition and growth and development of the future stand. At this time, a survey was taken of the number of light-seeded species occurring within 200 ft. of each plot. Seedlings from these fast-growing, intolerant species could ultimately suppress the development of desirable species, oaks and ash, in the future stand.

Other factors affecting stand development (i.e., high water, severe vegetative competition, etc.) were recorded at each plot visit. Also, newly germinating oak or ash species were marked with a combination of colored expansion rings, designating a particular height class and a particular year of germination, so further development can be followed.

RESULTS AND The CONCLUSIONS cen

There were 118 plots established on nine bottomland hardwood tracts in east central and south Mississippi. The tracts used in the study were logged between August and September of 1989 and 1990. Table 1 gives the total number of desirable stems, by species, that were initially sampled in this study.

Based on form 1, the sampled stands shown in table 2 exhibit differing trends of stocking success following harvest. The decline in percentage stocking of those stands shown in table 2 can be attributed to the high degree of mortality, due to logging, of oak regeneration ≤ 1.0 ft. tall (table 3). Seedlings ≤ 1.0 ft. tall accounted for the bulk of points assigned to plots within these stands before

	Percent plot	Years Since		
'ract	Preharvest	Present	Harves	
Noxubee Refuge				
U1DLR	55.6	33.3	2	
U3KTR	60.0	40.0	2	
U2TPR	68.8	68.8	2	
U1KTR	50.0	50.0	2	
U3CGR	60.0	40.0	3	
U1CGR	100.0	100.0	3	
U4KTR	55.5	77.8	3	
Scott Paper				
#1	.0	.0	1	
#2	80.0	30.0	1	

Table 2—The use of Johnson's (1980) evaluation technique (form 1) to predict percentage of plots adequately stocked, with oaks and ash, per tract before and after harvesting

152 Oak Regeneration Symposium Proceedings . . . September 1992

	Seedling height classes (ft.)				
******	0.1-1.0	1.0-2.9	<u>></u> 3.0		
Percent	62.9	40 .1	41.1		
(# Dead/Total #)	(166/264)	(78/192)	(69/164)		

Table 3—Percent mortality among permanent sample seedlings of desirable species, i.e., oaks and ash, due to logging damage

harvest, and the loss of these seedlings following harvest resulted in lower point totals for these plots. Accordingly, form 1 may need to be revised to reflect the need for greater numbers of regeneration ≤ 1.0 ft. tall for adequate regeneration when this is the only source of regeneration.

The one stand (table 2) that showed an increase in stocking percentage following harvest contained a majority of desirable stems > 1.0 ft. tall before harvest. Desirable stems, 1.1-2.9 ft. tall and \geq 3.0 ft. tall, have a much greater chance for survival (table 4). Also, in this stand individual stems have progressed in height class, increasing the points allocated to that particular seedling, and likewise increasing the point total for that plot.

Years after	Seedling height classes (ft.)		
harvest	0.1-1.0	1.0-2.9	<u>>3.0</u>
1	25.0	54.4	63.7
2	9.8	46.5	63.7

Table 4—Percent survival rates of seedlings of desirable species, i.e., oaks and ash, by height classes 1 and 2 years after harvest

Four stands remained constant as far as percentage stocking is concerned. Two stands appear adequately stocked, one stand is borderline, and one stand is considered nonstocked.

One point we are trying to determine is the cut-off point where a stand should be classified as suitable or not for regeneration based on initial stocking. Twelve points is now considered the threshold. However, some plots that do not make 12 points contain several stems that are large enough to successfully compete for growing space and some day become a component of the overstory. If that is the case plots containing less than 12 points but containing larger regeneration may be adequately stocked. Observation of stand development over time is needed to clarify this point.

Logging damage following harvest was assessed using the permanent sample trees located within each plot. Saplings and/or seedlings were categorized as to the extent of logging damage, i.e., intact, bent, broken, missing, or dead. Stems that were classified with severe logging damage (i.e., broken, missing, or dead) which were dead at the next remeasurement were assumed to have died as a result of logging damage (table 3). Some stems in the sapling class, 2.0-4.4 in. d.b.h., were pushed completely out of the ground exposing their rootstocks and ultimately resulting in death. Loss of this type of regeneration can be reduced by chainsawing of saplings prior to harvest.

Development of oak regeneration after clearcutting followed similar patterns to those reported in other regions (Loftis 1983, Sander 1972) where large advance regeneration exhibited better survival than small advance regeneration. Survival data (table 3) indicate that desirable seedlings ≤ 1.0 ft. in height cannot be counted on to provide many stems to the newly regenerated stand. Also, field data and observation showed that these seedlings ≤ 1.0 ft. tall which did survive had large root collar diameters, indicating that such stems had probably died back and resprouted several times. The larger rootstocks enabled these smaller stems to produce a vigorous, competitive sprout.

Stems in the 1.1-2.9 ft. and \geq 3.0 ft. height classes exhibited much better survival rates following harvest (table 3). It appears these two size classes will contribute the bulk of the stems to the new regeneration stand.

One growing season after harvest, 162 new oak and ash germinants, ≤ 1.0 ft. tall, and seven new ash germinants, 1.1-2.9 ft. tall, were found. Forty percent of the seedlings ≤ 1.0 ft. tall survived their first growing season while six of the seven ash germinants 1.1-2.9 ft. tall survived. In the second year following harvest there were 22 new germinants of oak and ash which were ≤ 1.0 ft. tall, and six new ash seedlings, 1.1-2.9 ft. tall. They have yet to be evaluated after a full growing season so their fate is undetermined. However, most research indicates that contribution of the new seedlings will be insignificant and that advance regeneration will provide the bulk of the stems to the new stand.

Seedling height growth was as follows: seedlings ≤ 1.0 ft. tall averaged 7.7 in. tall preharvest. One year after harvest, average height for these seedlings was 10.9 in., and after 2 years average height was 17.7 in. Seedlings 1.1-2.9 ft. tall, similarly, realized a 10.0-in. increase in average height over 2 years, growing from 20.6 to 30.6 in. However, the desirable stems in the ≤ 1.0 ft. and 1.1-2.9 ft. height classes are competing with desirable stems from the ≥ 3.0 ft. height class, as well as undesirable stems, for growing space in the newly developing stands. A number of stems in these height classes are free-to-grow and may eventually achieve a dominant position.

Seedlings ≥ 3.0 ft. tall before harvest showed a decrease in average height after the first growing season, from 82.2 in. to 47.3 in. This was primarily a result of logging damage, but there is evidence that exposure to low temperatures in the spring following release can result in top die-back (McGee 1988). After the second year, average height increased to 60.1 in. This is a result, mainly, of the vigorous sprout growth of these larger stems of advance regeneration. Some stems were also able to respond rapidly to the increased light conditions created by clearcutting. A majority of these stems are already in a free-to-grow position and have assumed a dominant position in the new stand. Their vigorous, rapid height growth may assure a dominant crown position as the stands develop.

Preliminary data show that stump sprouting for trees >1.5 in. d.b.h. were as follows: 45.6 percent for the 2.0-5.0 in. d.b.h. class, 42.9 percent for the 6.0-10.0 in. d.b.h. class, 33.3 percent for the 11.0-15.0 in. d.b.h. class, 50.0 percent for the 16.0-20.0 in. d.b.h. class (only two trees in this class), and the >20.0 in. d.b.h. class had no sprouting. The trend seems to be a decrease in sprouting with an increase in tree (stump) size as shown in other studies (Johnson 1977, McQuilkin 1975). Also, as trees increase in size their numbers decrease on a per-acre basis and, coupled with their reduced sprouting ability, further reduces their contribution to the regeneration stand.

RECOMMENDATIONS A model that predicts regeneration success was developed for southern bottomland hardwood forests, but should be suitable throughout the Southeastern United States. It is applicable only where complete overstory harvest is intended, meaning that all trees 2 in. d.b.h. and larger are cut from an area sufficient in size to allow full sunlight to reach the ground.

Because the number and vigor of understory oaks change rapidly, prediction by the model is useful for 1 year only. If harvesting is delayed, the stand should be re-sampled.

Based on experience it appears that one sample plot per acre might be needed for tracts under 50 acres, whereas for larger areas one plot per 2 or 3 acres may do. Stratified sampling is suggested where different stand conditions can be recognized and delineated. A systematic sampling scheme is practical and adequate.

The model is not refined enough to predict which tree, among the many mixed species, may eventually dominate. But, in the same growth environment (same growing space and micro-site) a tree with three points is much more likely to survive and compete than a 1-point tree. Growth environments differ, however, even over a foot or two in space, so there may be considerable variation in the way one 3-point tree compares to three 1-point trees in long-term development among competitors.

A point total of less than 12 per plot doesn't necessarily mean the sample area will not regenerate satisfactorily, but it does signal caution. For example, it seems likely that if oaks contribute only two points per plot, then they will not be a major part of the new stand. On the other hand, large advance regeneration appears highly likely to survive the rigors of harvesting to become a component of the new stand. Possibly, only two large stems per plot may be necessary to ensure successful regeneration.

A plot with more than 12 points, made up mostly of seedlings ≤ 1 ft. in height, may not be adequate for regeneration due to heavy mortality among seedlings in this height class. More seedlings ≤ 1 ft. in height may be needed to qualify plots as adequately stocked.

LITERATURE CITED

Deen, Robert T.; Hodges John D. 1991. Oak regeneration in abandoned fields: presumed role of the blue jay. In: Coleman, Sandra S.; Neary, Daniel G., Comp. and eds. Proceedings of the Sixth Biennial Southern Silvicultural Conference; 1990. October 30 - November 1; Memphis, Tn: Gen. Tech. Rep. SE-70. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 84-93.

- Dey, Daniel C. 1991. A comprehensive Ozark regenerator. Columbia, MO: University of Missouri-Columbia. 283 pp. Dissertation.
- Johnson, Paul S. 1977. Predicting oak stump sprouting and sprout development in the Missouri Ozarks. Res. Pap. NC-149. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 11 pp.
- Johnson, R. L. 1975. Natural regeneration and development of nuttall oak and associated species. Res. Pap. SO-104. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 12 pp.
- Johnson, R. L. 1980. New ideas about regeneration of hardwoods. In: Proceedings, Hardwood Regeneration Symposium; 1980. January 29; Atlanta, Ga. Southeastern Lumber Manufacturing Association; Forest Park, GA. 17-19.
- Loftis, David L. 1990. Predicting post-harvest performance of advance red oak reproduction in the Southern Appalachians. Forest Science. 36(4): 908-916.
- Marquis, David A.; Bjorkborn, John C. 1982. Guidelines for evaluating regeneration before and after clearcutting Allegheny hardwoods. Res. Note NE-307. Warren, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 4 pp.
- McGee, Charles E. 1988. Spring weather, canopy removal, and early budbreak threaten oak seedlings. The Journal of the Elisha Mitchell Scientific Society. 104(3):108-115.
- McQuilkin, Robert A. 1975. Growth of four types of white oak reproduction after clearcutting in the Missouri Ozarks. Res. Pap. NC-116. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 5 pp.
- Sander, Ivan L.; Johnson, Paul S.; Watt, Richard F. 1976. A guide for evaluating the adequacy of oak advance reproduction. Gen. Tech. Rep. NC-23. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 7 pp.
- Waldrop, T. A.; Buckner, E. R.; Shugart, H. H., Jr.; McGee, C. E. 1986. FORCAT: A single tree model of stand development following clearcutting on the Cumberland Plateau. Forest Science. 32(2):297-317.

Decision-Making for Natural Regeneration in the Northern Forest Ecosystem

David A. Marquis, USDA Forest Service, Retired, Homosassa, FL 34446

Mark J. Twery, Northeastern Forest Experiment Station, USDA Forest Service, Morgantown, WV 26505

ABSTRACT Failure to obtain prompt regeneration of desired species after a harvest cut can leave a stand unproductive for many decades, cost excessive amounts to reclaim through artificial means, and severely limit the suitability of the stand for a wide range of forest values. But prescribing silvicultural treatments that ensure successful regeneration most of the time is a difficult task.

Forest managers need to consider all the many factors that could affect regeneration success, weigh the many available silvicultural techniques available to accomplish the task, consider the characteristics of the particular stand under consideration, and make an informed decision that is most likely to achieve the desired conditions. In this paper, decision-making procedures are described that provide a systematic way to analyze stand potential and prescribe regeneration treatments. In the Allegheny region where the system has been used extensively, successful regeneration has been obtained in over 90 percent of the stands harvested, compared to only about 50 percent prior to development of this system. The procedures can be adapted for use in any geographic region or forest type.

INTRODUCTION

Natural regeneration of oaks is difficult to obtain in the northern forest ecosystem, just as it often is in most other parts of the oak range. To the common problems resulting from infrequent acorn crops, acorn depredations, slow juvenile oak seedling growth, and intense competition from other fast-growing plants, northern ecosystems add additional obstacles. Many oak stands in the north are found on sites better adapted to northern hardwoods, and the natural tendency is for them to revert to northern species now that fire and grazing have been removed as factors favoring the oaks. In addition, many northern ecosystems support excessive deer populations that make regeneration of any woody species difficult.

Research now underway is providing much information that will help us develop silvicultural procedures to perpetuate the oaks. We hope that this research will also define the range of sites where the battle to maintain oaks—rather than allowing conversion to northern species—is economically and ecologically feasible. There is still a long way to go before reliable procedures to regenerate oak on good northern sites are available, but progress is being made. One point is already quite clear, however: there will be no simple or universal treatment that guarantees consistent oak regeneration across the wide range of stand and site conditions found throughout the oak region. Instead, success will depend upon the careful prescription of treatments tailored to each individual situation. Stand and site conditions and all potential obstacles to oak regeneration will need to be evaluated systematically before a specific treatment is recommended. This decision-making process will require considerable knowledge and judgment on the part of the forest manager.

About 12 years ago, a systematic procedure for making silvicultural prescriptions for individual cherry-maple, northern hardwood, and oak stands in the Allegheny region was developed, and that procedure has been refined as new knowledge has accumulated. This stand evaluation and prescription process, known as the SILVAH system (Marguis and others 1992, Marguis and Ernst 1992), has been widely used in the Allegheny region, where it has helped forest managers consistently prescribe successful treatments in areas where deer browsing and other factors had traditionally prevented regeneration. That basic decision-making procedure is now being expanded to cover other forest types and geographic regions of the Northeastern United States (Marquis 1991a). In this paper, decisionmaking procedures we've developed for regeneration prescriptions in the Northeast are described. The process could be applied anywhere in the oak region if decision criteria appropriate to the geographic area are substituted for the Northeastern criteria. These decision-making procedures are also being expanded within the Northeast Decision Model for purposes beyond timber production. The successful establishment of new young trees after disturbance is important for all values derived from forests. Although the decision criteria may differ, the same data and decision procedures may be used in making regeneration decisions where wildlife, aesthetics, water, or other forest values are the management goal.

THE DECISION-**MAKING SYSTEM** The decision-making system involves the use of (1) a set of decision charts, (2) specific criteria to be used in making each decision, and (3) data on which to determine the condition or characteristics of the stand in question.

Decision Charts Decision charts are simply an outline of the decision-making process. All of the factors known to influence regeneration are included, and these factors are arranged in an order that systematically determines that stand's suitability for particular silvicultural techniques. As implemented in SILVAH and the new NE Decision Model, these are dichotomous charts that ask a series of yes/no questions about the stand and site under consideration.

A portion of the new NE Decision Model chart for even-age silviculture serves as an example (charts 8, 8.1, 8.1a, 8.1b, and 8.1z, from Marquis 1991b). Other portions of the NE charts deal with intermediate even-age culture and with other silvicultural systems.

In these charts, specific symbols are used to denote their function in the decision chart. Circles denote the start of the process. Five-sided boxes are connecting

symbols that route you to (or from) other charts. Arrows denote the sequence of travel within the chart. Horizontal lines with a short question above the line represent the main decision points in the chart. Rectangular boxes at either end of the decision question represent the possible answers—in this case, either "Yes" or "No." Ovals represent prescriptions, or recommended treatments; once you reach an oval, you have completed the decision process for that stand. Note in the legend for chart 8 (next page) that some prescription ovals are shaded or cross-hatched, indicating different degrees of confidence in the recommendation and/or various levels of investment required.

Target Species

Target species, as used in the Prescription Charts, refers to the species or species group that you wish to reproduce and/or manage for. For example, you may choose to manage for the oaks, or for white pine, or all northern hardwoods. If the prescription derived from the

charts requires planting or protection against browsing, or other unacceptably expensive treatments, you may wish to redefine the target species to include a broader group of species already present on the site, then re-determine the prescription to evaluate that alternative.



All decision criteria in the NE Decision Model are expressed as Yes or No answers to the questions asked at each decision point in the prescription charts. Amounts, levels, or values that constitute a Yes or No answer are defined in the decision criteria tables. There is a separate table for each forest type.



These prescriptions generally produce the desired results in the opinion of the NE Decision Model experts. They require no cash investments, and are highly recommended.



These prescriptions generally produce the desired results, but require an investment. If such investments meet your organization's economic criteria, we recommend them. If not, we recommend that no cutting be done in this stand. In the case of regeneration prescriptions, stands generally will not reproduce without the recommended treatment. For timber goal

4 (maximize timber value) these treatments will be recommended if you choose to maximize income, but not if you choose to maximize internal rate of return.



These prescriptions will not always produce the desired results, and they are quite expensive. The experts who developed the NE Model do not recommend them under either option of timber goal 4, but they are shown in case you feel you want to do something. We generally recommend that you devote your attention to more productive stands, and avoid expenditures in these difficult ones.

An alternative to both shaded and cross-hatched prescriptions is to shift your regeneration objective to less difficult target species.


Legend-Prescription Charts.

Note also that the charts are divided into zones by horizontal lines, and that there is a decision question on the left edge of the chart for each of these zones. The zones and their associated questions are included only to make the charts easier to understand.

Portions of the decision process that led to chart 8 are omitted because they do not involve regeneration considerations. However, to arrive at chart 8, you would have had to determine that even-age silviculture was the appropriate silvicultural system to use in this management unit. In the NE Decision Model, eight separate silvicultural "systems" are recognized, of which traditional even-age silviculture is number eight. This decision on the overall silvicultural system to be used is based primarily on the specific management goals selected, plus the forest type.

The eight systems in the NE Decision Model are viewed as a continuum on two different directions: intensity of disturbance and frequency of disturbance. The even-age system described in chart 8 is at one end of each continuum—intense but infrequent disturbance. Of course, the system selected is very important in


Chart 8 -- Even-age Silvicultural System



determining which of the potentially desirable species are likely to succeed under a prescription made by the model. Tracing a few prescriptions through the decision chart will illustrate its use.

The first chart in this series, chart 8, has only one zone, and its purpose is to determine the stage of development of this stand. If the stand is mature and ready to be regenerated, you will be routed to the 8.1 section of the chart. If regeneration treatments have already been applied, but the stand is still in the process of regenerating, you will be routed to the 8.2 section of the chart where the need for weeding, protection against animal depredations, and similar matters

Chart 8.1 -- Mature Stand to be Regenerated Possible Removal Cut



are considered. If the stand is in the sapling or larger stage but not yet mature, you will be routed to section 8.3 of the chart where the need for intermediate thinning and similar treatments will be evaluated.

Starting at the top of chart 8, the first decision point we reach contains the question: "Is this stand mature?" If we answer "Yes," we are routed to chart 8.1, which is the beginning of the section on regeneration prescriptions.

Remember that the decision charts are simply the outline for the decision process. To decide whether to answer the question about stand maturity "Yes" or "No,"

we'll need specific criteria on what constitutes stand maturity. Those criteria will be covered later.

At the top of chart 8.1, the five-sided box completes the connection from the previous chart. As the title of chart 8.1 indicates, it will consider the possibility that a removal cut is appropriate in this stand. The first zone deals with the adequacy of existing advance regeneration plus expected sprouts to regenerate the stand if harvested. There are three specific decision points or questions in this zone.

The simplest case involves those stands that already have regeneration of sapling size established in adequate amounts to form a new stand. Such a stand might have resulted from a heavy cut 10 to 25 years earlier, perhaps a diameter-limit cut or the second cut of a three-cut shelterwood. All that needs to be done is to remove the remaining overstory of the old stand to release the new stand. This prescription is reached if you answer "Yes" to the first decision question, "Is Sapling Regen Adequate?" Again, the criteria by which you decide whether or not sapling regeneration is adequate will be discussed later.

If sapling regeneration is not adequate, follow the route through the "No" box to the question "Is Advance Regen + Sprouts of Target Species Adequate?"

A "Yes" answer here moves us through the first zone to additional zones that deal with interfering plants and site limitations. A final removal cut, or clearcut, will be recommended only if interfering plants and site are not limiting.

If interfering plants are dense, removal cutting is not likely to provide satisfactory regeneration of the desired species, and you are routed to chart 8.1a where alternative techniques are considered. Interfering plants here include any plants that may interfere with successful regeneration of desired or target species. These may be herbaceous plants like fern and grass, woody plants of no commercial timber value like striped maple, dogwood, and sassafras, or commercial tree species such as yellow-poplar and red maple that outgrow desired oaks species under full sunlight. The criteria for evaluating these interfering plants deal with both those plants present in the understory before cutting and those present in the forest floor as dormant seed banks.

If certain site factors alone are limiting to regeneration, the recommendation will be for the removal harvest to be done in a sequence of two cuts that ameliorate the site limitations (wet or extremely rocky surface soil), with the first removal cut to be made now. Retaining 35 to 40 percent overstory density until the advance Page 224 of 404 seedlings and sprouts grow to small sapling size and have established roots in deeper soil layers has been found effective in maintaining a transpiration pump to avoid saturated conditions on wet sites, and in stabilizing the surface organic soil on rocky areas, as opposed to single-cut removals which accent the soil limitations.

Going back to the first zone, a "No" answer will lead to a repeat of the question about adequacy of advance regen + sprouts, assuming the area is fenced against deer. In areas of high deer population, the numbers of advance seedlings and sprouts needed to assure successful regeneration is very high, since deer browsing destroys many of them. A fence to exclude deer has the effect of reducing the numbers of advance seedlings and sprouts needed, and may permit removal cutting to occur where it would otherwise not be feasible. This section of chart 8.1 is identical to that discussed above, except that any prescription for a removal cut now includes erection of a deer fence as well.

If advance regen + sprouts of target species are not adequate even if fenced, you are routed to chart 8.1 where alternative techniques to encourage advance regeneration establishment are considered.

The remaining charts work in much the same way, but deal with other circumstances.

Chart 8.1a considers the possibility that a shelterwood sequence will permit advance seedlings to develop. Decision points here include evaluations of interfering plant density, overstory density or shade level, seed source availability, and deer browsing. Note that the prescriptions here include both natural and artificial regeneration, that shelterwood seed cuts may be recommended alone or in combination with either artificial regeneration and deer protection (which could include tree shelters in the case of oaks if ongoing research proves them effective). Note also that a "wait" (wait a few more years for advance seedlings to develop) prescription occurs when the desired level of shading or shelter already exists; this prescription also may be combined with planting and/or protection.

Chart 8.1b is a duplication of 8.1a, except that it deals with stands where interfering plants are expected to cause trouble. Thus, the prescriptions in this chart all include a herbicide recommendation along with the shelterwood, wait, plant, and/or protect recommendations.

Chart 8.1z is to be used in combination with all of the prescriptions obtained earlier, and includes supplementary treatments to be applied along with the primary prescription. These include the need for grapevine control, the need to treat unmerchantable saplings and poles, and the need or desirability of selecting individual stems from the existing stand to be retained after final harvest for specific purposes. If residuals are to be retained, it is important to identify them during any cuts prior to final harvest, so that they are not removed inadvertently at the earlier cut.

Thus, the decision charts provide an outline for the decision process. Major factors such as the adequacy of advance regeneration, expected sprouts, seed source availability, interfering plants, deer browsing, site limitations, and overstory shade conditions are evaluated, and the questions lead you quickly through the decision process.

Some factors of known importance may appear to be missing from the decision charts. For example, we know that acorn insects can be a major factor in oak regeneration. Although not included individually in the decision charts, losses of acorns are included in the criteria used to determine whether or not seed supply is adequate. Other pest problems, and many other factors are or could be included in a similar manner, as part of the decision criteria.






Chart 8.1a -- Mature Stand to be Regenerated



Page 227 of 404



Decision Criteria

they are encountered in the decision charts. Allegheny region. The decision questions are listed in the approximate sequence answer, are illustrated in table 1 (from Marquis 1910). Table 1 is from the Tables of these parameters, and the levels that qualify for either a "Yes" or "No" $\,$ on the basis of specific parameters that are measured in the stand under question. intent of the system that these determinations be made as objectively as possible, "Yes" or "No." Although the questions and answers appear subjective, it is the When using the decision charts, one must answer each decision question either

where success requires the regeneration of specific target species (such as oaks). Similar tables are being developed for other geographic regions, and for situations

(91661 Table 1- Decision criteria for the Allegheny Hardwood Forest Type (from Marquis

Decision question	Parameter used	VES Decision	I criteria NO
Is this stand	MDM (Medial Stand Diam-	WDW ∑ 18. <u>O</u> B	<u>anv</u> "81 > Man
mature?	eter of the Merchantable	AGS RD <35%	¥G2 KD ≥32%
	trees 5.5+); and AGS RD	(for timber pro-	
	(Relative Density in Ac-	duction goal;	
	ceptable Growing Stock)	for other goals, see regulation	
		parameter table)	
Is this stand in the	Regen. Stocking: % of	MOM $3ud$, $807 \leq$	%0L>
process of regene- ration?	stems over 5 ft, tall	ADM in regu- lation parameter	
		alder	
-snogon guilqes el	Sapling Regen. Stocking:	%0L <	%0L>
fation adequate?	% of plots with desirable sapling regen. See Re-		
	generation Stocking Cri-		
	teria tables in Input		
	Data documentation		
ls advance regene-	Advance Regen. Stocking:	%0 <i>L</i> <	%0L>
ration plus sprouts	muminim this stold to %	~~~~	
of target species	number of desirable stems		
sdequate?	(weighted by height),		
	supplemented by desirable		
	sprouts expected (from		
	overstory data). See Re-		
	generation Stocking Cri-		
	teria tables in Input Da- ta documentation. Use		
	column for actual deer		
	impact index in this stand.		
			<i><i><i>b</i> 0<i>C</i> ></i></i>
ls advance regene- ration plus sprouts	Advance Regen. Stocking: same as above, but using	%0L<	%0L>
of target species	the column for fenced ar-		
sdequate if fenced?	eas (deer impact index =		
	numinim to sldut sht ni (I		
	weighted numbers attached.		

Decision question	Parameter used to quantify	YES	n criteria NO	
Are interfering plants extremely dense?	Interfering Plant Stock- ing: % of plots stocked with minimum numbers or percent coverage of in- fering plants. See Re- generation Stocking Cri- teria tables in Input Data documentation	<u>></u> 70%	<70%	
Are interfering plants dense?	Interfering Plant Stock- ing: same as above.	<u>></u> 30%	<30%	
Do site factors limit regeneration?	Site Limitations: % of plots on which there is evidence of either poor soil drainage or extrem- ly rocky surface soil	<u>≥</u> 30%	<30%	
Is sunlight limiting to seedling es- tablishment?	%RD (Relative Stand Density), as the percent- age of the maximum basal area expected for stands of this species composition and tree size.	<u>≥</u> 75%	<30%	
Is the seed source of target species adequate?	Seed Source Index for the target species: cal- culated from overstory data	Deer Index 1-3 1, 2, or 3 Deer Index 4 1 or 2 Deer Index 5 1	Values other than those under YES	
Is the seed source of target species adequate if protect- ed against deer?	Seed Source Index for the target species: calculated from overstory data.	1, 2, or 3	4	
Are grapevines a problem?	Grapevine Stocking: % of plots with minimum numbers of grapevines. See Regeneration Stock- ing Criteria tables in Input Data documentation.	plots with minimum mbers of grapevines. e Regeneration Stock- g Criteria tables in Input		
Are unmerchantable saplings and poles dense?	Basal Area/acre in un- merchantable sapling- and pole-sized trees	<u>≥</u> 10 sq. ft.	<10 sq. ft.	
Are residuals de- sired or needed?	 a) by user choice, in the User Profile, or b) Advance Regen. Stocking plus Residuals Stocking: % of plots stocked with qualifying residuals. See Regeneration Stocking Criteria tables in Input Data documentation 	Advance Regen. Stocking \geq 50% but <70% and Residuals Stock- ing plus Advance Regen. Stocking total \geq 70%	Any other values	

Table 1-Decision criteria for the Allegheny Hardwood Forest Type-Continued

The question "Is sunlight limiting to seedling establishment?" illustrates how specific stand parameters are used to arrive at an objective answer. The stand parameter used to evaluate the density of the overstory and thus the amount of sunlight reaching the forest floor is relative stand density or stocking percent. This measure works well in stands of any species composition, is easily calculated from standard cruise data, and requires no subjective judgment. In stands with more than 75 percent relative density, the overstory is dense enough that sunlight may limit seedling establishment. At lower densities, seedlings can usually become established, although shade-intolerant species may not grow much there. So, 75 percent relative density is the breaking point; at densities above 75, sunlight is considered limiting and shelterwood cutting becomes a possible treatment. At densities below 75, light is probably adequate for establishment and any lack of advance seedlings is probably due to other limiting factors.

Some of the decision questions require much more complex criteria to evaluate adequately. The amount of advance regeneration that is adequate to ensure successful reproduction following harvest is a good example.

In the Allegheny region, adequate advance regeneration requires that at least minimum numbers of seedlings be present over at least 70 percent of the stand area. The minimum numbers of seedlings depends on the species, their size, and the amount of deer browsing expected. Table 2 (from Marquis and others 1992) shows the *weighted* minimum numbers of seedlings required on each plot, as a function of species and deer impact index.

Weighted numbers are derived by counting (or estimating) seedling numbers in each of three size classes, and applying weights to each size class to arrive at a weighted total number. Larger seedlings have a better chance of survival, and thus carry a higher weight than small seedlings.

		Advan	ce regen speci	es group	
Deer impact index	Black cherry	Small oak	Other desirable species	All desirable species	Large oak
		Weighte	d number per 6	6-ft. plot	
5	50	60	200	200	1
4	25	40	100	100	1
3	20	30	50	50	1
2	15	20	30	30	1
1	10	10	15	15	1

Table 2-Number of seedlings required for regeneration plots to be stocked, for different levels of deer pressure (from Marquis and others 1992)

The deer impact index is a value determined from a chart showing the estimated deer population (in deer per square mile) for various habitat conditions. Deer population estimates may be obtained from the state game agency, or from a

sample census conducted for this purpose. Habitat condition is determined from the proportions of the 1-square-mile area surrounding the stand in question (the deer home range roughly) in such conditions as: agricultural fields, open land, recent harvest cuts, thinned and unthinned forest, etc. All of these vegetative conditions affect the amount of deer food available in the vicinity to reduce browsing pressure on regeneration in any new harvest cuts.

Thus, the evaluation of advance regeneration adequacy is based on the percentage of understory that meets certain required minimum conditions. These conditions vary tremendously depending on such factors as tree species, seedling size, and deer pressure. To illustrate: in the Allegheny region, the minimum number of maple or ash seedlings per 6-ft. radius plot is 100, if deer pressure is high (level 4) and all seedlings are 2 in. to 1 ft. tall; but, the minimum number is only 7 or 8 if deer pressure is low and all seedlings exceed 1 ft. in height.

Since the decision question in the chart considers both advance regeneration and sprouts, the sprouting potential of all trees to be cut must also be evaluated. Then, estimates of the proportion of the area that will be regenerated by sprouts must be added to the advance regeneration estimates.

Thus, a simple "Yes" or "No" answer in this case involves a whole host of factors that must be carefully evaluated to make effective use of the decision charts. Other decision criteria, such as the density of interfering plants and seed supply adequacy, are evaluated in a similar manner. Several factors must be included in determining whether or not these factors are limiting to or are adequate for regeneration. Details on these and other factors for the Allegheny region are presented in the SILVAH literature (Marquis and others 1992), and are not repeated here.

Developing suitable decision criteria for a specific forest type and geographic region is a major task. In the Allegheny region, it took nearly 15 years of research aimed specifically at that goal to complete the SILVAH system. Information needed to develop decision criteria for other regions and forest types that have received less focused research effort will not always be available.

Nevertheless, many of the general principles and procedures from the Allegheny region can be used elsewhere, and experienced silviculturists can probably make informed estimates of the critical parameters in their areas. Such estimates provide a starting point that is certainly better than no system at all. At the very least, it forces the decision-maker to systematically consider all of the many factors that affect regeneration success, and to make a judgment on each based upon the best information currently available.

Furthermore, the system itself serves as a model or framework on which to evaluate research needs. Those factors or criteria on which knowledge is inadequate become quickly apparent as one attempts to define the decision questions and criteria. The relative importance of each research need also can be assessed in terms of its relative importance and frequency of occurrence in the overall oak regeneration process.

The decision charts presented here are intended for general use over a wide range of forest types and geographic areas. As a result, some of the decision points and decision criteria will be of little importance in some types or regions. Rather than $P_{age 232 \text{ of } 404}$

creating a whole series of specialized charts that eliminate these factors entirely for the charts for that region, it is best to simply set the criteria at a level that will effectively ignore the factor. For example, deer browsing may not be a factor at all where deer populations are low across a wide geographic area. In these cases, the criteria table can simply suggest that deer impact index be set to 1 (very low) or 2 (low) anywhere in that region.

Additional decision criteria are being formulated to extend this system of decision making to a wide variety of resource values. For example, visual qualities within a forest stand are highly dependent on numbers and sizes of woody stems close to the ground, as is habitat quality for many wildlife species. Criteria that ensure the desired stems densities for these objectives fit very comfortably in this system.

Data Requirements To function consistently, any stand analysis and prescription system must have reliable data on which to answer the decision questions. In SILVAH and the NE Decision Model, that data come from a cruise in the stand under consideration. Data on overstory and understory vegetation and site factors are collected and summarized for use in the decision charts. Computer programs ease the calculation job and permit extensive analyses to be performed as needed.

The amount of detail collected in the cruise can be varied to meet the needs of a variety of organizations and individuals, although there are certain essential data items that must be collected in all cases. The recommended cruise procedure involves:

1. A variable-radius (prism) cruise of the overstory, recording trees by species and diameter at the minimum, with other observations on individual trees as desired (such as merchantable height, timber quality, grade, defect, wildlife value, etc.).

2. A fixed-plot sample of understory conditions, with several options in the amount of detail recorded. At the minimum, a simple checkmark is used to indicate the presence or absence of 15 to 20 specified conditions (such as the presence of various categories of advance seedlings and interfering plants). Estimated numbers or percent of coverage may also be recorded for each of these classes. Or, complete enumerations may be made of all vegetation by species and size classes.

3. A fixed-plot sample of site conditions. At the minimum, this involves a checkmark to indicate the presence of certain site factors known to limit regeneration. It may also include site index or site class determination and recording of other important site factors.

Because understory and site conditions vary considerably within a stand where overstory conditions are uniform, we recommend that twice as many understory plots be sampled as overstory plots. Understory and site data can easily be collected at the same time as overstory data. Understory and site plots are taken at each overstory plot, and additional understory/site plots are located half way between each overstory plot.

When the SILVAH system was first introduced, there was some reluctance on the part of forest managers to collect understory and site data. Traditional cruises

included only the overstory (often only the merchantable size trees), and it was assumed that the extra data collection would add considerably to cruise costs. However, studies by both the Allegheny National Forest and Hammermill Paper Co. revealed that collecting the minimum understory and site data did not add much to the time required. Recording simple checkmarks at each understory plot took only a few minutes per plot, and was insignificant compared to the time required to lay out cruise lines, walk from plot to plot, and collect prism data.

Of course, more detailed understory data collection would add significantly to cruise time and costs. Although the minimum data collection provides sufficient information for prescription preparation, extra data may be desirable for a wide variety of reasons. Each organization must tailor data collection to its own needs and budget. The system is designed in such a way that it can utilize the added data if available, but can still function for writing prescriptions with the minimum data.

SUMMARY Failure to obtain prompt regeneration of desired species after a harvest cut can leave a stand unproductive for many decades, cost excessive amounts to reclaim through artificial means, and severely limit the suitability of the stand for a wide range of forest values. But prescribing silvicultural treatments that ensure successful regeneration most of the time is a difficult task.

Forest managers need to consider all the many factors that could affect regeneration success, weigh the many available silvicultural techniques available to accomplish the task, consider the characteristics of the particular stand under consideration, and make an informed decision that is most likely to achieve the desired conditions. The decision-making procedures described here provide a systematic way to analyze stand potential and prescribe regeneration treatments. In the Allegheny region where the system has been used extensively, successful regeneration has been obtained in over 90 percent of the stands harvested, compared to only about 50 percent prior to development of this system. The procedures can be adapted for use in any geographic region or forest type.

- LITERATURE CITED Marquis, D.A.; Ernst, R.L.; Stout, S.L. 1992. Prescribing silvicultural treatments in hardwood stands of the Alleghenies (Revised). Gen. Tech. Rep. NE-96. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 101 pp.
 - Marquis, D.A.; Ernst, R.L. 1992. User's Guide to SILVAH: Stand analysis, prescription, and management simulator program for hardwood stands of the Alleghenies. Gen. Tech. Rep. NE-162. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 124 pp.
 - Marquis, D.A. 1991a. The Northeast Decision Model: a multi-resource silvicultural decision model for forests of the Northeastern United States. In: Systems Analysis in Forestry; 1991 March 5-7; Charleston, S.C.: 358-365.
 - Marquis, D. A. 1991b. Current specifications, NE Decision Model. Unpublished report; 1991 December 19. Available from the Northeastern Forest Experiment Station, Morgantown, WV. 101 pp.

Page 235 of 404

Natural Regeneration of Oaks

Regenerating Oaks in the Central States

Ivan L. Sander, USDA Forest Service, Retired, Columbia, MO 65202

David L. Graney, Southern Forest Experiment Station, USDA Forest Service, Fayetteville, AR 72701

ABSTRACT Although oaks are the most abundant overstory species in many upland stands in the Central States, they are often replaced in the reproduction that follows harvesting of current oak stands because of the lack of adequate oak advance reproduction. Obtaining adequate oak regeneration is especially difficult on highly productive sites where understories are often well developed and dominated by shade-tolerant species. Recommended regeneration methods for these upland stands are based on an evaluation of the oak regeneration potential. Where the regeneration potential is low because oak reproduction is absent, sparse, or too small, stand treatments to enhance oak reproduction establishment and growth are described. Silvicultural treatments for individual stands will vary with site quality, competing species (especially fast-growing competitors such as yellow-poplar) and physiographic-ecological region.

INTRODUCTION In the Central United States, geologic history, physiography, and climate vary greatly and many different associations of soils, water, plants, and animals are present. Braun (1950) described the forest regions of Eastern North America using remnants of the "presettlement" forests. The Central United States includes parts of her Oak-Hickory, Western Mesophytic, Mixed Mesophytic, Oak-Pine, Oak-Chestnut, Beech-Maple, and Maple-Basswood forest regions (figure 1), each with its own distinctive characteristics but with wide variation of forest vegetation. Oaks have always been an important stand component in these regions and currently dominate most of the mature or nearly mature stands. Even so, the oak component of new stands following harvesting of many mature stands is often less than expected or desired.

It is generally accepted that a key to replacing these stands with new oak stands is having well-established oak advance reproduction in place when the final harvest is made. Sprouts from stumps of cut overstory trees are also important. It is now apparent that existing oak advance reproduction plus stump sprout potential is not adequate to replace many current oak stands, especially those on the most productive sites. How then do we ensure that when the stand is ready to harvest the oak advance reproduction will be adequate to replace it?




Figure 1-Area of central hardwoods with Braun's Forest Regions.

EVALUATING OAK REGENERATION POTENTIAL

When planning regeneration harvest cuts in oak stands, the important first step is to evaluate the potential of the oak advance reproduction plus stump sprouts to replace the current stand. Oak advance reproduction differs greatly in number and size of individuals. Some of this difference can be attributed to site quality; in general, oak advance reproduction in the Central States is more abundant on average to poor sites than on good sites (table 1). Size of individual stems is important because growth after final harvest is related to both stem height and basal diameter. Thus, height and basal diameter are indicators of total plant size and the potential of the root systems to support vigorous growth after overstory removal.

Assessing oak regeneration potential involves making an inventory of advance reproduction and trees over 2 in. d.b.h. in the midstory and overstory. The advance reproduction data are used to determine the probability that a stem of a given size will attain a specified height by the time it is 5 years old. These probabilities are then adjusted to account for drop-out from mortality and suppression to age 20. Probabilities have been developed for the Ozark Forests in Missouri and are related not only to stem size, but also to aspect and slope position. For stems of equal height and basal diameter, the probabilities are highest on southeast- and northwest-facing middle slopes and lowest on lower, northeast slopes (Sander and others 1984). The coefficients in the regeneration potential equation apply specifically to Missouri and must be used cautiously in other areas.

The midstory and overstory inventory data are used to estimate the proportion of stump sprouts from cut trees that can reasonably be expected to produce dominant or codominant oaks at age 20. This estimate, together with the advance reproduction probabilities, is used to determine if the new stand can be expected to develop into a predominantly oak stand by age 20.

This evaluation procedure is for oaks only. The new stand will contain other species, but we do not have a procedure to estimate their contribution to future stand stocking. We can get a general idea of what the new stand composition might be by looking at the total composition of the advance reproduction and the composition of the overstory (table 1).

State	Oaks ¹	Hickory	Maples	Yellow poplar	White Ash	Other Trees ²	Total Trees	Under- Story ³	• Total Stems
				(Good Site				
Arkansas	940	820	1420		430	1640		4420	
Indiana	500	360	4350	40	1570	1000	7820	3570	11390
Kentucky	1460	610	1840	150	0	690	4750	2920	7670
Missouri	1430	640	380		0	420		2060	
Ohio	1960	550	860	240	80	390	4080	3770	7850
				A	verage Si	ite			
Arkansas	1950	600	1960		460	1560		3370	
Indiana	1580	100	1640	50	800	400	4570	5330	9900
Kentucky	1070	580	870	140	490	630	3780	1510	5290
Missouri	1510	514	329		0	440		1540	
Ohio	3640	700	420	140	0	840	5740	7310	13050

Table 1-Advance reproduction per acre on good and average sites in some Central States

¹ White, black, northern red; includes scarlet oak in Missouri and scarlet and chestnut oaks in Indiana, Kentucky and Ohio.

² Good Sites: Elm, blackgum, scattered black cherry and black walnut. Includes American beech in all states except Missouri. Also includes sweetgum in Arkansas. Average Sites: Primarily blackgum in all states.

³ Good Sites: Primarily flowering dogwood, redbud, hazelnut, witch-hazel, paw-paw, and hophornbean. Includes sourwood in Kentucky and Ohio. Includes Carolina buckthorn in Arkansas. Average Sites: Primarily flowering dogwood, sassafras, and serviceberry. Includes sourwood in Kentucky and Ohio, Carolina buckthorn in Arkansas, and hawthorn in Missouri.

When yellow-poplar is an overstory component, its regeneration potential will be substantial. This potential exists because numerous yellow-poplar seedlings will appear, even after partial harvests, from the large quantity of viable seed in the litter and humus. The primary source of reproduction of species such as white ash, black cherry, hickories, maples, elm, blackgum, dogwood, sourwood, and sassafras is advance reproduction. Like the oaks, their growth after harvest is related to their pre-harvest size; relatively large advance reproduction stems have a higher potential for competing successfully than small ones. Stump sprouts are also an important source of regeneration for these species. Although we do not yet have a system for estimating success probabilities for species other than oaks, we can expect at least some of them to become a dominant component of the reproduction stand after harvest if they are present as relatively large advance reproduction.

CHOOSING A REGENERATION METHOD

The choice of which regeneration method to use must be based on the outcome of the regeneration potential analysis, the basic ecology of the forest region in which the stand is located, and the overall management objective. On many sites in the areas east of the Mississippi River it is not realistic to expect to be able to naturally regenerate essentially pure oak stands even though the current stand is dominated by oak. The ecological trends due to effective wildfire control, and results of past Page 239 of 404

regeneration efforts, indicate many future stands will be mixtures of a number of species especially on high-quality sites. An oak component can be retained readily on average sites, but on good sites the stands will most likely be dominated by species other than oaks.

If the oak reproduction potential is adequate to replace the current stand, clearcutting is the best method to use. Although understory species may appear to dominate the stand for about 10 years following clearcutting, the oaks and other overstory species begin to assert dominance, and by age 10-15 the understory species are generally in a subdominant position.

If oak reproduction potential is not adequate and the stand is clearcut, the new stand will be dominated by a varying mixture of species. With the exception of yellow-poplar, the species that dominate the advance reproduction will be predominant in the new stand. Yellow-poplar will also be abundant if it is present in the overstory, and some oaks will probably be present.

When oak advance reproduction is small, scarce, or absent, the regeneration method most likely to produce the best results is the shelterwood method. However, the method must be tailored to produce the micro-environments required by oaks for successful seedling establishment and early seedling growth. These micro-environments are still largely unknown and are likely to differ among differing ecological regions. For example, because of the differences in the basic ecology between the Mixed Mesophytic and the Oak-Hickory forest regions, what is required in one of these areas may not be required in the other.

In 1979, a study of the shelterwood method with overstory densities of 40, 50, and 60 percent stocking and three understory treatments that ranged from heavy (all non-oaks treated with herbicide) to none, was started in Missouri's Ozarks. At the time of treatment the number of oaks per acre varied from about 300 to 1,100 per acre on good sites (Black oak site index 75+) and from about 600 to 1,200 per acre on average sites (Black oak site index 60-75) (Sander 1987). However, a large majority of them were less than 1 ft. tall, much too small to compete if the stands were clearcut. Ten years after treatment the total number of oaks increased and ranged from about 600 to 1,800 and about 2,500 to 3,500 per acre on good and average sites respectively (figure 2), although the total number of oaks increased in all treatments at all overstory stocking levels.¹ Even so, when the advance reproduction evaluation criteria were applied to these plots, stocking values were less than one-half that required for successful oak regeneration on the good sites. Furthermore, about 50 percent of the large reproduction is understory species, primarily dogwood and sassafras. Maple, hickory, and blackgum account for an additional 15 to 25 percent of the large reproduction. Thus the oaks, particularly the smaller stems, are still at a competitive disadvantage.

On the average sites understory treatment had little effect on increasing the number of large oaks, but there were more large oaks on the 40 percent stocking plots than on the plots with higher stocking¹. The advance reproduction stocking values are

¹ Schlesinger, R.C., Sander, I.L., and Davidson, K.R. 1992. Unpublished draft on file: North Central Forest Experiment Station, Columbia, MO.




Figure 2—Reproduction per acre 10 years after shelterwood cutting in the Missouri Ozarks. Species: O—oaks, N—non-oak overstory species, U—Understory species. Treatments: Overstory—40-, 50-, and 60-percent stocking; Understory—H, Kill all non-oaks; M, Kill all non-oaks taller than 6 ft.; N, Kill nothing.

now close to those required for successful stand regeneration on the 40 percent stocking, but only about one-half to two-thirds of the required value on the higher density plots. The large reproduction is still dominated by hickory, black gum, sassafras, and dogwood. However, the large oaks appear to be at least holding their own and should in a few more years be adequate to replace the original stands.

Results from a similar study in southern Indiana show that all treatments failed to increase the number of oaks. There were few oaks present before treatment so a major objective was to get new oaks established. Two things occurred that precluded new oak seedling establishment. One was the lack of any acorn production the first 3 years after treatment, and the other was the very rapid regrowth of the understory. Nine years after treatment a major windstorm damaged nearly all of the plots and the study had to be abandoned. Although these early results are somewhat discouraging, they provide valuable information and insight for future research, and are useful for making immediate recommendations for using shelterwood cutting to regenerate oaks.

Interest in uneven-age regeneration methods, particularly the group selection system, has increased greatly in recent years. This interest has come about because of political and sociological pressures against clearcutting. Research on the effect of opening size on regeneration has demonstrated that oaks can be reproduced in small openings as well as in large ones (Minckler and Woerheide 1965, Sander and Clark 1971, Smith 1981). However, the requirements that apply to clearcutting also apply to group selection. Creating an opening, even a small one of one or two tree heights in diameter, will not in itself ensure oak reproduction. Oak advance reproduction of adequate size and in adequate numbers must be present where openings are created in order to achieve successful oak stand replacement.

We cannot, however, consider only the regeneration when we use group selection. Because it is a variation of the single-tree selection system, we must also strive to achieve and maintain a reasonably well-balanced diameter distribution. We have only begun to address the problem of how the reproduction in groups eventually contributes to overall future stand structure, and how to maintain balanced diameter distributions in oak stands has not been determined.

REGENERATION GUIDELINES The first step in planning for oak regeneration is to evaluate its regeneration potential. The results of this evaluation will determine which even-aged regeneration method should be used. Or, if group selection is planned, such an evaluation can help determine where to locate groups and can provide information on which species are reproducing and how the reproduction is growing.

Even-Age Methods Clearcutting. If the oak advance reproduction plus stump sprouting is adequate, clearcut.

When clearcutting:

1. Determine size of area to be designated as a stand. Stand size can vary but should be at least 2 acres. Stands smaller than this have a large proportion of their area in a zone around the stand border where reproduction growth will be slow

because of the influence of the surrounding trees. A stand should preferably be restricted to a single condition or size class of timber and site quality. Size of the forest property will also influence stand size. In areas where deer populations are high, stand size may need to be relatively large in order to reduce the impact of browsing on reproduction growth.

2. Arrange and shape clearcuts so they mingle with uncut stands and blend into the landscape as much as possible.

3. Plan, construct, and maintain skid trails and logging roads to minimize erosion.

4. Harvest all merchantable trees.

5. Cut or kill remaining culls and small trees larger than about 2 in. d.b.h. Killing the unwanted trees instead of cutting them reduces sprouting and provides snags for nesting holes and perches for birds. Cutting some of them will provide habitat for other wildlife species such as ruffed grouse.

6. There may be some other options for harvesting and getting rid of the unwanted and unmerchantable trees. These options are generally used for special purposes, such as to soften the aesthetic impact or provide special wildlife habitat. Their use depends on owner policy rather than silvicultural desirability.

If the clearcut stand is on southeast or northwest middle and upper slopes, we can expect to have a stand at about age 20 that can be molded into an essentially pure oak stand by thinning. On north and east aspects and lower slopes, the stand composition may be highly variable. In the mixed and western mesophytic forest regions, yellow-poplar will likely be abundant. Other species such as white ash, black cherry, and red and sugar maples will also be present. However, if the oak advance regeneration is adequate, we can expect to have a predominantly oak stand 20 years after clearcutting.

Shelterwood. When the regeneration potential of the existing oak advance reproduction is not adequate to replace the stand, use the shelterwood method. Oak advance reproduction is most likely to be inadequate on the middle and lower north- and east-facing slopes, where it will be difficult to obtain oak regeneration. Costs will be high because of the intensive site preparation needed on these high-quality sites.

When applying the shelterwood method:

1. Determine stand size, arrangement, and shape the same as for clearcutting.

2. Control the understory by cutting or preferably killing the non-oak species that will compete with the small oaks in the oak-hickory and oak-pine forest regions.

3. Reduce the overstory to 40-60 percent stocking. In the other forest regions, a somewhat higher stocked residual stand (70-80 percent stocking) might be desirable to inhibit yellow-poplar seed germination and rapid redevelopment of the understory. Leave the best dominant and codominant oaks as uniformly spaced as possible. Kill all stems of unwanted species.

4. If possible, apply the understory and overstory treatments before seedfall in a good seed year.

5. Monitor seedling establishment and growth and make additional light cuts to keep the overstory from restricting growth. Apply additional understory control if the understory redevelops to a point where it restricts the oak reproduction growth. This control may be desirable 5-10 years after the original treatment, particularly on high-quality sites.

6. When the regeneration potential of the oak reproduction is adequate to replace the stand, remove the remaining overstory trees in one cut.

The length of time required to establish oaks and grow them to adequate size under a shelterwood is not yet known, but will probably be 10 to 20 years or more.

Uneven-Age Methods Single-tree selection. The establishment and development of oak regeneration is not possible using the single-tree selection system. Harvesting single trees to achieve and maintain a specific diameter distribution does not provide the microclimate needed for oak regeneration, but does provide the conditions needed for the establishment and growth of shade-tolerant species. Over time, single-tree selection will convert the stand from oaks to shade-tolerant species.

Group selection. This method can be used to reproduce oaks satisfactorily, but the diameter of the circular openings created should not exceed 1 to 2 times the height of the surrounding dominant trees in order to maintain the uneven-age character of the stand.

When applying group selection cutting:

1. Evaluate the potential of the oak advance reproduction to fill each opening (group) that will be created by cutting.

2. If the oak advance reproduction is adequate, harvest the merchantable trees in the group and cut or kill remaining culls and small trees as described under clearcutting. The reproduction response will be similar to the responses after clearcutting except that reproduction growth will be retarded in a large part of the opening area because of the influence of the surrounding stand.

3. If the oak advance reproduction is not adequate to fill the opening, cutting the trees to create the opening will not result in oak reproduction and the opening will be filled by whatever species are present in the understory. In this case, follow the procedure for the shelterwood method where the openings will be located. The percentage stocking goal may not be attainable in the small groups, so make sure seed-producing oaks are left where the groups will be located. Treat the understory as in the shelterwood system. Removing sub-canopy overstory trees and the understory competition may be all that is needed to increase the amount of light reaching the forest floor and enhance oak reproduction establishment and growth.

4. Cut trees also between the groups to maintain the selected diameter distribution and to enhance oak reproduction establishment and growth throughout the stand.

5. Group selection may not be successful in areas with high deer populations because of excessive browsing in the small openings.

IN CONCLUSION There is no guarantee that applying the above guidelines will regenerate oaks successfully in all situations. However, doing nothing in the current mature or nearly mature stands will not increase the chances of regenerating oaks. On high-

and Beech-Maple forest regions, regenerating essentially pure oak stands on high-quality sites may not be possible. However, oaks have always been components of stands in these forest regions on all sites and it is likely that oaks will at least oak regeneration should not be difficult to obtain on average sites; however, on quality sites in particular, regenerating oaks is a difficult task. In some parts of the Central States, particularly in the mixed and western mesophytic, Maple-Basswood, be a component in future stands. In the Oak-Hickory and Oak-Pine forest regions, high-quality sites intensive treatments are probably going to be necessary.

The		
ΡA		
ITTERATURE CITED Braun, E. Lucy. 1950. Deciduous Forests of Eastern North America. Philadelphia, PA: 1ne		
America.		-
North		:
Eastern		•
of		1
Forests		
sn		
iduo	pp.	'
Dec	296	
1950.	Blakiston Company. 596 pp.	
Lucy.	n Com	
щ	cisto	
Braun,	Blal	
CITED		
С Ш	, 	
UR)	
AT		
ER.	i	
L		

Minckler, Leon S.; Woerheide, John D. 1965. Reproduction of hardwoods 10 years after cutting as affected by site and opening size. Journal of Forestry. 63:103-107.

Sander, I. L. 1987. Oak reproduction establishment and early development following shelterwood cutting in Missouri. In: Proceedings, fifteenth annual hardwood symposium; 1987 May 10-12; Memphis, TN: The Hardwood Research Council: 17-30.

Sander, I. L.; Clark, F. B. 1971. Reproduction of upland hardwood forests in the Central States. Agric. Handb. 405. Washington, DC: U.S. Department of Agriculture. 25 pp. Smith, H. C. 1981. Diameter of clearcut openings influence Central Appalachian hardwood stem development. Res. Pap. NE-476. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 8 pp. Clark, F. B., Technical Editor; Hutchinson, J. Editor. 1989. Central Hardwood Notes. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. Other Selected References Graney, D. L.; Rogerson, T. L. 1985. Growth of oak, ash, and cherry reproduction following overstory thinning of upland hardwood stands in the Boston Mountains of Arkansas. In: Dawson, J. O.; Majlrus, K. A., eds. Proceedings fifth central hardwood forest conference; 1985 April 15-Urbana-17; Urbana-Champaign, IL. Society of American Foresters Publication 85-02. Champaign, IL: University of Illinois, Department of Forestry: 4-10.

Proceedings fifth biennial southern silvicultural research conference; 1988 November 1-3; Graney, D. L. 1989. Growth of oak, ash, and cherry reproduction following overstory thinning and understory control in upland hardwood stands of northern Arkansas. In: Miller, J., compiler. Memphis, TN. Gen. Tech. Rep. SO-74. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 245-252.

Challenges in oak management and utilization; 1985 March 28-29; Madison, WI: University of Johnson, P. S. 1985. Regenerating oaks in the Lake States. In: Johnson, J. E., ed. Proceedings Wisconsin, Extension Service: 98-109.

and shelterwood removal cutting. Res. Pap. NC-202. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 5 pp. Johnson, P. S.; Jacobs, R. D. 1981. Northern red oak regeneration after preherbicided clearcutting

Johnson, P. S.; Jacobs, R. D.; Martin, A.J.; Godell, E. D. 1989. Regenerating northern red oak: three successful case histories. Northern Journal of Applied Forestry. 6:174-178.

Lorimer, C. G. 1983. Eighty-year development of northern red oak after partial cutting in a mixedspecies Wisconsin forest. Forest Science. 29:371-383.

- Lorimer, C. G. 1989. The oak regeneration problem: New evidence on causes and possible solutions. Forest Resource Analyses No. 8. Bull. R3484. Madison, WI: University of Wisconsin-Madison, Department of Forestry. 31 pp.
- Sander, I. L. 1977. Managers handbook for oaks in the North Central States. Gen. Tech. Rep. NC-37. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 35 pp.
- Sander, I. L.; Johnson, P. S.; Rogers, R. R. 1984. Evaluating oak advance reproduction in the Missouri Ozarks. Res. Pap. NC-251. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 16 pp.
- Schlesinger, R. C. 1976. 16 years of selection silviculture in upland hardwood stands. Res. Pap. NC-125. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 6 pp.

Regenerating Oaks in the Bottomlands

W. K. Clatterbuck, Tennessee Department of Agriculture, Division of Forestry, Nashville, TN 37204 J. S. Meadows, Southern Forest Experiment Station, USDA Forest Service, Stoneville, MS 38776

- **ABSTRACT** Bottomland oaks depend on advance reproduction and stump sprout potential for successful natural regeneration. Large seedlings must be present or be developed prior to harvest to ensure successful regeneration of oaks. Small oak seedlings, with their slow initial growth rates, are not able to compete with the faster initial growth of other species. This paper addresses the effects of the various methods of reproduction on the regeneration of bottomland oaks.
- **INTRODUCTION** Bottomland forests are some of the most productive and diverse ecosystems in the United States, yet they are poorly understood. Hardwood regeneration in the bottomlands can be established by most reproduction methods. Problems occur when foresters want to regenerate stands to specific species, particularly oaks. A myriad of different species with different site requirements and growth habits makes regeneration of oaks extremely complex and variable.

Although regenerating bottomland oaks is a major concern in these forests, Hodges (1989) stated that

"the answer to the question of how to ensure adequate oak regeneration in bottomland hardwood stands is not the development of some radically new method of cutting, but recognition that all cutting operations in the stand, from the very first, should have as some of their objectives creation of an environment, largely light conditions, favorable for oak regeneration" . . . and furthermore . . . "ensure that cuttings occur frequently enough to maintain growth of oak regeneration."

Unfortunately, most bottomland stands in the South are not managed. Stand structure typically ranges from closed canopies with little advance reproduction to cutover, mismanaged areas where species composition and tree quality are poor. Regeneration of such stands is certainly complex, variable, and difficult to predict.

This paper discusses various species-site relationships and reproductive mechanisms that should be considered when regenerating bottomland oaks and the various methods of reproduction that can be used to obtain successful oak regeneration. The term "bottomland oaks" in this paper refers to: cherrybark oak, Nuttall oak, Shumard oak, swamp chestnut (cow) oak, water oak, and willow oak. Maps showing the distribution of these bottomland oaks can be found in Burns and Honkala (1990).

SPECIES-SITE When natural regeneration methods are used to perpetuate bottomland oaks, the first step should be a detailed evaluation of site and stand conditions to determine RELATIONSHIPS the potential for regeneration (Johnson and Deen, in press). Matching appropriate species to suitable sites is especially important considering the wide variety of landforms (bars, levees, ridges, flats, sloughs, terraces) found in bottomlands and the different species of oak, each with specific silvical characteristics and site requirements (McKnight and others 1981). Slight, nearly undetectable elevational changes in bottomlands alter species-site suitabilities (Hodges and Switzer 1979). Other factors affecting regeneration are the frequency, depth, timing, and duration of flooding and the species' tolerance to flooding. Baker and Broadfoot (1979) published a site evaluation guide for commercially important bottomland Typical species compositions for particular landforms have been hardwoods. developed for the major bottoms of the lower Mississippi alluvial plain and the minor bottoms of the Gulf coastal plain (figure 1) (Hodges and Switzer 1979) and the red river and black river bottoms of the Atlantic coastal plain (Kellison and others 1988).

> Along with flood tolerance, shade tolerance is another variable in the regeneration equation. Bottomland oaks range from intolerant to intermediate in their tolerance of shaded conditions (Putnam and others 1960) and most require some direct overhead sunlight to survive and grow (Johnson 1979). Even though oak seedlings will become established in almost any size of opening, continued growth and development require large openings. In small openings, sunlight becomes a limiting factor, especially as canopies close, and actually favors the more tolerant species. Consequently, bottomland oaks are managed most successfully by implementing an even-aged silvicultural system (Kellison and others 1988).

REPRODUCTIVE MECHANISMS

When planning for natural regeneration of bottomland oaks, the presence of advance reproduction in both size and number is essential for success (Hodges and Janzen 1987). Oak regeneration consists of three forms of reproduction:

(1) New seedlings from seed in place.

(2) Seedlings that developed over a period of years prior to harvest (i.e., advance reproduction).

(3) Seedling and stump sprouts.

New oak seedlings generally cannot compete with sprouts and previously established reproduction (Johnson and Krinard 1976). Sprouts of oak trees less than 10 in. in diameter are a dependable source of regeneration because of their well-established root systems and vigorous growth potentials (Janzen and Hodges 1987). However, stems greater than 10 in. in diameter exhibit a markedly reduced sprouting potential. Consequently, in stands composed primarily of large trees, stump sprouts are not a reliable source of regeneration, so that a build-up of advance reproduction in size and number over a period of years appears to be necessary before final harvest (Johnson 1979, Loftis 1990). These larger seedlings compete more successfully with other vegetation when the overstory is removed and show good sprouting potential if severed (Janzen and Hodges 1987).




Figure 1—Cross-section of a major and minor stream valley showing relationship of relief features and typical species association (Hodges and Switzer 1979).

A key to establishing and developing advance reproduction for natural regeneration is in the regulation of light reaching the forest floor (Hodges 1989). Too much or too little light can have detrimental effects on regeneration efforts. Full sunlight may promote faster-growing species rather than oak, while limited light will benefit more shade-tolerant species. In the absence of fire and grazing, most bottomland stands have dense midstories and understories of tolerant species which hinder the regeneration of oak. Research and experience indicate that failure to treat this undesirable and unmerchantable component causes reduced stocking and growth of oak regeneration (Deen and others, in press).

Johnson and Deen (in press) developed a numerical rating system to evaluate the potential for successful regeneration of southern bottomland hardwood stands. The system is based on the number and size of advance reproduction as well as on stump and seedling sprout potential. If regeneration potential is adequate, the stand can be harvested. If not, additional sources of regeneration must be secured naturally by manipulation of the overstory and/or lower canopies to allow development of advance reproduction, or by artificial means, such as direct-seeding or planting. Similar evaluations of regeneration potential have been developed for Page 249 of 404

the Northeast (Marquis and Bjorkham 1982) and the Midwest (Sander and others 1976).

Tables that show the occurrence, shade tolerance, flood tolerance, and reproductive characteristics of many bottomland species including oaks have been synthesized from the literature by McKnight and others (1981). The bottomland oak components of these tables are shown in table 1. An understanding of these species-site relationships and reproductive strategies is helpful when making decisions on oak regeneration and management.

Table 1-Occurrence, shade and flood tolerance, and reproductive characteristics of the bottomland oak species¹

Common and scientific name	Occurrence in bottomlands	Shade tolerance	Flood tolerance	Reproductive characteristics
Cherrybark oak (Quercus falcata var. pagodifolia Ell.)	Widely on best loamy sites on all river bottom ridges and better drained creek bottoms and hummocks. Mostly on older alluvium. Occasionally on tight, silty clay, but grows poorly there.	Moderately intolerant to intolerant.	Weakly tolerant to intolerant. Seed- lings can withstand very little flood- ing. Viability of acorns greatly re- duced by submer- sion.	Seed dispersed Sept Dec. by gravity, birds, and animals— seldom by water. Seedlings may start in shade, but cannot survive long. Not a good stump sprouter.
Nuttall oak (<i>Q. nuttallii</i> Palmer)	Widely on flats, low ridges, shallow sloughs, and near margin of swamps, in recent alluvial sites. On older alluvium mainly restricted to wet, heavy but not impervious sites. Restricted to major strear entering the Gulf and the large tributaries.		Moderately tolerant. Seedlings killed by high water during growing season. Viability of acorns not reduced by 34 days of submergence.	Starts readily in either shade or openings but soon dies in shade. Persists against all ground cover ex- cept heavy pep- pervine. Many large trees are of sprout origin.
Shumard oak (<i>Q. shumardii</i> Buckl.)	Mainly on terraces in older alluvium and out- wash from upland and well drained creek bottoms and hummocks. Rare on newer soils. Widely distributed but scattered.	Intolerant.	Weakly tolerant. Seedlings rela- tively intolerant to flooding.	Seed dispersed Sept Dec. by gravity and animals, largely by squirrels and rarely by water. Establishes as scattered individ- uals in shade or openings. A poor sprouter.
Swamp chestnut oak (cow oak) (Q. michauxii Nutt.)	Common in large creek bottoms and hummocks on best, well drained loamy ridges. Occa- sionally on a wet, silty clay, high flat. Rarely on best, most mature recent alluvium, but typically a tree of older alluvium sites.	Moderately intolerant.	Weakly tolerant. Seedling intolerant to flooding.	Seed dispersed Oct Dec. by gravity and animals, primarily squirrels. Seed germ- inates soon after seed fall. Best seedbed is moist and well- drained with a light cover of leaves. Seedlings require full sunlight for best development. Small stumps sprout well.

Common and scientific name	Occurrence in bottomlands	Shade tolerance	Flood tolerance	Reproductive characteristics	
Water oak (Q. nigra L.)	Widely on loam ridges in first bottoms and on any ridge and silty clay flats in second bottoms or terraces.	Intolerant.	Weakly to moderately tolerant. Prolonged submergence of seed- lings during growing season will kill the trees.	Seed dispersed Sept. Nov. by gravity, birds, animals, and water. Seedlings establish best on moist, well-aerated soil. Small stumps sprout readily.	
Willow oak (Q. phellos L.)	Widely on ridges and high flats of major streams. Less common in creek bottoms. May form nearly pure stands of poor quality on hardpan terrace soils. Grows best on flats of old alluvium and on clay loam ridge of new alluvium.	Intolerant.	Weakly to moderate- ly tolerant. Seedlings among oaks are one of the more tolerant to water, but pro- longed submergence during growing season is fatal.	Seed dispersed Oct Dec. by gravity, birds, animals and water. Good seedling crops not unusual. Stumps under 12 in. in diameter sprout well.	

Table 1—Occurrence, shade and flood tolerance, and reproductive characteristics of the bottomland oak species¹—Continued

¹ Adapted from McKnight and others 1981.

SILVICULTURAL SYSTEMS AND METHODS OF REPRODUCTION

Silvicultural systems integrate harvesting, regeneration, and intermediate operations in an orderly process for managing forest stands. Silvicultural practices are traditionally divided into two systems: even-aged and uneven-aged. A third system, two-aged, has been used with some success with Appalachian oaks (Beck 1987, Sims 1992). Although there are no published studies, the two-aged system also may have some application with bottomland oaks. The methods of reproduction employed under even-aged silviculture include seed tree, shelterwood, and clearcutting. Single-tree and group selection are methods of reproduction under uneven-aged silviculture. A discussion of each of these various methods of reproduction in relation to regeneration of bottomland oaks follows. A brief synopsis of the use of artificial regeneration to supplement natural regeneration of oaks is also presented.

Regeneration by the Seed-Tree Method In the seed-tree method of reproduction, a small number of trees are retained to provide seed after harvest. This method favors wind-blown, light-seeded species (Toliver and Jackson 1989) and not heavy-seeded species, such as oak, in which seeds are dispersed for relatively short distances from the parent tree by gravity, birds, rodents, or sometimes water. Because most bottomland oak regeneration develops from advance reproduction present at the time of cutting or from the sprouting of stumps, there is little need or justification for seed trees (Johnson and Krinard 1976). Twenty-nine years after a seed-tree cut in bottomland sweetgumred oak stands in Arkansas, Johnson and Krinard (1988) reported that the resulting stands were essentially similar to those produced through clearcutting. Most of the oak regeneration developed from advance reproduction and stump sprouts. Seed trees did not significantly contribute to the establishment of new seedlings. Regeneration by the Shelterwood Method

The shelterwood method of reproduction is used to enhance the development of large oak seedlings prior to the final harvest cut. The classical shelterwood method usually employs a minimum of three cuts (Smith 1986):

(1) A preparatory cut of lower-crown-class trees to prepare a suitable seedbed.

(2) One or a series of partial cuts to encourage development of existing small seedlings.

(3) A removal cut of the overstory to release the advance reproduction.

Experience with oaks has indicated that the preparatory cut generally does not result in the establishment of new oak seedlings (Janzen and Hodges 1987, Loftis 1990). Although oak seedlings may appear after an occasional bumper crop of acorns, most seedlings become established as advance reproduction only over a period of years (Beck 1977). The shelterwood method for oak regeneration is primarily used to encourage the development of large seedlings from existing smaller ones, so that they will be able to successfully compete with other vegetation when the overstory is removed (Hodges and Janzen 1987, Loftis 1990).

Large red oak advance reproduction has been successfully cultured with shelterwood treatments in the Southern Appalachians (Loftis 1990). However, results with bottomland oaks have been highly variable. Hodges (1989) reported that, in practice, heavy shelterwood cuts favor the development of fast-growing, intolerant species rather than the oaks, while lighter cuts may encourage more tolerant and less desirable species. Factors contributing to the variable success of the shelterwood method to regenerate bottomland oaks as compared to oak regeneration in the Southern Appalachians include (1) higher site productivities; (2) a preponderance of tolerant species such as sugarberry, boxelder, beech, elms, maples, hickories, hornbeam, and hophornbeam; and (3) the unique hydrologic regime of bottomlands.

Shelterwood is the most flexible of all the methods of reproduction (Toliver and Jackson 1989); however, the timing of various cuts is critical for the continued development of oak reproduction. Treatment of the undesirable midstory and understory vegetation by herbicide or cutting is essential for the development of oak seedlings (Janzen and Hodges 1987, Loftis 1990). More research is needed before recommendations can be made on the shelterwood method of regenerating bottomland oaks, particularly in terms of the degree of competition control necessary, the number and intensity of cuts, and the interval between cuts.

Regeneration by the Clearcutting Method The clearcutting method of reproduction involves the removal, in a single cut, of all trees larger than seedlings. However, this "complete clearcut" is different from a "commercial clearcut" in which unmerchantable stems that may interfere with the future development of regeneration are not removed. The primary advantage of the clearcutting method is that it provides the sunlight required for the development and growth of moderately intolerant to intolerant species such as oak (Kellison and others 1988).

> Clearcutting is the most proven method of successfully regenerating bottomland oaks (Johnson 1979); however, there are qualifications. The success of clearcutting in regenerating bottomland oaks is dependent on the presence of adequate advance reproduction in the stand prior to cutting and the development of stump sprouts Page 252 of 404

following cutting. Large seedlings of advance reproduction are essential to successful oak regeneration. The thousands of small oak seedlings that frequently occur in older stands simply cannot compete with faster-growing species upon release. Clearcutting has been successful in managed stands where one or more earlier thinnings have resulted in the development of large seedlings (Hodges 1989). Most oak regeneration failures following clearcutting are due to the lack of large, well-established advance reproduction and/or stumps capable of producing oak sprouts.

When clearcutting to regenerate oaks in bottomlands, the harvest should take place during the dormant season to maximize stump sprouting (Kellison and others 1988). Control of residuals either by herbicides or cutting is advocated, especially after a commercial clearcut (Golden and Loewenstein 1991), to reduce competition from undesirable midstory and understory species (Janzen and Hodges 1987). Clearcutting is also recommended where tree quality and stocking are poor (such as in high-graded stands) and there is little potential to upgrade the stand (Kellison and others 1988).

Clearcutting tends to initially favor the establishment and growth of shadeintolerant, fast-growing, light-seeded species, but species composition will gradually shift toward more mid-tolerants over time (Toliver and Jackson 1989). Oaks are often present, but typically inconspicuous, in the jungle of woody and nonwoody vegetation found during the early stages of development following clearcutting. However, two separate research studies on oak stand development have shown that bottomland oaks, although inconspicuous in the regeneration, eventually became prominent in the mature stand (Clatterbuck and Hodges 1988, Johnson and Krinard 1988).

Regeneration by the Single-Tree Selection Method

Single-tree selection involves cutting trees in all size classes to achieve a preplanned distribution of diameters and to create stands with an uneven-aged structure. This method is best-suited for shade-tolerant species that have the ability to regenerate and grow in the shade, such as beech and maple in the Northeast. When single-tree selection is continually applied to stands containing intolerant species, composition will gradually shift to more tolerant species (Johnson and Krinard 1989). When this method of reproduction is applied to mature bottomland oak stands, it usually leads to the development of future stands dominated by tolerant species, such as sugarberry, boxelder, hickories (pecan), elms, maples, and bays.

High-grading and diameter-limit cutting are often touted as single-tree selection. These abusive cutting practices of "taking the best and leaving the rest" are not true regeneration methods and are not to be construed as variations of the single-tree selection method of reproduction. These abusive cutting practices are not directed toward obtaining regeneration, and cutting does not occur in all size classes to maintain an uneven-aged structure. The assumption of taking the big trees to allow the little trees to grow perpetuates the development of advanced-aged, poorly formed, shade-tolerant trees that result in a progressively less-valuable stand.

There are few commercially valuable shade-tolerant bottomland species. Most bottomland oaks range from intolerant to intermediate in tolerance of shade Page 253 of 404

(Putnam and others 1960) and require some direct sunlight for successful seedling establishment and continued development. The light available to the forest floor in the small openings created by single-tree harvesting simply is not enough to ensure successful regeneration of oaks. Therefore, the single-tree selection method of reproduction is not recommended for bottomland oaks (Toliver and Jackson 1989).

Regeneration by the Group Selection Method

The group selection method of reproduction involves the removal of trees in small groups. The textbook interpretation (Smith 1986) is that cutting in small groups is regulated through volume control within size classes, following the uneven-aged system of silviculture. Size of opening generally ranges from 0.1 to 0.5 acre. The small openings created during these cuts usually fail to allow sufficient light to the forest floor for satisfactory establishment and development of bottomland oak reproduction. In bottomland forests, the trees that are able to regenerate in these small openings are usually of either a commercially undesirable understory species or a shade-tolerant species of low potential value.

Patch cutting is a combination of uneven-aged (group selection) and even-aged (clearcutting) silviculture where larger groups of 0.5 to 5 acres (usually 1 to 3 acres) are removed (Marquis 1989). An uneven-aged stand condition is produced by patch cutting consisting of many small, variously shaped, even-aged groups or patches within the stand. Management of timber is regulated by forest area. The forest stand under patch cutting exhibits "uneven-aged characteristics even though it is managed as a collection of small, even-aged groups" (Marquis 1989). Light available to the forest floor in these larger openings is sufficient to successfully regenerate bottomland oaks. The Tennessee Division of Forestry uses patch cutting as one management technique on its State Forests (Applegate 1991).

Anderson-Tully Company of Memphis, TN, successfully uses variations of group selection and patch cutting in its management of bottomland oak forests. Anderson-Tully practices classical uneven-aged silviculture (Smith 1986) by maintaining at least three age classes in its stands and by using volume regulation of the cut. The company's biggest challenge has been the regeneration of oaks under the uneven-aged system (Stephano 1992). Larger openings are used to provide optimum conditions for oak regeneration. This company also practices intensive control of midstory vegetation to further enhance establishment and development of oak reproduction.

Group selection requires a "hands-on" intensive silvicultural approach. Regulation of stand structure and development is complicated, time-consuming, and expensive. If used judiciously, however, group selection can produce the biological conditions necessary to allow the successful establishment and further development of bottomland oak reproduction.

Artificial Regeneration Because large advance reproduction of oak seedlings is necessary for successful natural regeneration on bottomland sites, planning for regeneration is critical prior to harvest. When advance reproduction is insufficient, several to many years may be required to build up the amount and size of reproduction needed before harvest. Artificial regeneration can be used when advance reproduction is lacking and when the 5 to 10 years needed to establish advance reproduction are not available.

Artificial regeneration of oaks through direct-seeding, plantation culture, or enrichment/supplemental planting is discussed elsewhere in this symposium. A bulletin by Allen and Kennedy (1989) is an excellent source of reforestation information for bottomland hardwoods.

SUMMARY AND RECOMMENDATIONS

Natural oak regeneration in the bottomlands has been inconsistent. Most regeneration failures have been attributed to lack of advance oak reproduction. In general, advance reproduction of oaks must be secured in both sufficient size and number before the final harvest cut to successfully regenerate oaks on bottomland sites. The key consideration is to create favorable light conditions on the forest floor prior to final harvest. Control of the shade-tolerant competition in the midstory and understory is required for culturing bottomland oak reproduction. A pretreatment regeneration evaluation for bottomland oaks is necessary. When sufficient advance oak reproduction and sprout potential are present, a complete harvest or clearcut of all stems will usually regenerate the stand to oak.

Most stands range from closed-canopy stands with little advance reproduction to cutover, mismanaged, or unmanaged areas where species composition and tree quality are poor. In theory, the shelterwood method should nurture oak reproduction if oak seed sources are present. However, in practice, a heavy shelterwood cutting that creates gaps in the overstory usually favors the development of faster-growing intolerant species rather than oak, while in other instances shelterwood cutting actually encourages the growth of undesirable tolerant species already established in the midstory and understory. Both scenarios exclude optimum oak regeneration. Application of the shelterwood method for regenerating bottomland oaks has been discouraging with many more failures than successes. Single-tree selection also is not recommended for regenerating bottomland oaks because this method favors the growth and establishment of shade-tolerant species. Group selection can be used to regenerate bottomland oaks if the size of the opening is adequate to provide sufficient light to the forest floor to encourage the establishment and development of oak reproduction.

Figure 2 shows a generalized prescription for regenerating bottomland oaks. The pretreatment regeneration evaluation of figure 2 follows that presented in this symposium by Johnson and Deen (in press). No cookbook treatment for bottomland oak regeneration can totally account for the numerous biotic, abiotic, and anthropogenic factors which affect oak regeneration potential (e.g., site, species mixtures, past disturbances, flooding, fire, grazing, etc.). Practitioners interested in perpetuating oaks in the bottomlands need to evaluate existing stand and site conditions, the ecological requirements of the species, and the various prescriptions that are suitable for not only the establishment but the continued development of bottomland oaks.

Pre-treatment Regeneration Evaluation




Figure 2-Regeneration procedures for bottomland oaks.

LITERATURE CITED

Allen, James A.; Kennedy, Harvey E., Jr. 1989. Bottomland hardwood reforestation in the lower Mississippi valley. Stoneville, MS: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station, and Slidell, LA: U.S. Department of the Interior, Fish and Wildlife Service, National Wetlands Research Center. 28 pp.

- Applegate, H.W. 1991. Group selection---An alternative for management of small, privately owned woodlands in Tennessee. Nashville, TN: Tennessee Department of Agriculture, Division of Forestry. 9 pp.
- Baker, James B.; Broadfoot, W.M. 1979. A practical field method of site evaluation for commercially important southern hardwoods. Gen. Tech. Rep. SO-26. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station, 51 pp. Page 256 of 404

- Beck, Donald E. 1977. Twelve-year acorn yield in southern Appalachian oaks. Res. Note SE-244. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 8 pp.
- Beck, Donald E. 1987. Management options for southern Appalachian hardwoods: the two-aged stand. In: Phillips, Douglas R., comp. Proceedings of the fourth biennial southern silvicultural research conference; 1986 November 4-6; Atlanta, GA: Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 451-454.
- Burns, Russell M.; Honkala, Barbara H., tech. coord. 1990. Silvics of North America: 2. Hardwoods. Agric. Handbook 654. Washington, DC: U.S. Department of Agriculture, Forest Service. 877 pp.
- Clatterbuck, Wayne K.; Hodges, John D. 1988. Development of cherrybark oak and sweetgum in mixed, even-aged bottomland stands in central Mississippi, U.S.A. Canadian Journal of Forest Research. 18(1):12-18.
- Deen, R.T.; Hodges, J.D.; Janzen, G.C. [in press]. The influence of midstory and understory control treatments on regeneration of hardwoods following clearcutting. In: Proceedings of the seventh biennial southern silvicultural research conference; 1992 November 17-19; Mobile, AL. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station.
- Golden, Michael S.; Loewenstein, Edward F. 1991. Regeneration of tree species 7 years after clearcutting in a river bottom in central Alabama. In: Coleman, Sandra S.; Neary, Daniel G., comps., eds. Proceedings of the sixth biennial southern silvicultural research conference; 1990 October 30-November 1; Memphis, TN. Gen. Tech. Rep. SE-70. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 76-83.

Hodges, John D. 1989. Regeneration of bottomland oaks. Forest Farmer. 49(1):10-11.

- Hodges, John D.; Janzen, Greg. 1987. Studies on the biology of cherrybark oak: recommendations for regeneration. In: Phillips, Douglas R., comp. Proceedings of the fourth biennial southern silvicultural research conference; 1986 November 4-6; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 133-139.
- Hodges, John D.; Switzer, George L. 1979. Some aspects of the ecology of southern bottomland hardwoods. In: North America's forests: Gateway to opportunity: 1978 Joint Convention of the Society of American Foresters and the Canadian Institute of Forestry; 1978 October 22-26; St. Louis, MO. Washington, DC: Society of American Foresters: 360-365.
- Janzen, G.C.; Hodges, J.D. 1987. Development of advanced oak regeneration as influenced by removal of midstory and understory vegetation. In: Phillips, Douglas R., comp. Proceedings of the fourth biennial southern silvicultural research conference; 1986 November 4-6; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 455-461.
- Johnson, Robert L. 1979. Adequate oak regeneration a problem without a solution. In: Management and utilization of oak: Proceedings of the 7th annual hardwood symposium of the Hardwood Research Council; 1979 May; Cashiers, NC. Asheville, NC: Hardwood Research Council: 59-65.
- Johnson, R.L.; Deen, R.T. [in press]. Prediction of oak regeneration in bottomland forests. In: Proceedings of the symposium on oak regeneration: Serious problems—practical recommendations; 1992 September 8-10; Knoxville, TN.
- Johnson, R.L.; Krinard, R.M. 1976. Hardwood regeneration after seed tree cutting. Res. Paper SO-123. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 9 pp.

- Johnson, Robert L.; Krinard, Roger M. 1988. Growth and development of two sweetgum-red oak stands from origin through 29 years. Southern Journal of Applied Forestry. 12(2):73-78.
- Johnson, Robert L.; Krinard, Roger M.. 1989. Survival and growth of Nuttall oak seedlings following selection cutting. Southern Journal of Applied Forestry. 13(1):43-46.
- Kellison, R.C.; Martin, J.P.; Hansen, G.D.; Lea, R. 1988. Regenerating and managing natural stands of bottomland hardwoods. Bulletin APA 88-A-6. Washington, DC: American Pulpwood Association. 26 pp.
- Loftis, David L. 1990. A shelterwood method for regenerating red oak in the southern Appalachians. Forest Science. 36(4):917-929.
- Marquis, David A. 1989. Alternative silvicultural systems east. In: Silvicultural challenges and opportunities in the 1990's: Proceedings national silviculture workshop; 1989 July 10-13; Petersburg, AK. Washington, DC: U.S. Department of Agriculture, Forest Service: 36-43.
- Marquis, David A.; Bjorkbom, John C. 1982. Guidelines for evaluating regeneration before and after clearcutting Allegheny hardwoods. Res. Note NE-307. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 4 pp.
- McKnight, J. Sid; Hook, Donal D.; Langdon, O. Gordon; Johnson, Robert L. 1981. Flood tolerance and related characteristics of trees of the bottomland forests of the southern United States. In: Clark, J.R.; Benforado, J. eds. Wetlands of bottomland hardwood forests: Proceedings of a workshop on bottomland hardwood forest wetlands of the southeastern United States; 1980 June 1-5; Lake Lanier GA. New York: Elsevier Science Publishing Co.: 29-69.
- Putnam, John A.; Furnival, George M.; McKnight, J.S. 1960. Management and inventory of southern hardwoods. Agric. Handbook 181. Washington, DC: U.S. Department of Agriculture, Forest Service. 102 pp.
- Sander, Ivan L.; Johnson, Paul S.; Watt, Richard F. 1976. A guide for evaluating the adequacy of oak advance reproduction. Gen. Tech. Rep. NC-23. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 7 pp.
- Sims, Daniel H. 1992. The two-aged stand: a management alternative. Management Bulletin R8-MB61. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 2 pp.
- Smith, David Martyn 1986. The practice of silviculture. 8th Edition. New York: John Wiley and Sons, Inc. 527 pp.
- Stephano, Peter J. 1992. Logging along the mighty Mississippi. Wood Magazine. 62-67 pp.
- Toliver, John R.; Jackson, Ben D. 1989. Recommended silvicultural practices in southern wetland forests. In: Hook, Donal D.; Lea, Russ, eds. Proceedings of the symposium: The forested wetlands of the Southern United States; 1988 July 12-14; Orlando, FL. Gen. Tech. Rep. SE-50. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 72-77.

Relative Growth of Oaks and Pines in Natural Mixtures on Intermediate to Xeric Piedmont Sites

F. Thomas Lloyd and Thomas A. Waldrop, Southeastern Forest Experiment Station, USDA Forest Service, Clemson, SC 29634

ABSTRACT Increment cores and stem analyses from a growth and yield study in naturally regenerated pine-hardwood stands in the Piedmont were used to compare relative growth of oaks, pines, and other non-oak hardwoods on intermediate to xeric sites in that region. The primary focus in this paper is on the relative growth of only the oak and pine components. The stem analyses show that pines early in stand life grow faster in height than the oaks (a widely observed result), reaching an average maximum height advantage of 20 ft. by an average age of 32 years. Beyond this age, the stem analyses showed annual pine height growth slowing dramatically, falling below the rather steady 2 ft. per year observed for the oaks through age 70, thus reducing the average 20-ft. cumulative height advantage of pine was cut in half, to 9.6 ft. We show that oaks attain basal area growth comparable to that of pines as early as age 15, and that beyond age 15, the oak growth advantage increased through stand age 70, outgrowing the pines by 70 percent between ages 60 and 70.

INTRODUCTION Pine-hardwood mixtures are common in the Piedmont physiographic regions of Virginia, North Carolina, South Carolina, and Georgia. Unpublished data from the Forest Inventory and Analysis (FIA) Unit of the Southeastern Forest Experiment Station show that 1/3 of the Piedmont forest—7.1 million of 22 million acres—is in stands where 30 to 90 percent of the total basal area is in hardwoods, or 10 to 70 percent in pines. Although loblolly pine (*Pinus taeda* L.) is the dominant pine species, other yellow pine species are included in the estimates.

On these intermediate to xeric Piedmont sites, it is well known that pine growth exceeds that of oaks and other hardwoods early in stand life. This dramatic pine growth advantage is why we often hear these areas described as "pine sites." At the same time, oak coppice and advanced regeneration do well on these sites when they are present and are not aggressively controlled during site preparation. Oaks also outlive pines on these dry sites. Barring major disturbance, therefore, the oak component normally increases as stands on these sites age. In fact, it is common on eroded Piedmont sites for mortality of dominant and codominant pines to begin as early as age 40, thus speeding the composition toward oaks. Jones (1991) in his research on landscape ecosystem classification found oaks abundant in the late-successional stands he studied in the Piedmont. He also found that the oak

species associations were indicators of a site-quality gradient. In summary, oaks regenerate naturally and persist on these sites. Their survival is site-specific, but they can generally be thought of as well-adapted to these sites.

Given this suitability of oaks for these sites, it is logical to ask how well they grow in long rotations relative to pines and non-oak hardwoods. Relative growth dynamics of the pine, oak, and non-oak species groups have not been examined for sawtimber rotations of naturally regenerated mixtures of pines and hardwoods. One objective of a major study we are installing, and the major focus of this paper, is to examine these growth dynamics.

METHODS A growth and yield study in naturally regenerated mixtures of hardwood and pine has been initiated to study growth dynamics (Lloyd 1991). Fifty circular, 1/5-acre permanent plots measured in the first phase of this study form the dataset for this paper. Sampled stands contained from 93 to 182 sq. ft. of basal area per acre in merchantable and unmerchantable trees, and stand ages ranged from 20 to 79 years. Twenty one plots are on the Piedmont Ranger Districts of the Sumter National Forest and 29 are on the Clemson University Experimental Forest.

Diameters at breast height were measured on all trees, and merchantable sized trees (4.6+ in.) were tagged and mapped by azimuth and distance from plot center. Separate samples of hardwood and pine trees covering the diameter range on each plot were selected for measurement of total height. Although growth and mortality ultimately will be estimated by remeasuring these plots, recent growth was calculated from increment cores taken from merchantable-sized trees. Radial growth for the last 5 and 10 years was measured with a Bannister incremental measuring instrument. Radial growth data were used to estimate basal area of trees 5 and 10 years prior to plot establishment. Only survivor growth can be studied in this way.

Ten-year basal area growth of surviving trees was estimated as the difference between basal area of merchantable trees at measurement time and the calculated basal area of the same trees 10 years earlier. It was divided into components for pines, oaks, and non-oak hardwoods, and separate prediction models were developed for each species component. The same model form was fit to all groups. The predictor variables screened for these models were: (1) initial merchantable basal area in the given species group, (2) the species group's basal area as a proportion (ratio) of the total merchantable basal area, (3) the reciprocal of stand age, and (4) the cross-products of (1), (2), and (3).

Cumulative height growth differences were examined next. It has been widely observed that early height patterns favor pine. The goal here was to examine cumulative height over a longer time period. These height data were obtained by analyzing stems of pairs of one dominant or codominant oak and one such pine located near (not in) each permanent plot. Thus, two height/age curves were plotted for each plot. Each potential stem analysis tree was cored prior to felling to determine age and to seek evidence of previous suppression of growth. Trees with previous suppression were excluded. Substitutes were examined in the same way until a free-to-grow tree was found. Finding suitable trees was not difficult; we rarely had to go beyond the first choice. The resulting cumulative height/age

curves were used to examine height-over-age patterns. In order to assess long-term patterns, the cumulative height data set was screened to include only plots in stands over 54 years old. Twenty plots (40 trees) met this criterion.




Figure 1—An example of a typical cumulative height pattern observed in stem-analyzed pairs of oaks and pines located near each of 50 permanent growth and yield plots in naturally regenerated hardwood and pine mixtures in the Piedmont physiographic region. The sample pine was always a loblolly, and the oak could be a dominant or codominant tree of good form from any of the oak species found on these sites.

The first step was to examine graphs of the height/age data for the pairs of oaks and pines on each plot. Figure 1 illustrates the predominant pattern observed for all plots. Because we estimated height at the end of every growing season by interpolation, it was easy to compute the height difference between oaks and pines over the life of the stand. Since pines where generally taller, we arbitrarily subtracted oak height from pine height, and then fit to each of the 20 data sets the simple quadratic polynomial expression

$$H_{d} = c_{0} + c_{1}t + c_{2}t^{2}$$
(1)

where H_d is the difference in cumulative height between the pine and oak at an each age (t). We then used the estimates of c_0 , c_1 , and c_2 and the calculus of extreme values to estimate for each of the 20 plots the age at which the difference between pine and oak height was maximum and what the maximum difference was at that time in stand life.

The final analytical procedure looked at the average shape of the oak cumulative height pattern. All height/age data for the 20 sampled oaks were pooled into a single data set, and then the model

$$\mathbf{h}_{\mathbf{0}} = \mathbf{e}^{\mathbf{a}}\mathbf{t}^{\mathbf{b}} \tag{2}$$

was fit using nonlinear least squares. The variable h_0 is cumulative oak height at age t, and a and b are the model parameters to be estimated.

RESULTS The model used to predict 10-year pine basal area growth
$$(b_P)$$
 is
 $b_P = g_0 + g_1 B_P + g_2 P_P + g_3 B_P/t$ (3)

where B_{P} is the initial basal area of presently surviving pines as it occurred 10 years prior to plot installation, P_P is the proportion of total initial merchantable basal area (B) represented by pines (that is, $P_P = B_P/B$), and t is initial stand age. There are similar models for oaks (b_0) and non-oak hardwoods (b_N) , where

and

$$b_0 = g_0 + g_1 B_0 + g_2 P_0 + g_3 B_0/t$$
 (4)

$$b_N = g_0 + g_1 B_N + g_2 P_N + g_3 B_N/t.$$
 (5)

To further illustrate the definitions of the independent variables, the following relationships hold across the three models: $B_P + B_O + B_N = B$

and

$$P_{\rm P} + P_{\rm O} + P_{\rm N} = 1.$$
 (7)

(6)

The \mathbb{R}^2 statistics of fit for Equations (3), (4), and (5) above are 0.91, 0.75, and 0.77, respectively. R² values of 0.8 for regression models of periodic growth are considered good. The corresponding estimates of the model parameters $(g_0, g_1, g_2, g_2, g_3)$ g₃) are (-3.079134, -0.21006, 32.38342, 6.670954) for pines, (1.879838, -0.0051275, 8.509635, 5.819824) for oaks, and (0.4226926, 0.1848599, -9.917444, 2.909263) for non-oak hardwoods. These estimated parameters were used in the appropriate model to predict components of 10-year growth for selected values of initial stand ages listed in table 1. It is not the goal in table 1 to predict for the actual initial stand conditions, but rather, to compare relative oak and pine growth performance. For this reason, the same set of P-values was used for each initial age, that is, $P_P = 0.4$, $P_O = 0.4$, and $P_N = 0.2$.

Table 1-Predicted periodic (10 years) basal area growth of survivors at four ages in pine-hardwood mixtures which have merchantable basal area composed of 40 percent pine, 40 percent oak, and 20 percent other hardwoods

	Initial stand age ¹						
Species group	15		45	60			
	ft. ² /a						
Pine	17.4	10.3	7.4	5.5			
Oak	17.5	12.1	10.3	9.3			
Other hHardwood	4.5	3.5	3.4	3.6			
Total	39.4	25.9	21.1	18.4			

¹ Total merchantable basal area at the beginning of each period was 80 ft.² at age 15, 90 ft.² at age 30, 100 ft.² at age 45, and 110 ft.² at age 60.

These proportions are all within the ranges observed in the data. Fixing these proportions permits direct comparisons because the oak and pine predictions of basal area growth come from the same initial basal area.

Table 1 shows that by age 15, the oaks were producing as much basal area growth as the pines for the same initial basal area. As the stands aged, the oak increasingly outgrew the pines in basal area through age 70. At that time, oaks were growing 70 percent more per 10-year period than the pines. The footnote for table 1 explains how the corresponding values of initial basal area were calculated. For example, for stand age 30, the observed data averaged about 100 sq. ft. of initial merchantable basal area (that is, B). Since P_P and P_o are both set equal to 0.4, the initial basal area components were 40 sq. ft. for both the oaks and pines. Thus, table 1 shows that for initial stand age 30, the 40 sq. ft. of oak grew 12.1 sq. ft. of basal area in the next 10 years (that is, from age 30 to age 40), while the 40 sq. ft. of pine basal area grew 10.3 sq. ft. in the same period. This growth advantage of the oaks increased with increasing initial stand age.

Figure 1 illustrates how pines on these Piedmont sites outgrow oaks. From Equation 1, we found that the average stand age of maximum pine/oak height difference was 32 years, and the average height difference at that point was 20.0 ft., with a quartile range of 17 to 22 ft. However, by stand age 55, the average pine/oak cumulative height difference was cut in half, to 9.6 ft. The pattern of dramatically slowing pine height growth after age 30 and steady oak height held generally across plots. It should be kept in mind that our working definition of pine-hardwood mixtures is not a closed pine overstory with a hardwood understory. We only work with stands in which the pine component is sparse enough to allow some light from above for the largest hardwood, even though they are shorter than the pines.

Visual examination of the pooled oak height data suggested a steady growth rate through stand age 70. We examined this average trend by fitting Equation (2) to the cumulative height data for the 20 plots that were 55 + years old. The nonlinear least squares estimates of the model parameters were 0.71 and 0.98 for a and b, respectively, and the value of e^a was 2.03. Since b was nearly equal to 1, the analysis suggests a rather steady 2 ft. of height growth per year. This average 2-ft. rate was suggested independently in another study (Geisinger and others 1989) of pine and hardwood regeneration. Table 2 shows oak height growth of 3.5 and 3.4

Growing season	Pine	Oak	Hickory	Blackgum	Other hardwood
1988	1.0 ¹	3.5	ft 1.7	2.8	4.0
1988	1.5	6.9	3.8	2.8 5.0	4.0 7.3
1990	3.3	8.9	5.2	5.9	9.1
1991	7.5	11.1	7.0	7.4	11.2

Table 2—Average heights of five species groups after four growing seasons for the winter-fell, no-burn treatment of the pine-hardwood regeneration study.

¹ Average height of seedlings planted in the 1988-1989 growing season.

ft. per year during the first two growing seasons, followed by a slowing to around 2 ft. per year the third and fourth growing seasons. Table 2 also reinforces the pine height growth pattern of lagging for around 3 years, and then dramatically accelerating. In this case, the pine grew 4.2 ft. in the fourth growing season.

CONCLUSIONS As stated numerous times in this symposium, getting oak regeneration on xeric to intermediate sites is not hard when oak root stocks and advanced reproduction are present in clearcut stands. The data re-emphasize that after a growth lag, pines on these sites clearly outgrow oaks early in stand development. However, a longer-term look at height development tells a different story. It shows how pine growth slows dramatically after age 30, with oak rapidly cutting the height deficit. Although we do not have basal area growth data for very young mixed stands, this analysis shows oak basal area growth equaling that of pine by age 15. From that point, oak basal area growth increasingly surpasses the pines, reaching a 70-percent advantage between ages 60 and 70.

Given the increasingly important values of oaks for aesthetics and wildlife, the increasing stumpage prices for high-quality oaks, and the growing markets for low-grade oaks, managers of relatively dry upland Piedmont sites should take a close look at our results. We know that oaks are ecologically suited for these sites because they regenerate and live long lives there. Their growth performance in relation to pines is not impressive early in life, but our data indicate that they catch up later. Thus, they would appear to be sensible choices for sawtimber rotations.

LITERATURE CITED

- Geisinger, Donn R.; Waldrop, Thomas A.; Haymond, Jacqueline L.; Van Lear, David H. 1989. Sprout growth following winter and spring felling with and without summer broadcast burning. In: Waldrop, Thomas A., ed. Proceedings of Pine-Hardwood Mixtures: A Symposium on Management and Ecology of the Type; 1989 April 18-19; Atlanta, GA: 91-94. Gen. Tech. Rep. SE-58. Asheville, NC: Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 271 pp.
- Jones, Steven M. 1991. Landscape ecosystem classification for South Carolina. In: Mengel, Dennis L.; Tew, D. Thompson, eds., Proceedings: Ecological Land Classification: Applications to Identify the Productive Potential of Southern Forests; 1991 January 7-9; Charlotte, NC: 59-68. Gen. Tech. Rep. SE-68. Asheville, NC: Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 149 pp.
- Lloyd, F. Thomas. 1991. Forecasting growth of pine-hardwood mixtures from their ecological land class. In: Mengel, Dennis L.; Tew, D. Thompson, eds., Proceedings: Ecological Land Classification: Applications to Identify the Productive Potential of Southern Forests; 1991 January 7-9; Charlotte, NC: 93-95. Gen. Tech. Rep. SE-68. Asheville, NC: Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 149 pp.

Regenerating Northern Red Oak on High-Quality Sites in the Southern Appalachians

David L. Loftis, Southeastern Forest Experiment Station, USDA Forest Service, Asheville, NC 28806

ABSTRACT On high-quality sites in the Southern Appalachians, dominant and codominant oak stems will be present in regenerated stands only when adequate regeneration sources exist in the mature stands to be harvested. These regeneration sources may be present in mature stands as advance reproduction or as stump sprout potential. Since oak trees tend to be larger and fewer in number on high-quality sites than on lower quality sites, and since large stumps are less likely to produce viable sprouts than are smaller stumps, advance reproduction is the most important oak regeneration source on high quality sites. The potential for advance oak seedlings to become dominant or codominant stems in the next stand increases with seedling size. Thus, the expected contribution of trees (as stump sprouts) and advance reproduction are both related to size. Methods are currently available to assess oak regeneration potential on high-quality sites in the Southern Appalachians. If the oak regeneration potential (and the potential for other species) in an existing stand is sufficient to meet management objectives, the overstory can be removed in one or more harvest cuts. If, however, the oak regeneration potential is not sufficient to meet management objectives, oak regeneration potential must be increased. Since very little can be done to increase the stump sprout contribution in the short run, the focus of activities must be to increase the contribution of advance reproduction. A technique currently recommended reduces stand basal area of the mature stand from below with herbicides, leaving a main canopy with no canopy gaps. This treatment allows existing, small oak seedlings to grow, but does not encourage vellow-poplar establishment and growth. The herbicide treatment prevents tolerant, mid-story stems from sprouting and removes them as a source of competition both before and after overwood removal. About 10 years after treatment, oak seedlings should be large enough to compete, and overwood removal can begin.

INTRODUCTION

Northern red oak is a very important tree species in the Southern Appalachians. The stands in which it occurs range from those dominated by red oak and other upland oaks to those dominated by deciduous species other than oaks. High-quality wood and relatively rapid diameter growth make red oak one of the most desirable sawtimber species, and its good, but infrequent, acorn crops are an important wildlife food. The problem of regenerating northern red oak on high-quality sites in the Southern Appalachians is well-documented and has been the subject of much research over the past several decades. In this paper I will discuss the overall research and the development of a successful regeneration technique.

THE PROBLEM

.(b00e). we subsequently found important for red oak in the Southern Appalachians (Loffis advance reproduction prior to harvest (Sander 1971, Sander 1972), a relationship of advance reproduction following harvest was positively correlated with size of oak regeneration (Sander and Clark 1971). And Sander was reporting that growth of pre-existing vegetative structures—advance reproduction and stump-spouts—for in the Central States by Ivan Sander and Bryan Clark emphasized the importance essentially the same as would be expected after clearcutting. Concurrent research, regeneration under shelterwoods, over a broad range of residual basal areas, was shade-tolerant subcanopy species, on high-quality sites after clearcutting; and (3) competition with other species, particularly yellow-poplar and sprouts of occurs); (2) oaks, and particularly northern red oak, performed poorly in after clearcutting on lower quality, xeric sites (where northern red oak seldom 1983a). The findings from these studies indicated that(1) oaks regenerated well 1970, McGee 1975, Beck and Hooper 1986, Della-Bianca and Beck 1983, Loftis (2) would provide important direction for future research (McGee 1967, Beck studies on the Bent Creek Experimental Forest that (1) identified the problem and Thirty years ago, Charles E. McGee and Don Beck began a series of regeneration

McGee (1967) observed that under mature mixed oak stands, "seed germinate, the seedlings live for a few years, die, and are replaced by new ones. Cutting the overstory interrupts this cycle and stimulates growth." A study installed by Beck (1970) provides a quantitative description of the dynamics of red oak seedling become established whenever there is a good acorn crop, but after 10 years fewer than 10 percent of the seedlings have survived. And those survivors have grown very little (Loftis 1983b).




Figure 1-Survival curve for a cohort of red oak seedlings growing under undisturbed conditions under full canopy.




Figure 2—Mean height attained over time by a cohort of red oak seedlings growing under undisturbed conditions under full canopy.

It is now clear that regenerating red oak on high-quality sites is a problem in large part because the large advance reproduction necessary for maintaining a red oak component in the next stand is seldom present at the time of heavy regeneration cuts. Further, large red oak advance reproduction will not develop in the absence of disturbance sometime prior to harvest.

SOLUTION

Assessing Oak Regeneration Potential In a stand where regeneration is being considered, and where maintaining a component of red oak in the regenerated stand is a management objective, the first step in meeting this objective is an assessment of existing red oak regeneration potential. As Johnson (this Proceedings) has indicated, regeneration potential is the expected contribution of existing advance reproduction and stump sprouts to the next stand.

Based on the relationship between preharvest size of advance reproduction and its development after harvest, we have developed a model that predicts the probability that an advance red oak stem will become dominant or codominant in the next stand as a function of its size and the site quality on which it is growing. These probability values and comparable values for stump sprouts adapted from Sander and others (1984) are presented in table 1.

Basal diameter of		Site index		
advance reproduction	70	80	90	
(In.)		4		
0.1	0.01	0.00	0.00	
0.2	0.02	0.01	0.00	
0.3	0.03	0.01	0.01	
0.4	0.05	0.02	0.01	
0.5	0.07	0.03	0.02	
0.6	0.10	0.05	0.02	
0.7	0.13	0.07	0.03	
0.8	0.17	0.09	0.04	
0.9	0.20	0.12	0.06	
1.0	0.24	0.14	0.07	
1.1-1.5	0.34	0.24	0.14	
1.6-2.0	0.44	0.38	0.28	
d.b.h.	Stump sprouts ¹			
2-5	0.49	0.42	0.35	
6-11	0.46	0.39	0.32	
12-16	0.38	0.31	0.24	
17+	0.24	0.17	0.10	

Table 1—Dominance probabilities, at age 20, for northern red oak advance reproduction and stump sprouts.

¹ Adapted from Sander and others (1984).

where

These probability values can be used to estimate the contribution of existing advance reproduction and sprouts produced by stumps if the stand were regenerated immediately by:

$$\mathbf{N} = \Sigma \, n_{ij} \, p_{ij}$$

N = the expected number of dominant and codominant red oaks per acre at age 20 in the new stand

 n_{ij} = the number of red oak per acre in the *i*th size class on the *j*th site index class

 p_{ij} = the probability of red oak in the *i*th size class on the *j*th site index class becoming dominant or codominant at age 20.

For example, consider a mature stand on a site index of 90 with the following distribution of red oak:

Basal diameter	n(i) (stems/acre)	p(i) (from table 1)	(expec	N ted number)
0.1	700	0.00		0
0.2	200	0.00		0
0.3	50	0.01		0.5
d.b.h.				
2-5	0	0.35		0
6-11	2	0.32		0.6
12-16	2	0.24		0.5
17+	5	0.10		<u>0.5</u>
			Total	2.1

Obviously, the stand under examination must be sampled to provide estimates of n(i) for advance reproduction (stems 2 in. or less in diameter at groundline) and for stems 2 in. d.b.h. and larger. Site index (j) must also be determined. Given the age of the study on which the probability model is based, the results are tentative. However, the models should allow the silviculturist to recognize, at the very least, those stands where the current red oak regeneration potential is inadequate to meet management goals.

If the current red oak regeneration potential is deemed adequate, the stand can be harvested in a single cut. However, if management objectives indicate the need to retain some overstory trees for a period of time, oak regeneration can still be successful if more than one cut is used to harvest the stand.

Increasing Red Oak Regeneration Potential If current red oak regeneration potential is deemed inadequate to meet management objectives, silvicultural treatment must be used to increase the regeneration potential.

Shelterwood methods have long been recommended for regenerating oaks. However, as stated earlier, our experience with shelterwood cuts was disappointing. In shelterwoods with residual basal areas ranging from 25 sq. ft. to 66 sq. ft., oak seedlings did grow. But oak seedlings were soon overtopped by yellow-poplar seedlings and by sprouts of tolerant subcanopy species. Oaks did not benefit, relative to other species, from higher residual basal areas or longer periods of overstory retention. These results suggest that a shelterwood method to encourage red oak regeneration must (1) provide for the development of large advance red oak reproduction while preventing the establishment and growth of yellow-poplar, and (2) control competition from shade-tolerant, midstory species.

To address this question, we installed plots in well-stocked, mature stands with basal area reductions ranging from 0 to 40 percent of initial basal area. The basal area reduction was accomplished from below, with herbicides and deals in the most straight-forward manner with sprouting of understory species. A complete description of the methodology used in these studies and a presentation of early results can be found in Loftis (1990b).

The results of these studies after 13 years are very promising. We have defined residual stand densities that impede the establishment and growth of yellow-poplar while still providing enough light for the growth of small red oak seedlings that exist at the time of treatment. We did not find in these studies that shelterwoods enhance the establishment of new red oak seedlings.

If advance reproduction exists in the stand, but is not adequate to provide the desired oak component in the next stand (based on the model noted above), basal area reductions in the ranges of 25-30 percent on SI=90, 30-35 percent on SI=80, and 35-40 percent on SI=70 sites will result in development of larger advance reproduction. This treatment, should leave a stand with *no gaps* in the main canopy. Rarely would this initial treatment result in the removal of commercial material. The cost of application of this method is likely to be only marginally greater than the cost incurred when the same material is removed in other methods

used to create well-stocked young hardwood stands. The initial treatment controls essentially the same vegetation that must be controled regardless of the choice of regeneration method. The difference is primarily one of timing.

Mean basal diameter growth of red oak seedlings resulting from the above treatment can be predicted (figure 3), and a future basal diameter distribution can be projected. For example, in a mature stand on a site index of 90 in which basal area is reduced by 27.5 percent, the predicted mean basal diameter growth would be 0.42 in. (figure 4). The data suggest that growth about this mean is normally distributed.




Figure 3—Mean basal diameter growth of red oak seedlings 10 years after treatment, as a function of site index and basal area reduction.




Figure 4—Predicted mean basal diameter growth of red oak seedlings 10 years after reducing basal area by 27.5 percent on site index 90.

By recovering the standard deviation from the equation:

$$SD = 0.08930 + 0.4314$$
 (Growth)

a vector of transition probabilities based on the normal distribution can be generated. These transition probabilities are the probabilities that a seedling will grow 0.0 in., 0.1 in., 0.2 in., etc., in 10 years.

10-year growth transition	probabilities (SI=90)
Growth	Probability
0	0.086
0.1	0.073
0.2	0.106
0.3	0.133
0.4	0.146
0.5	0.141
0.6	0.117
0.7	0.087
0.8	0.055
0.9	0.031
1.0	0.015
1.1	0.007
1.2	0.002
1.3	0.001

For a given population of seedlings, say:

Basal diameter		<u>Number/acre</u>
0.1	-	750
0.2	-	200
0.3	-	50

adjusted for survival:

<u>10-</u>	<u>ear survival</u>
750	525
200	x 0.7
50	35

a future basal diameter distribution of red oak advance reproduction can be projected:

	Trans	ition pro	<u>bability</u>				Future al diame <u>stributic</u>	 Basal <u>diameter</u>
10	0.086	0	0	1			45.2	.1
	0.073	0.086	0				50.4	.2
	0.106	0.073	0.086				68.9	.3
	0.133	0.106	0.073				87.2	.4
).146	0.133	0.106		525		99.0	.5
	D.141	0.146	0.133				99.1	.6
	D.117	0.141	0.146	x	140	=	86.3	.7
).087	0.117	0.141				67.0	.8
).055	0.087	0.117		35	1	45.2	.9
).031	0.055	0.087				27.0	1.0
).015	0.031	0.055				14.1	1.1
).007	0.015	0.031				6.9	1.2
	0.002	0.007	0.015				2.6	1.3
10	0.001	0.002	0.007				1.1	1.4

	<u>Nu</u>	<u>mber/acre</u>	x	Dominance probability		pected number of dominant & codominant trees <u>20 years after harvest</u>
.1	-	45.2		0.00		0
.2	-	50.4		0.00		0
.3	-	68.9		0.01		0.7
.4	-	87.2		0.01		0.9
.5	-	99.0		0.02		2.0
.6	-	99.1		0.02		2.0
.7	-	86.3		0.03		2.6
.8	-	67.0		0.04		2.7
.9	-	45.2		0.06		2.7
1.0	-	27.0		0.07		1.9
1.1-1.5	-	24.7		0.14		<u>3.5</u>
					1	Γ otal = 19

Then, by applying the dominance probabilites to this projected distribution we can predict how much oak we would expect if this treatment is applied:

By collecting information about the size distribution of red oak—from small seedlings to large trees—in an existing mature stand, the forester can explore the consequences of two potential prescriptions. First, the dominance probabilities (table 1) can be applied to the existing size distribution to predict the red oak component expected if the overstory removal in one or more cuts is begun immediately, with appropriate treatment of competing vegetation. Second, the existing size distribution can be used in the projection method presented above to predict the expected red oak component in the next stand if overstory removal is preceded by a basal area reduction from below followed by 10 years of advance reproduction development, as described above.

Limits of Application The results are based on studies primarily of northern red oak in the Southern Appalachians. The stands in which this study were conducted would be classified as Northern red oak-White oak-Yellow-poplar, by far the most common mesic association in the Southern Appalachains. In this association, yellow-poplar is a very aggressive competitor. The association lacks a truly shade-tolerant canopy species, but does contain a number of shade-tolerant subcanopy species. Further, these studies were conducted in stands that were initially well-stocked, with complete closure of the main canopy and significant density of subordinate crown layers. While similar results have been reported in bottomland hardwoods (Janzen and Hodges 1985) and in more xeric oak systems (Graney 1989, Schlesinger and others 1993), the details of the method and almost certainly the models will be different.

Based on observations and measurements in some of the study plots (Loftis, unpublished data), black oak, scarlet oak, and chestnut oak respond to release similarly to red oak. White oak, however, does not.

CONCLUSION If an objective is to maintain a red oak component on a site where we are considering regeneration, we have (1) a model to predict the future red oak component if the current stand were regenerated immediately by clearcutting or a heavy shelterwood cut; (2) a method to use when the predicted oak component falls short of that desired; and (3) models to predict the outcome of this alternative prior to its application. The use of the models does require that data be collected to enter into the models, however. By doing so the manager should have the information necessary to make a prudent choice of regeneration methods.

LITERATURE CITED Beck, D.E. 1970. Effect of competition on survival and height growth of red oak seedlings. Res. Pap. SE-56. U.S. Department of Agriculture, Forest Service. 7 pp.

- Beck, D.E.; Hooper, R.M. 1986. Development of a Southern Appalachian hardwood stand after clearcutting. Southern Journal of Applied Forestry. 10(3):168-172.
- Della-Bianca, Lino; Beck, D.E. 1983. Selection Management in Southern Appalachian Hardwoods. Southern Journal of Applied Forestry. 9(3):191-196.
- Loftis, D.L. 1983a. Regenerating Southern Appalachian mixed hardwoods with the shelterwood method. Southern Journal of Applied Forestry. 7(4):212-217.
- Loftis, D.L. 1983b. Regenerating red oak on productive sites in the Southern Appalachians: A research approach. In: Jones, E.P., Jr. (ed.). Proc. 2nd Biennial Southern Silvicultural Research Conference. Gen. Tech. Rep. SE-24: 144-150.
- Loftis, D.L. 1990a. Predicting Post-Harvest Performance of Advance Red Oak Reproduction in the Southern Appalachians. Forest Science. 36(4):908-916.
- Loftis, D.L. 1990b. A Shelterwood Method for Regenerating Red Oak in the Southern Appalachians. Forest Science. 36(4):917-929.
- McGee, C.E. 1967. Regeneration in Southern Appalachian Oak Stands. U.S. Forest Service Res. Note SE-72. 6 pp.
- McGee, C.E. 1975. Regeneration alternatives in mixed oak stands. Res. Pap. SE-125. U.S. Department of Agriculture, Forest Service. 8 pp.
- Sander, I. L. 1971. Height growth of new oak sprouts depends on size of advance reproduction. Journal of Forestry. 69:809-811.
- Sander, I.L. 1972. Size of advance reproduction: Key to growth following harvest cutting. Res. Pap. NC-79. U.S. Department of Agriculture, Forest Service. 6 pp.
- Sander, I.L.; Clark, F.B. 1971. Reproduction of upland hardwood forests in the central states. Agric. Handb. 405. U.S. Department of Argiculture, Forest Service. 25 pp.

Schlesinger, [and others]. Northern Journal of Applied Forestry. [In Press].

Regenerating Oaks in the Central Appalachians

H. Clay Smith, Northeastern Forest Experiment Station, USDA Forest Service, Parsons, WV 26287

ABSTRACT In the Central Appalachians, regeneration of oaks on fair-to-poor sites is usually successful using even-aged silvicultural regeneration methods. However, for the good-to-better growing sites (red oak site index 70+), the establishment and development of natural oak regeneration is of major concern. Demonstrations of consistently successful methods for the establishment and subsequent development of oak reproduction, or both, on the better growing sites are limited. A promising shelterwood technique to stimulate oak seedling height growth has not been utilized to any extent. Keys to successfully establishing oak on the good growing sites include seedling establishment, stimulating seedling height growth, and controlling vegetation competition.

INTRODUCTION The Central Appalachian forest is an estimated 23,000,000 acres that includes portions of southern Pennsylvania, western Maryland, West Virginia, western Virginia, eastern Kentucky, and southeastern Ohio. The Appalachians are mountainous with steep topography and a wide range of stand and site characteristics. A variety of slopes, moisture regimes, and soils are common. Access is difficult; however, the Appalachians are well known for the production of quality hardwood timber products. Often 25 commercial species or more occur in a given inventory. Oaks seldom exist in pure stands but more in mixtures with other hardwoods and conifers.

Historical events have played a major role in the establishment of the presently mature oak stands in the Central Appalachians. Repeated fires, death of the American chestnut, probable changes in wildlife and insect populations, and perhaps climatic conditions related to flowering are some of the factors that may have influenced availability of acorns and oak seedling development. Crow (1988) suggested a possible cyclic pattern related to the establishment of conifers within mature oak stands and, conversely, oak reproduction occurring within mature conifer stands. In the eastern forest, Carvell and Tryon (1961) concluded that disturbances reduce canopy densities and create openings, thus improving chances for the development of advance reproduction. Their hypothesis was influenced by a good correlation between amount of oak regeneration and amount of light reaching the forest floor.

Though oaks can be readily regenerated on fair-to-poor oak sites (northern red oak site index 50, 60, or below), oak regeneration problems occur on cove hardwood sites or stands with a site index of 70 and above. These good-to-excellent sites often contain a good source of available oak seed in the overstory. Because regenerating oak on the fair-to-poor sites is not a major problem in the Central Appalachians, information in this paper will focus on the better hardwood growing sites.

Establishment of oak simply cannot occur without available reproduction from stump sprouts, advanced reproduction, planted seedlings, or acorns. On these better oak sites, there are at least three oak reproduction understory situations: (1) oak seedlings or saplings not present; (2) some oaks present but inadequate (usually not tall enough); and (3) oaks present and tall enough (4.5 ft.) to respond after removal of the overstory. However, in reality, the situation where numerous oak understory stems are greater than 4.5 ft. rarely occurs. Of course, when oak stems are not present in the understory, there are two obvious choices—wait for a bumper seed crop to occur or plant seedlings or perhaps acorns. Assuming overstory trees are mature enough to produce acorns, a bumper seed crop means having enough acorns to feed all the wildlife, insects, etc., and still have acorns available for seedling establishment. However, these bumper seed crops are rare and not predictable. Regardless, the presence of oak regeneration does not assure success.

Another factor to establishing oak is to ensure that existing oak seedlings and saplings produce an increased and continuous height growth response. Getting oak seedlings to increase in height is costly and not an easy task. Too often, the release methods used stimulate competing vegetation at the expense of the oak stems or the oak seedlings respond poorly to the release technique. Released oak seedlings are exposed to adverse weather and often damaged by spring frosts and freezes (McGee 1973).

Another key to establishing oak is the need to keep the oak stems competitive as the young stand develops so that some oaks will become a component of the mature stand.

SUCCESSFUL REGENERATION METHODS AND SILVICULTURE SYSTEMS

On fair-to-poor growing sites, oaks can be consistently regenerated by even-aged silvicultural methods such as clearcutting or shelterwood (Roach and Gingrich 1968, Sander and Clark 1971). On the better oak sites, regenerating oak becomes more difficult and inconsistent. There has been much theoretical discussion about using commercial shelterwoods to reproduce oak seedlings on the better growing sites (Sander and others 1983). However, successful on-the-ground demonstrations are limited. In most cases, seed crops were inadequate or the initial cut of the shelterwood stimulated competing stems, such as yellow-poplar, that quickly overtopped established oak seedlings. Wolf (1991) reported oak regeneration success (red and white oak families) using the shelterwood system on Glatfelter Pulp Wood Company land in south-central Pennsylvania. Failures were related to not controlling understory vegetation, the white-tailed deer, or both. Oak site index for the areas ranged from 55 to 75 (fair-to-good growing sites). Site index 75 is near the lower range for oak regeneration problems in the Central Appalachians.

Success using Wolf's shelterwood approach included:

• Start with mature, fully stocked stands with a number of seed-producing oak trees.

- Use stands having terrain well suited for maneuvering harvesting equipment.
- Complete initial seed cut within 3 years after a bumper seed crop.
- Leave 55 to 60 percent oak crown cover in the residual overstory.
- Select residual trees based on distance between crowns.
- Control undesirable shade-tolerant understory competing vegetation.

• Complete overstory removal cut when 70 percent of milacre plots are stocked with at least one red oak seedling 2 ft. tall. This seedling height usually occurs within 5 to 10 years after the initial cut.

Loftis (1990b) used a "noncommercial" shelterwood method to stimulate growth of existing advanced red oak reproduction in the Southern Appalachians. Success was found by reducing basal areas in mature, fully stocked stands to 60, 65, and 70 percent of the initial stand basal area for red oak site index areas of 70, 80, and 90, respectively. Trees were removed from beneath the main crown and herbicides applied to the cut stumps or stems were basal sprayed. Little to no cutting was done in the main canopy. These treatment methods prevented yellow-poplar, a primary competitor of red oak, from becoming established before the final removal cut (Loftis 1990b, O'Hara 1986). Also, the herbicide treatment eliminated most sprout competition from shade-tolerant understory species after the final removal cut. However, application of this technique is minimal in the Central Appalachians due, perhaps, to the establishment of other desirable (non-oak) reproduction, transfer of technical knowledge, gypsy moth, few to no oak seedlings present, cost, or just lack of confidence that the technique works or will work for the given situation.

While consulting with others in the Central Appalachians, a few success stories were reported related to regenerating red oak on good-to-excellent sites. For instance, there are some successful oak regeneration areas on Westvāco properties in south-central West Virginia. Initially, almost no advanced reproduction was present in these stands. Over the years, partial cuttings were done with the idea of eventually clearcutting the stands. Forest managers believe that these partial cuttings increased the establishment of the advanced oak reproduction. There are a few on-the-ground examples where some overstory trees were removed, and advanced oak stems are successfully competing with other regeneration for the future stands. Currently, it appears that some oaks will be codominant trees in the future stands. Precommercial cultural techniques, such as using herbicides to favor the oaks, are being tried experimentally in these stands. Most of the competing vegetation develops foliage earlier than the oak; thus, foliar-active herbicides are used to remove the competition before oaks break bud. Preliminary results are encouraging.

Where no oak seedlings are present in the understory, some managers suggest eliminating the heavy shade understory to allow patches of sunlight to reach the forest floor. This treatment may allow more opportunity for acorn germination and seedling development. Though this application might work for some situations, it is very uncertain. Where seed-bearing oaks are present and when a bumper acorn seed crop occurs, oak seedlings will become established everywhere throughout the understory on all sites. These seedlings will usually remain for a couple of years and then rapidly begin to die. Thus, managers have a few years are used available techniques to stimulate understory oak height growth. Unfortunately, bumper red oak acorn crops do not occur often. During the last 30 years, there has been a maximum of 3 to 4 bumper crop years in the Fernow Experimental Forest near Parsons West Virginia—when walking in the woods is like roller skating on acorns during the winter and spring after the fall seeding.

Some forest managers indicated that getting established advanced oak regeneration to respond might be done by felling the understory trees, such as sugar maple, striped maple, and possibly hemlocks. Hardwood sprouts could be a problem and, if so, herbicides could be used as necessary. This is similar to the Loftis (1990b) approach.

On the Fernow Experimental Forest, a shelterwood study was established 8 years ago for the purpose of establishing red oak seedlings and/or stimulating seedling development. Initially, 16 two-acre plots were established on good growing sites (red oak site index 70) and the following year 12 additional plots were established on excellent growing sites (site index 80). Various overstory residual stocking percent treatments were used including 45, 60, 75, and 100 percent (no cutting). Also, understory treatments were done using herbicides for seedlings, saplings, and pole-size trees up to 10 in. d.b.h. (excluding oaks). Replicated plot treatments included no understory or overstory treatments; understory but no overstory treatments; understory and overstory treatments combined. Each plot was underplanted with 50 red oak seedlings during the spring immediately after the dormant season cutting treatment. Red oaks were the main residual trees after the initial shelterwood cuts. The fall before the initial treatment period appeared to be a reasonable acorn seed year. Before the initial shelterwood treatment, two of the plots on the site index 70 areas had hundreds of advanced red oak seedlings 2 ft. tall and taller. At the time, it was thought a clearcut would have been as successful as a shelterwood due to the height of the advanced oak stems. A few other plots on this site index 70 area had some small advanced red oak seedlings present before treatment.

The two plots with oak stems 2 ft. tall and taller were treated with a 45 percent residual shelterwood overstory removal. Also, an understory basal spray herbicide treatment was used. Oak reproduction dominated these areas and after 5 years, a major blowdown occurred in this portion of the study area. Eight years after treatment, the remaining overstory trees were removed. Red oak will dominate the future stand. However, for the remaining shelterwood and/or herbicide treatment plots, oak regeneration is absent; if present, it is less than 2 ft. tall and often shaded by a dense layer of vegetation. Recently, on some plots, the reproduction layer of competing vegetation was treated with herbicides in an effort to maintain and stimulate the oak seedlings to respond. Unfortunately, for the Fernow study, the timing of the initial understory-overstory treatments did not coincide with a bumper acorn seed crop. We should have waited until natural seedlings were well established before applying the study treatments; however, we thought the planted seedlings would respond. Presently, after 8 years, surviving planted oak seedlings average less than 2 ft. tall, but most died due to deer browsing, weather, and insects.

Thus, there has been some success in the Central Appalachian region in regenerating oak, primarily red oak, on the good and better growing sites. However, in general, specific methods to establish oak on these sites tend to be Page 277 of 404

more happenstance or theoretical than actual. Establishing oak seedlings is the most important initial concern. Once natural oak seedlings (resulting from a bumper acorn seed crop) are established, techniques to stimulate height growth of the seedlings become most important. Methods of stimulating the height growth of existing understory oak seedlings involve controlling (removing-killing) other competing understory seedlings, saplings, and pole stems. Though expensive, the technique has potential. The purpose is to provide enough light for oak seedlings to respond yet minimize the establishment of yellow-poplar. Kittredge (personal comments) indicates that stimulation of oak regeneration height growth is important, but he believes the oak seedlings have to spend some portion of time initially growing slowly in the understory while developing good roots. He also indicates there may have been some oak regeneration mistakes made in the past where foresters had excellent seedlings established from bumper acorn crops but the overstory canopy was opened too much, too soon. Kittredge indicated the new seedlings were not able to take advantage of the light due to poor root development.

A history of repetitive fires to regenerate oaks has been related to the presence of oaks in the current overstory stands on good-to-excellent Appalachian hardwood growing sites. However, fire specifically to regenerate oaks may not be practical in the Central Appalachians.

ANTICIPATED PROBLEMS

Anticipated problems to the establishment of oaks on the good-to-better growing sites are related to the use of herbicides and possibly the use of fire. Herbicides and fire are means to control the competing vegetation. Continued political constraints and social concerns related to using herbicides may result in limited to perhaps no use of herbicides on public forest lands. Fire, if successful, would be a difficult tool to apply from a practical basis. Many of the good and better growing sites would be difficult to burn under prevailing weather conditions; more importantly, when these moist sites would burn, the fire hazard in the general area would be too high for safe use of controlled burning. Residual stand damage as a result of the hot fire would be major (Wendel and Smith 1986). Too, air pollution resulting from prescribed fire is of great public concern.

First Key Factor Problems in establishment and development of oak regeneration relate to seed production, predation, competing vegetation, and stimulating seedling height growth. Seed production for oaks, particularly red and white oak, is very sporadic and unpredictable. To be successful in establishing oak seedlings, good-to-bumper acorn crops are a necessity. When major acorn crops are produced, there are enough acorns to feed the wildlife, etc., and still have many acorns to germinate new seedlings. However, planning forest management practices around production of acorns has too much uncertainty in the Central Appalachians to be effective.

If techniques to establish oaks on these good-to-better oak growing sites are perfected, there are still two overriding factors that will have a major impact on the oak regeneration programs: deer and the gypsy moth. Deer populations in certain areas of the Central Appalachians are so high, that understory browse lines are quite visible. As the deer population increases, there will be public, and thus political, resistance to increasing deer control measures for the sake of forestry Page 278 of 404

purposes. In the future, much of the Appalachian forests could have a browse line similar to parts of the northeastern forests. When this occurs, it is too late for quick remedial actions.

Gypsy moths are another concern. This insect presently has no effective natural controls in the United States, though there is a fungus from Japan that has possibilities (Andreadis and Weseloh 1990). Thus, once present, the gypsy moth remains in the forest indefinitely. Oaks are the preferred food source for the gypsy moth and tree mortality is usually high following one or two periodic defoliations. As the moths steadily progress from New England into the Southern Appalachians, and if the Asian gypsy moth begins to progress from the west to the east coast, it is reasonable to ask why do we even consider establishing and growing oaks in the future stands? The answer to this pertinent question is not obvious. Forest managers will continue to assess the situation and will provide the answer by their actions. For those wanting to manage oaks, the establishment of advanced oak reproduction and the tending and culturing of these stems will be an acceptable management objective. Methods for controlling gypsy moths will become part of the management scheme. Also, efforts could be made to stimulate oaks to retain vigorous crowns in mixed hardwood stands that might withstand periodic defoliations (Gottschalk 1988). However, these recommendations/concepts are in the research stage.

In reality, the presence of gypsy moths will have a long-term effect on the silvicultural practices used in eastern hardwood forest management. However, some natural resistance to defoliation will occur and, though the oak composition in the Central Appalachian forest stands will be significantly reduced, it is not expected to be completely eliminated by the moths. Many oak trees still remain associated with the gypsy moths. The forest manager's inability to reproduce oak on these good growing sites may have a negative impact on moth populations. The stands will become more mixed with other hardwoods, thus minimizing the attractiveness to the moths. Stands on forests with fewer oak should not be able to sustain the moths at frequent, epidemic levels (Steve Jones, personal comments).

Second Key Factor Once acorns germinate and seedlings become established, competing vegetation often can become a major deterrent to their development. Oak seedlings commonly have very slow height growth during their early years of development. Treatments to stimulate oak height growth often trigger an overwhelming response by competing vegetation or can expose the seedlings to frost or freezing problems. There is a fine line between just enough light only to stimulate oak seedling growth and too much light, thus stimulating shade-intolerant reproduction. Apparently shaded oaks will start growth in the spring before the canopy fully leafs out (McGee 1974). A noncommercial shelterwood technique described by Loftis (1990b) is the proposed approach for use in the Southern Appalachians.

COSTS, BENEFITS, AND RELIABILITY

Establishing natural oak regeneration on the good-to-better growing sites can be expensive and results are uncertain. There are a number of practices to consider. In the Central Appalachians, I suggest waiting until the natural oak seedlings are established before applying any cultural practices. This could prevent future problems, especially if the anticipated seed crop is a bust. Felling saplings and Page 279 of 404

poles in the noncommercial shelterwood technique can cost from \$30 to \$40 per acre, while the cost to treat poles and saplings with herbicides approaches \$60 per acre or more. However, effective herbicide treatments eliminate stump sprouts. Remember, do not consider this non commercial shelterwood approach until natural seedling establishment has occurred.

If natural seedling establishment is rare, consider planting bare-root seedlings following clearcutting. All saplings and poles should be cut during or following removal of the commercial stems. Cutting during the mid to late summer and early fall is preferable in that sprouting would be minimized. If saplings and poles are felled during logging and left on the ground, cost or credit to the logger would average about \$30 to \$40 per acre. If this site preparation occurs after logging, the contract cost would increase to at least \$60 per acre due to the presence of slash. However, depending on the local market situation, the minimum size for merchantable trees could range from 6 to 12 in. d.b.h. Some or all of the pole size trees might be sold, thus reducing the cost.

Also, use plastic tree shelters on the planted and even any natural seedlings that might be present. These shelters protect the seedlings from deer and other animal damage while providing a greenhouse effect that stimulates tree growth. Currently 2-0 special root-pruned seedlings cost about \$150 per 1,000 (15¢ per seedling). These "special" seedlings are usually root pruned in early August during the first year in the nursery seedbed and only 2-0 seedlings with at least 0.4 to 0.5 in. root collar diameter are planted. On a good planting site, one person could plant about 200 seedlings per day. Thus, if the hourly wage were \$7, it would cost \$56 to plant 200 seedlings per 8-hour day, plus another \$30 for the seedlings, for a total cost of \$86 (43¢ per seedling).

A 5-ft. tree shelter would need to be installed around each planted seedling. If the deer population were low, perhaps shorter shelters could be used. Current costs of 5-ft. shelters vary from about 2 to 6 depending on brand and quantity purchased. Also, in most instances, 1- x 1-in. sharpened white oak stakes are used to hold the shelter in place. These stakes cost approximately 50¢ to 60¢ each. Two people can install about 200 shelters per day at a labor cost of 14 per hour using the 7 per hour rate. Using a 4.50 cost for each shelter-stake, planting 30 seedlings per acre, and installing a shelter on each seedling would cost 164.70 per acre (table 1). With the planted seedling-shelter on a forested site, there is a strong possibility that one or more herbicide applications will be necessary to control competing vegetation on the good-to-excellent Appalachian hardwood sites. This could involve applying herbicides in a 10-ft., or larger, radius around each shelter.

Planting acorns using shelters has potential. If acorns are used, planting cost will be reduced; however, the cost of purchasing and installing shelters would be similar.

The artificial regeneration technique has some promise, and on the Fernow Experimental Forest, approximately 40 percent of planted (2-0) and a few 1-yearold natural seedlings have grown out the top of 5-ft. shelters in clearcut areas during the first year. However, there is uncertainty as to whether this seedling height growth will continue sufficiently to keep above, or at least compete with, the natural reproduction on these good-to-excellent cove hardwood sites.

Materials, Labor	Cost
2-0 special root-pruned seedling	\$150 per 1,000
Labor rate	\$7 per hour
Shelters, 5-ft.	\$2-\$6 per shelter
Stakes, 5-ft. 1- x 1-in. white oak	50¢-60¢ per stake
Example:	
Plant 30 seedlings per acre (15¢ per seedling)	\$ 4.50
Labor cost (28¢ per seedling)	8.40
Install 30, 5-ft. shelter-stakes per acre	
(\$4.50 per shelter-stake)	135.00
Labor cost (56¢ per shelter-stake)	_16.80
Total cost per acre	\$164.70

 Table 1—Estimated cost of planting red oak seedlings and plastic shelters

BEST RECOMMENDATIONS TO ACHIEVE VARIOUS OAK REGENERATION GOALS

On drier, poor-to-fair sites, where mature oak trees occur, oaks can be reproduced through clearcutting or any other form of even-aged silviculture. However, for the moist, good-to-better growing sites, it has been difficult to reproduce oak from mature oak stands. With change in site quality, there is a change in species composition and the species differ in their ability to sprout. Sprouting is a function of size, age, growth rate, etc. On poor-to-fair sites, chestnut and scarlet oaks may dominate. On fair-to-medium sites, scarlet, chestnut, black, and white oaks may be dominant. On good sites, northern red, white, and black oaks will occur, and on the best sites, northern red oak is usually the only one that will compete (David W. Smith, personal communication).

Ideally, when wanting to reproduce oak stands, if 435 well-established, advanced oak stems per acre, over 4.5 ft. tall are present (including potential stump sprouts) and free from shade of a mid-canopy, then clearcutting will be successful (Sander and Clark 1971, Sander and others 1983). However, it is rare to find advanced oak reproduction more than a foot tall on the good-to-excellent Appalachian hardwood sites. Where these small oak stems occur, a potential practice is to reduce the overstory canopy by applying herbicides to all sapling and pole trees (1 to 10 in. d.b.h.) to give these small oak seedlings room to grow (Loftis 1990b). Usually no trees are cut in the upper crown canopy. The objective is to provide just enough light to let the oak grow without allowing other competing vegetation such as yellow-poplar to become established. Northern red oak advanced seedling growth response is based on root collar diameter and the probability of these stems being in the dominant or codominant crown class 20 years after clearcutting (Loftis 1990a). Using this probability concept, silvicultural decisions, such as when to complete the final cut, are based on the desired number of dominant and codominant red oak stems the landowner wants in 20 years. This is where the uncertainty occurs. Once the overstory is removed, will the oak stems compete with the associated vegetation? In the Central Appalachians, there are few stands on these good-to-better growing sites that have oak reproduction of proper size to consider removing the overstory.

Another technique suggested for use in openings is to clip the tops of established natural seedlings near groundline, then to protect these clipped seedlings by installing plastic shelters (Kittredge and Kelty 1992). On the Fernow, tree shelters were installed on a few natural seedlings after clearcutting and the seedlings responded.

For most of the Appalachians, a realistic objective for reproducing of oaks on the good-to-excellent growing sites is to reproduce *some* oak in the next stand. Thus, guidelines will be less stringent than those recommended for reproducing a fully stocked oak stand.

In general, bumper seed crops are needed to provide enough acorns to overwhelm the rodents, mammals, insects, and other users to allow for the establishment of the young seedlings. Where no oak seedlings exist in the understory, choices are limited if an oak component is an objective in the next stand. Managers will have to wait for a bumper seed year. Planning forest management programs based on climatic or natural occurrences is uncertain. There is another and yet unproven technique that may work. Plant "special" 2-year-old northern red oak seedlings that are root-pruned during early August the first year in the nursery seedbed. Taproots are pruned 6 to 8 in. below the ground, resulting in the development of a more fibrous root system (Johnson and others 1986). Since the special seedlings have a large root system, it is very important that good planting techniques be used. A major problem in planting these large seedlings (root collar diameter at least 0.4 in.) is providing a planting hole of sufficient size to accommodate the root mass. If a hole is too small, the roots are easily cramped and become "j-shaped," and seedlings die or have little growth potential.

Planting perhaps 30 to 40 "special" red oak seedlings per acre in a clearcut area before the start of the first growing season after a dormant season cut is worthy of consideration. Immediately after planting, install at least a 5-ft. plastic shelter around each planted seedling. Before installing the shelters, seedlings could be top pruned about 6 to 8 in. above the ground. However, to date, the best results we have on the Fernow Experimental Forest are where the planting stock was top pruned at 1.5 ft. This was done in the seedbed immediately before the 2-0 seedlings were lifted for distribution. Our ideal objective is to have each seedling growing out the top of a 5-ft. shelter in 1 year. Though specific results are not available, recently stratified red oak acorns were planted during two different spring seasons on the Fernow Experimental Forest. Shelters were installed immediately after each planting. Rodents did not damage or remove the acorns and the acorns developed into seedlings.

CONCLUSIONS In general, I believe successful regeneration of red oak would be to have 15 to 20 codominant oak stems per acre on good-to-excellent Central Appalachian hardwood sites when the even-aged stands are 20 years old. A probability of survival for planted seedlings without shelters is to plant four seedlings to obtain one (Johnson and others 1986). With shelters, this proportion should be less; however, this is conjecture and not based on research information. Remember, the objective is not to convert these good-to-excellent central hardwood sites to oak stands, but simply to ensure that there is an oak component in the future stands—perhaps at levels similar to the present portions.

Thus, there are two general objectives concerning oak regeneration on the better sites in the Central Appalachians:

1. Establish oak seedlings where none exist;

2. Stimulate oak seedlings to respond in height growth following establishment, in order to compete with associated vegetation and become part of the future stand overstory.

If natural oaks are present in the understory, a possible technique to stimulate these seedlings to develop is the noncommercial shelterwood technique reported by Loftis (1990b) for the Southern Appalachians. If natural oak seedlings are not present in the understory, wait for their establishment as a result of a bumper seed crop or plant oaks immediately after clearcutting. In both instances, once seedlings are established and if clearcutting is used, install a plastic tree-shelter around each seedling. Do not install brown plastic shelters on red oak seedlings where a portion of the overstory shelter remains after cutting.

The establishment of natural oak seedlings and their subsequent stem development are keyed to bumper acorn seed crops and timing of cultural practices. Cultural practices should not be applied until after the natural seedlings are established in the understory. In the past, understory-overstory cultural techniques have been applied in anticipation of bumper seed years. In most cases, these seed crops did not occur and their absence resulted in additional understory regeneration problems.

LITERATURE CITED

- Andreadis, Theodore G.; Weseloh, Ronald M. 1990. Discovery of *Entomophaga maimaiga* in North American gypsy moth, *Lymantria dispar*. In: Proceedings, national academy of science USA. 87:2461-2465.
- Carvell, K.L.; Tryon, E.H. 1961. The effect of environmental factors on the abundance of oak regeneration beneath mature oak stands. Forest Science. 7(2):98-105.
- Crow, T.R. 1988. Reproductive mode and mechanisms for self-replacement of northern red oak (Quercus rubra)--a review. Forest Science. 34(1):19-40.
- Gottschalk, Kurt W. 1988. Gypsy moth and regenerating Appalachian hardwood stands. In: Smith,
 H. Clay; Perkey, Arlyn W.; Kidd, William E., Jr., eds. Proceedings: Guidelines for regenerating Appalachian hardwood stands; 1988 May 24-26; Morgantown, WV. SAF Publication 88-03. Morgantown, WV: West Virginia University Books and U.S. Department of Agriculture, Forest Service: 241-254.
- Johnson, Paul S.; Dale, Charles D.; Davidson, Kenneth R.; Law, Jay R. 1986. Planting northern red oak in the Missouri Ozarks: A prescription. Northern Journal of Applied Forestry. 3:66-68.
- Kittredge, David B., Jr.; Kelty, Matthew J. [in press]. The use of tree shelters with northern red oak natural regeneration in southern New England. Northern Journal of Applied Forestry.
- Loftis, David L. 1990a. Predicting post-harvest performance of advanced red oak reproduction in the Southern Appalachians. Forest Science. 36(4):908-916.
- Loftis, David L. 1990b. A shelterwood method for regenerating red oak in the Southern Appalachians. Forest Science. 36(4):917-929.
- McGee, Charles E. 1973. Is variation in bud bread of red oak the result of heredity or environment? Proceedings twelfth southern forest tree improvement conference: 185-189.

- McGee, Charles E. 1974. Elevation of seed sources and planting sites affect phenology and development of red oak seedlings. Forest Science. 20(2):160-164.
- O'Hara, Kevin L. 1986. Developmental patterns of residual oaks and yellow-poplar regeneration after release in upland hardwood stands. Southern Journal of Applied Forestry. 10:244-248.
- Roach, B.A.; Gingrich, S.F. 1968. Even-aged silviculture for upland central hardwoods. Agric. Handb. 355. Washington, D.C.: U.S. Department of Agriculture, Forest Service. 39 pp.
- Sander, I.L.; Clark, F.B. 1971. Reproduction of upland hardwood forests in the Central United States. Agric. Handb. 405. Washington, DC: U.S. Department of Agriculture, Forest Service. 26 pp.
- Sander, Ivan L.; McGee, Charles E.; Day, Kenneth G.; Willard, Ralph E. 1983. Oak-Hickory. In: Silvicultural systems for the major forest types of the United States. Russell M. Burns, Technical Compiler. Agric. Handb. 445. Washington, DC: U.S. Department of Agriculture, Forest Service: 116-120.
- Wendel, G.W.; Smith, H. Clay 1986. Effects of prescribed fire in a Central Appalachian oakhickory stand. NE.RP-594. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 8 pp.
- Wolf, Wilbur E., Jr. 1991. Planned timber harvests provide for natural regeneration. Pennsylvania Forests. Winter:25-28.

Page 285 of 404

.

Artificial Regeneration of Oaks

A Historical Perspective of Planting and Seeding Oaks: Progress, Problems, and Status

Phillip E. Pope, Department of Forestry and Natural Resources, Purdue University, West Lafayette, IN 47907

ABSTRACT Historically, the planting and seeding of oaks was done for species conversion on forest lands and for reforestation of old fields. More recently artificial regeneration has been used to supplement natural regeneration prior to or following stand removal, to introduce genetically superior individuals, to reforest drastically disturbed lands, and to provide species diversity in naturally occurring or man-made wetlands.

This paper describes the progress that has been made since 1971 in solving the problems which inhibit the successful regeneration of oaks by planting and direct seeding. Reference to current literature is used to describe the state of knowledge or status of artificial regeneration.

Advances in oak regeneration have greatly improved the chances of a successful planting, but a degree of uncertainty remains. Artificial regeneration offers the best solutions to oak regeneration problems in the future. It is imperative to press forward for breakthroughs in genetics, economics, and the biology of early growth. Today, land managers should proceed with the planting and seeding of oaks but they should consider all the information provided in this symposium and proceed with caution.

INTRODUCTION The purpose of this paper is to present the current status of artificial regeneration of oaks by direct seeding and seedling planting. In the course of this presentation I have attempted to identify some of the problems that remain and provide some suggestions for future research. I have divided artificial regeneration into two major topics—direct seeding and seedling planting. Information on direct seeding is applicable for bottomlands in the southern and southeastern United States. Information on seedling planting is divided into upland and bottomland sites.

In developing a historical perspective of planting and seeding oaks I selected 1970 as the reference year. A review article by Russell (1971) described the state-of-the-art of artificial regeneration and offered suggestions on research to enhance the success of this regeneration technique. I have summarized the article by Russell and have used it as a benchmark to see how far the science has advanced since 1970.

Russell (1971) concluded "that upland oaks could be established by seeding or planting, but additional experience was needed before these methods became economical alternatives to natural regeneration." Based on his research and the literature, he concluded the following: (1) Recently forested sites were generally more favorable for artificial regeneration than abandoned fields. (2) Lack of repellents to protect acorns from animals severely limited direct seeding, but oaks could be planted readily by conventional methods and would survive well on suitable sites. (3) Ample sunlight was required for the best growth of seedlings, and competing vegetation must be controlled. (4) At best, height growth was discouragingly slow. (5) Advances in cultural methods and the development of genetically improved stock were essential to make artificial regeneration practical.

While the greatest obstacle in direct seeding was the "lack of economical ways to protect acorns from animals," the limitation to planting was the slow initial growth of the seedling. Russell offered some possibilities for speeding growth:

- 1. Improved planting stock
- 2. Better methods for competition control
- 3. Fertilization
- 4. Genetic improvement

There has been an explosion of research dealing with artificial regeneration of oak species via direct seeding and/or seedling planting since 1971. Some of the accomplishments made since 1971 follow.

DIRECT SEEDING – BOTTOMLANDS

Johnson and Krinard (1985, 1987) reported that direct seeding of acorns may be a viable artificial regeneration method. They evaluated *Querqus falcata* var. *pagodifolia*, *Q. nigra*, *Q. nuttallii*, and *Q. shumardii* for their ability to germinate, grow, and survive in an abandoned agriculture field setting in the Mississippi delta and minor stream bottoms and in the silty uplands.

Advantages to sowing acorns as opposed to planting seedlings:

1. Less expensive (eliminates the cost of growing, transporting, storing, and planting the seedling).

2. Less time consuming than seedling planting.

3. Sowing is not seasonally dependent, though the best sowing is March-April 15 to minimize rodent depredation.

While results of direct seeding of oaks have been mixed, the successes have yielded the following:

- 1. Acorns must be covered with soil.
- 2. Fall and spring seeding are equally successful.

3. Animals, particularly squirrels and chipmunks, forage for and destroy and/or consume a high percentage of the seeded acorns. No suitable repellent was available.

Animal Damage The earliest seeding tests conducted at the Forest Service Laboratory at Stoneville, Mississippi, were performed under a full forest canopy and were unsuccessful due principally to rodent damage. The researchers later discovered that seeding tests Page 288 of 404 in site prepared forest openings of 2 acres or more in size or on agricultural fields had much less rodent damage. Research results and recommendations were based on seeding these types of sites.

Species Research trials included Nuttall (Q. nuttallii Palmer), Shumard (Q. shumardii Buckl.), cherrybark (Q. falcata var. pagodifolia Ell.) and water (Q. nigra L.) oaks. Commercial plantings have included the above species as well as swamp chestnut (Q. michauxii Nutt.), chinkapin (Q. muchlenbergii Engelm.), overcup (Q. lyrata Walt.), willow (Q. phellos L.) and bur (Q. macrocarpa Michx.) oaks. Nuttall oak was the most adaptable species tested and consistently had superior percentage germination and growth over a wide range in sites from flood plains to silty upland sites. As a group, white oak (swamp chestnut, chinkapin, overcup, and bur) have lower percent seed germination in the field than the red oak group. Four seedlings per 50 ft. of linear row is considered adequate stocking.

- **Site Selection** The same guidelines used for evaluating sites prior to planting seedlings can be used for direct seeding (Baker and Broadfoot 1979). The major consideration is the timing and duration of flooding. The influence of these factors is species specific. For example, Shumard and cherrybark oak acorns should be planted on well-drained soils that flood only for short periods (a week or less) during the growing season. Neither species should be planted in clay soils subject to flooding during the growing season. Nuttall oak acorns, on the other hand, grow well in clay soils and can tolerate flooding during the dormant and early growing season. Water oak acorns are less tolerant of flooding.
- Seed Collection and
StorageA key to successful artificial regeneration is the proper collection and storage of
acorns. This important topic is covered in detail by Bonner in these Proceedings.
 - **Time of Seeding** The percentage of seeds which germinate (seedling percentage) varies with the time of year the seeds are planted, but generally acceptable germination is attainable regardless of the time of year the seeds are sown. This finding is important because sites that are inundated with water or otherwise unworkable during the dormant season can be planted during the growing season after the water recedes.

Johnson and Krinard (1985) observed that acorns may remain dormant in the soil during the dry summer months and germinate when soil moisture is favorable, i.e., late in the growing season.

Depth of Seeding Regardless of species and hence acorn size, seedling emergence is attainable with seed planted to a depth of 6 in. However, planting depth and seed size will influence percent germination and early seedling size (Johnson and Krinard 1985). The greatest seedling percentage occurred for seed planted at the 2-in. depth. Johnson and Krinard (1985) offered two reasons for sowing deeper than 2 in.: (1) to discourage pilfering by rodents and (2) to reduce the variability of the soil environment, i.e., temperature and moisture fluctuations. The first-year seedling heights were inversely related to planting depths. Site preparation, for seedbed preparation and weed control, is important to enhance seedling percentage and subsequent seedling survival and growth.

- Method of Sowing Both machine and hand planting have been successful. Hand planting is suited for cutover areas, rocky sites, or sites on steep slopes. Hand seeding has been improved by a tool developed by the USDA Forest Service Laboratory at Stoneville. Machine planting is best suited for open fields. Machines can accommodate a range in the size of acorns and will plant at various spacings. Machine seeding is an option only when soil moisture conditions are suitable.
 - **Spacing** Johnson and Krinard (1985) reported that approximately 35 percent of planted oak seed germinate, that approximately 10 percent of the planted acorns produce a "free-to-grow-tree" in 10 years on cleared forested sites, and that 25 percent of the planted acorns produce "free-to-grow-trees" on old field sites. Therefore, sowing 1,500 acorns per acre would produce 500 seedlings the first year with 150-375 "free-to-grow-trees" in 10 years. Between- and within-row spacing should allow for approximately 30 sq. ft. per acorn (Johnson and Krinard 1985).
- **Survival and Growth** Growth rates differ by species and site but, in general, survival is generally good but growth is slow for the first 5 years with only 1 to 2 ft. of height growth for the best seedlings. From age 5 to 15, trees may average 2 to 3 ft. of annual height growth. Correspondingly, diameter growth is also slow for the first 5 years and more rapid from years 5 to 15 (Johnson and Krinard 1985).
 - **Case Studies** Johnson (1983) reported on an 11-year-old study in which Nuttall oak was direct seeded in April either by machine or manually at 2-, 4- or 6-in. depths in a 20-acre, intensively site-prepared clearing (double disked) in the Mississippi delta. Field germination averaged 36 percent and was not influenced by method or depth of sowing. After 11 years, two-thirds of the Nuttall oaks (551 per acre) were free to grow and averaged 2.1 in. in diameter and 17 ft. in height.

Francis and Johnson (1985) reported that direct-seeded sawtooth oak (Q. acutissima Carsuth.) grew well on the Mississippi flood plain despite high soil pH, short-term flooding, and poor soil aeration. Sites were cleared and disked before seeding in April. Approximately 60 percent of the sown seed germinated and survived. Trees averaged 31 ft. in height and 4.6 in. in diameter after 14 years.

Johnson (1984) reported on the 10-year results of cherrybark and Shumard oak direct-seeded in the silty uplands near Vicksburg, Mississippi. The 3-acre openings were created by cutting and removing all trees > 1.0 in. d.b.h. The field germination of winter-sown acorns was 42 percent for cherrybark and 55 percent for Shumard, about twice that of spring-sown acorns. After 10 years, the average height of codominant trees averaged 23.6 ft. for Shumard and 27.0 ft. for cherrybark. The average d.b.h. was 1.5 in. for both species. The dominants and codominants in the competing natural stand averaged 40 ft. in height and 3.5 in. d.b.h. Only 5 percent of the seed spots had oak in a free-to-grow position; most codominant oaks were on higher, drier sites where competition was less severe. Johnson (1984) concluded that "direct seeding to supplement stocking of natural oak has merit in spite of severe competition from natural regeneration."

Page 290 of 404

SEEDLING PLANTING -UPLANDS If hardwoods are to be planted in clearcuts, Johnson and Rogers (1982) recommended planting in the spring after harvest cutting to take advantage of favorable soil moisture and light conditions.

- **Site Preparation** Johnson and Rogers (1982) recommended the application of herbicides to increase planting success. Herbicides can be selected to control specific types/groups of vegetation (annual weeds, perennial grass and broadleaf species, and woody plants). The science of vegetation management has advanced since 1982 when Princep 80W¹, Tordon 101R¹ and Roundup¹ were the foundation of weed control in forestry. Products have become more species specific, have greater efficacy, and are more environmentally safe. Application technology minimizes drift to nontarget plants and is more effective at treating target plants with a minimal amount of chemical to be effective. "Although the effectiveness of one herbicide application in reducing competition is likely to be temporary, the impact on planted trees, especially slow-growing species like the oaks, may nevertheless be significant" (Johnson and Rogers 1982). Once established, oaks can become successful competitors.
- Number of Trees To
Be PlantedResults from West Virginia (Wendel 1980) indicate that after 7 years, 30 to 50
percent of the 2+0 red oak seedlings planted in a clearcut were judged successful.
This result suggests that 2 to 3.3 red oaks should be planted to obtain one
successful tree. This success ratio may be influenced by site, climate, competing
vegetation, and quality of the planting stock.
 - **Underplanting** Underplanting allows the root system of the planted trees to become well established before competition peaks after clearcutting. To be successful, both overstory and understory density control may be required to produce the light and soil moisture conditions necessary for adequate development of planted trees before the final harvest.

Johnson (1984) measured survival and height growth of northern red oak underplanted in the fall in mature stands and also planted in clearcuts. The overstory was removed between the third and fourth growing season. Among the underplanted trees, the 1+1 clipped seedlings grew the fastest. The greater growth for the clipped seedlings in the underplanting could be related to a more favorable root:shoot ratio.

The optimum length of time between planting and overstory removal is uncertain. Johnson speculated that, over long periods, mortality of underplanted trees would outweigh the growth advantage underplanting might provide and that a 2-3 year period would provide underplanted oak the time needed to increase its growth potential.

Tworkoski, Smith, and Parrish (1986) reported on seedling survival and growth of red and white oak underplantings in the Virginia Piedmont. Seedlings were underplanted followed by no canopy removal, partial removal (shelterwood), and

¹ Mention of trade names does not constitute endorsement of the product by the Department of Forestry and Natural Resources, Purdue University.

total removal (clearcutting). Seedling survival was high (90 percent). Growth increased with the degree of canopy removal.

Myers and others (1989) evaluated six types of northern red oak nursery stock planted in the understory of a mature stand of upland hardwoods in southern Indiana after 6 years. Overstory removal consisted of thinning from below to 60 percent stocking or a control. While thinning and understory control treatments resulted in the best survival (64 percent) and height growth (78 cm) it was not significantly different from the understory control treatment, i.e., understory control is important to the success of underplanted seedlings. Treatment differences associated with nursery stock type were nullified because of severe deer browsing.

Quality of Nursery Stock Johnson (1979) reported that 1+1 red oak transplants performed better than 1+0 seedlings even when the 1+0 seedlings were larger. Johnson suggested that undercutting seedlings in the nursery bed might result in the same advantages as observed with 1+1 transplants. The minimum size standard for northern red oak in the Lake States is: 50 cm (20 in.) height and 8mm (5/16 in.) caliper.

Stroempl (1985) described grading standards for northern red oak seedlings to be used in field plantings in southern Ontario. The grading criteria were based on root-collar diameter and stem length and stem form, bud number, and roots of large and small 2-0 nursery stock.

	Root-collar	Stem
Seedling size	diameter (mm)	length (cm)
Large	7.5-8.5	55-75
Small	4.5-6.5	30-45
Cull	<4.5	

Stem form. The stem should be well defined and straight, with relatively short branches on the previous year's growth. Long branches should be pruned and multiple leading shoots pruned to leave the sturdiest shoot on seedlings that have met all other criteria. Seedlings with multiple stems should be culled and the sources of acorns that do poorly identified and eliminated as a seed source.

Buds. The stem should have numerous buds, especially on the last year's growth, and a cluster of buds at the apex. Buds should not be swollen or flushed at the time of planting.

Roots. The "hockey stick" root which results from undercutting may make it more susceptible to root rot and may not properly support the developing crown. Root should be pruned to about 20-25 cm to facilitate field planting and ensure a high rate of survival.

Seed Source. Acorns should be collected from local sources and screened to select for adaptability and rapid juvenile growth.

Physiological Quality	Appropriate sized nursery stock has the potential for rapid root and shoot growth
of Nursery Stock	(Farmer 1975b, Johnson 1979) but this growth potential is often not realized

because of the poor physiological condition of the seedling. The physiological condition of nursery stock at planting is greatly influenced by the dormancy environment that immediately precedes planting.

Fall- and winter-lifted stock requires the proper storage or the physiological quality may decline. Seedlings should be kept cool $(1-5 \,^{\circ}C)$ and moist. The practice of heeling-in seedlings as a storage method is no longer recommended because of the inability to control root temperature and moisture conditions and the increased likelihood of the development of root diseases.

Spring-lifted seedlings and heeled-in seedlings are subject to desiccation resulting in shoot dieback. Because both shoot dieback and frost heaving may occur in fallplanted bareroot seedlings, Johnson (1979) recommends spring planting.

In oak, the winter shoot dieback is associated with reduced and delayed root growth capacity and reduced shoot growth (Johnson 1979). The cause of winter shoot dieback is linked to bud viability rather than the shoot, *per se*.

Buds play a major role in the initiation of root growth in oaks. Johnson and Rogers (1982) stressed the need to protect buds of dormant oak seedlings against both physiological and physical damage during lifting, handling, and storage. It is also important to plant trees as soon as possible after receiving them from the nursery and to do so with proper field handling and planting techniques (Johnson and Rogers 1982).

In 1971 Russell determined that root pruning and top clipping did not reduce the survival and growth of 1-0 northern red oak planted in bar-slits when compared to "ordinary nursery stock planted in the center-hole method."

Seedlings were root pruned to 5 in. and top clipped to 5 in. above the ground line. Ordinary seedlings had tap roots averaging 11 in. Toliver and others (1980) reported that after 5 years the combination of top pruning and root pruning of water oak and willow oak resulted in the best height growth, when compared to other combinations of pruning or no pruning and averaged 332 and 372 cm, respectively. Survival was 82 percent for water and 90 percent for willow oak.

The root system is believed to be the key to the above ground growth of planted seedlings (Farmer 1975a). Seedlings should have a favorable root:shoot ratio, and an abundance of first-order lateral roots, which are capable of developing new roots. Barden and Bowersox (1989) experimented with pre-sowing radicle root clipping and lateral root pruning in the nursery. When compared to the control, the combination of root treatments resulted in a doubling of the root growth capacity and larger diameter seedlings with greater leaf area.

Ruehle and Kormanik (1986) and Johnson (1989) have demonstrated that the persistent first-order lateral roots, stem height and diameter, and seedling vigor are important factors for the successful establishment of planted oak seedlings. Methods to produce high-quality seedlings in the nursery have been presented by Johnson (1988) and Crow and Isebrands (1986).

Ruehle and Kormanik (1986) stated that for oak seedling planting to be successful it was important "to identify at lifting time the individual seedlings that will develop an extensive root system after planting and grow well in height during the next 5 years." Gall and Taft (1973) and Webb (1969) cautioned on making early comparisons between seedlings since rankings often change after planting. More recently, Kormanik and others (1989) evaluated the frequency distribution of lateral roots of 1-0 bareroot white oak seedlings and correlated measurements of aboveground seedling growth in the nursery with lateral root development. Approximately 24 percent of the seedlings had 11 or more first-order lateral roots and offer the best chance for outplanting success (Kormanik and others, 1989). Evidence suggests that large oak seedlings with five or more lateral roots offer the potential for rapid growth (Schultz, personal communication). Others have not found this direct relationship (Kaczmarek 1991, Kaczmarek and Pope 1992).

The selection of genotypes with a high probability for rapid growth is still elusive. Maybe the phenotypic expression of the root system is a key. Techniques in DNA mapping and micropropagation provide new tools to identify and propagate superior genotypes.

Buchschacher and others (1991) recommended that oak seedling quality would be improved by using seed collected from superior trees or stands (registered for collection) rather than purchased from unrestricted public collections. Sowing beds by seed source and undercutting should decrease within nursery bed seedling variability and increase seedling quality and outplanting success.

Planting Prescription Johnson and others (1986) presented a four-step prescription for underplanting northern red oak in the Missouri Ozarks: (1) Control undesirable woody understory vegetation with herbicides before planting. (2) Create a medium density shelterwood. (3) Underplant large-diameter nursery stock (at least ½ in. diameter 1 in. above root collar) with clipped tops. (4) Remove the shelterwood three growing seasons after planting. Johnson and others stated that this prescription would result in the successful establishment of ½ to ½ of the planted trees 2 years after the shelterwood is removed.

On the average, shoot growth for planted upland oak seedlings is less than 10 cm per year. Typically, shoot growth of planted seedlings is restricted to the early part of the growing season when soil moisture is readily available. Weed control, including mulches, has generally not significantly improved oak seedling growth on forest sites (Wendel 1980, Hilt 1977). Fertilization has improved growth in some instances (Johnson 1980) but gains generally have not been large enough to justify the added cost. Approximately 10 percent of northern red oaks planted in forests grow 50 to 60 cm per year (Johnson 1976, Wendel 1980) which compares favorably to that of dominant natural reproduction in clearcuts on site indexes from 60 to 70.

Fall planting as a means of establishing seedlings prior to the spring growth period has been tried off and on for decades, always with mixed success. Johnson and Rogers (1982) reported that fall-planted container-grown seedlings had greater shoot growth and minimal shoot dieback when compared to 1-1 transplants and 1-0 seedlings respectively. In all cases the height growth of fall-planted seedlings exceeded that of spring-planted seedlings after 1 year. The minimal shoot dieback is particularly significant because gains made in shoot elongation are often negated by shoot dieback.

Containerized Seedlings

In 1978 only a small percentage of the more than 100 million container-grown seedlings produced were hardwoods. Container stock as large as a comparable bareroot seedling may cost four times as much; smaller container-grown seedlings may be as low in cost as bareroot seedlings (Tinus 1979).

Container stock costs more to transport and handle but the actual planting was easier and cheaper in some cases, i.e., when an automated tree-planting machine was used (developed by USDA Forest Service, San Dimas Equipment Development Center, J. L. Edwards). Hand-planting tools, which reduce planting effort and increase the planting rate by two to three times, have been developed for planting container-grown seedlings.

Container grown oak can be produced rapidly and survives well in the field (Tinus 1979). Container stock should be used on harsh sites, including south-facing slopes and sandy upland soils throughout the United States. Planting is most successful when the stock is dormant but ready to grow new roots. The keys to field performance are proper physiological condition, the right balance between shoot and root, and adequate size (Tinus 1979). The tree planter must have the proper equipment and must be motivated and know how to do a good planting job. Tinus (1979) concluded that it is the cost of the established seedling "free to grow" that counts, not the cost of the seedling at the nursery.

Marx (1979) introduced the idea of using specific mycorrhizal-forming fungi to Mycorrhizae and Oak improve the growth and physiological condition of oak seedlings produced in the Regeneration nursery and in containers. Marx projected the superior growth of these oak seedlings in field plantings. Based on his experience with loblolly pine (Pinus taeda L.), he was able to create an inoculation system that was successful in producing mycorrhizae on white oak (Marx 1979a) and red oak (Marx 1979b) seedling roots grown in the nursery. Other researchers (Dixon and others 1980, 1981; Garrett and others 1979) reported similar successes with black oak and northern red oak (Pope 1988, Dixon 1989). In all cases seedlings inoculated with the fungal species *Pisolithus tinctorius* were larger and had more root mass than control seedlings and those colonized by indigenous fungi. Similarly, fungal inoculations have been successful in forming mycorrhizae on a variety of oak species in container systems (Maronek and Hendrix 1979, Maronek and others 1981, Pope and Chaney 1985, Beckjord and others 1986, Kissee and others 1989, Reber 1991).

While fungal inoculation results in the formation of mycorrhizae and subsequent growth simulation of oak seedlings under control conditions, these same mycorrhizal seedlings rarely show any survival and growth advantages over control seedlings in field plantings. The early growth advantages of seedlings inoculated with mycorrhizal-forming fungi are often eclipsed by "non-inoculated" seedlings within 2 to 4 years after outplanting except on nutrient deficient and/or drought stressed sites (Beckjord and McIntosh 1983).

Tree Shelters Tree shelters are reported to increase height growth of planted 2-0 northern red oak (Lantagne and others 1990, 1991). Red oak seedlings planted in southern Michigan and grown in shelters were 53 cm taller than unsheltered trees after 3 years. The height growth of sheltered seedlings was related to the increased extension of the Page 295 of 404

initial growth flush without substantial increases in stem caliper and dry weight. This growth pattern is the major concern expressed for not using this technology. In Great Britain, however, the use of two shelters increased from 80 in 1979 to more than 6 million in 1986 (Potter 1987).

Case Studies Johnson (1980) described the 8-year growth response of five oak species planted in the Missouri Ozarks to fertilization with magnesium ammonium phosphate (8-40-0), a slow release amendment. The fertilizer was applied at the time of planting at a rate of 100 pounds of nitrogen and 500 pounds of phosphorus per acre. Fertilization increased height growth for black oak and scarlet oak but did not affect northern red oak, white oak, and post oak. Fertilization improved the survival of all species but black oak. In addition to an increase in mean height, fertilization increased the percentage of seedlings in the 6-, 8-, and 10-ft. height groups for black and scarlet oak and in the 10-ft. group for white oak. Competition control is essential to realize the growth potential from fertilizer applications (Johnson 1980).

> Based on the results of two long-term studies, McGee and Loftis (1986) concluded that planting oak on upland sites in North Carolina and Tennessee using conventional methods does not offer a solution to oak regeneration problems. In one study, 1-0 northern red oak seedlings were separated into three classes based on root collar diameter and planted in six clearcuts. Competing vegetation was controlled prior to planting and for 5 years thereafter. The site index for northern red oak was 75 to 98 (base age 50). Survival of the largest grade seedlings was slightly greater than grades 2 and 3, but height growth was the same. Survival averaged 88 percent after 5 years but declined rapidly after the cleanings stopped. By age 19, survival was 6 percent, height averaged 17.9 ft., and the free-to-grow percentage averaged 0.6.

> In another study located in the Cumberland Plateau (ridges and hollows) and Highland Rim (barrens and fragipans) of Tennessee, black oak seedlings were planted in clearcuts which had been chemically treated to control competing vegetation. Cleaning treatments were initiated when the planted trees were either 6, 7, or 8 years of age. Except for the Plateau hollows, cleanings had little beneficial effect on survival or height growth. Overall survival averaged 88 percent for five annual cleanings and 81 percent for uncleaned areas. Height after 12-14 years average 1.7 to 8.5 ft. and was related to site quality. Height growth of yellow poplar (*Liriodendron tulipifera* L.) planted in companion plots and not cleaned averaged 1.6 to 3.9 ft. per year.

McGee and Loftis (1986) concluded that, given the good early survival, planting could become a viable regeneration alternative for red and black oak if techniques could be developed to accelerate height growth.

Red oak seedlings (1-0 stock) were planted in an upland oak-hickory forest thinned to 60 percent stocking (Gingrich 1967). The competition was chemically treated prior to planting. Seedling survival was highest when the understory was controlled, but height growth was not affected. Deer browse was identified as a serious problem in this study and may have influenced the results in height growth (Myers and others 1989).

SEEDLING PLANTING -BOTTOMLANDS

Bottomland oak species are not satisfactorily regenerated after the harvest because small seedlings are not able to compete successfully with undesirable stump and root sprouts and/or faster growing seedlings (Johnson 1979, Johnson and Krinard 1976). Some regeneration problems can be solved with artificial regeneration Survival is often good but slow initial growth requires intensive control o competing vegetation, usually by cultivation, if planted seedlings are to develop rapidly and compete.

Seedling Morphology Research in artificial regeneration has focused on improving the early growth of outplanted seedlings. Early field performance was influenced by nursery practices and container production. Seedlings of acceptable size (2-1/2 to 3 ft. in height; 3/6 + in. root collar diameter) and vigor can be produced in the nursery in 1 year, provided that the nursery bed density is 10 seedlings per sq. ft. or less and the seeds are sown early enough in the spring (Hodges and Janzen 1987). Hodges and Janzen (1987) reported that undercutting and lateral root pruning increased root proliferation, but these results had no effect on height growth of outplanted seedlings.

Moorhead (1978, 1981) reported that containerized seedlings of bottomland red oaks had greater initial height growth after outplanting than nursery grown bareroot seedlings of the same age. The growth difference was not great enough to eliminate the need for competition control. The additional cost of growing containerized seedlings could not be justified in this study. Enrichment planting will require competition control in the lower canopy to successfully augment natural regeneration (Chambers and others 1987, Nix and Cox 1987). By 1986 research at the USDA Forest Service Laboratory at Stoneville had shown repeatedly that certain bottomland oaks can be successfully established via planting 1- to 3-year-old seedlings and controlling weeds.

Site Preparation Nix and Cox (1987) showed that preharvest competition control enhanced the probability of successful enrichment plantings with cherrybark oak. They evaluated overstory removal (clearcut and shelterwood) in combination with preharvest disking and post-planting, directed herbicide application. Disking increased survival and growth of seedlings planted in clearcuts. Similar results were evident in the shelterwood, but harvesting damage eliminated any preharvest disking treatment effect.

Nix (1989) evaluated the effect of mulching, and spring or fall application of herbicide on the survival and growth of 1-year-old cherrybark oak seedlings planted in a clearcut. Site preparation was done after clearcutting; no attempt was made to establish a cultivated plantation of oaks but rather to establish an enrichment planting. After six growing seasons, survival was greatest for the herbicide treatments (95 percent) followed by the mulch treatment (88 percent) and the control (75 percent). Treatments did not significantly affect height, or basal diameter, but all were significantly greater than the control. Spring herbicide treatment gave superior results.

Miller and Burkhardt (1987) tested herbicides and methods of application enrichment plantings of cherrybark oak. The trials were conducted on the loes bluff forests in western Mississippi. The test included injection of overtop

Page 297 of 404

trees and soil-active pelleted herbicides. After 3 years, the treatment of soil-active herbicide yielded seedling volumes significantly greater than those on the untreated plots.

Seifert and others (1985) evaluated the 5-year response of 1-0 swamp chestnut oak seedlings planted on a poorly drained upland flat in southeastern Indiana. Site preparation included disking and disking plus bedding. Chemical weed control was applied annually in the spring around all seedlings. Survival averaged 83 percent and was unaffected by treatment. Mean seedling height was greatest for the disked treatment. Fifty-two percent of the seedlings in the disked and 45 percent in the disked plus bedding treatments were greater than the overall mean seedling height, as compared to 16 percent for the control. Mechanical site preparation may be important for establishing oaks on poorly aerated loessial flats.

Plantation Management Plan

Based on 10 years of research, Malac and Heeren (1979) provided a plantation management plan for willow, water, and laurel oak in the southeastern United States. No speculation was made on the cost of implementing such a plan:

1. Seed should be collected from the best available local trees.

2. Sites selected for planting should have a sandy loam or loam surface fairly high in organic matter, and should have a water table within 4 in. of the surface during portions of the year.

3. Plant only large, healthy seedlings, with a minimum root collar diameter of **%** in. and a top height of at least 2 ft.

4. All sites should be intensively cleared, except for recently abandoned agricultural land. Some sites should be disked prior to planting, abandoned fields with plow pans or shallow topsoil should be subsoiled, and wet sites bedded.

5. Seedlings should be planted by a machine which has been modified to accommodate large seedlings.

6. Depending on site and weed growth, plantations should be disked two to three times a year for at least the first 2 years.

7. Fertilizer should be applied during the first cultivation; applications vary, but a general recommendation is 250 lb./acre triple superphosphate or diammonium phosphate.

CONCLUSIONS Regeneration of oaks by planting or direct seeding remains very uncertain. There are no foolproof methods or techniques that can guarantee success. The process, from seed collection to planting (seed or seedling) to young plantation management, is rife with pitfalls. The vagaries in weather, site conditions, and impacts of animals, insects, and disease all add to the uncertainty. Given all the potential problems, it is encouraging that successful artificial regeneration of oaks is occurring more frequently. The science/art of artificial regeneration has made significant advancements since 1971.

Research in direct seeding has provided methods that reduce animal predation of acorns and has identified oak species which are adaptable to a range of soil and site conditions. Research has led to better site selection for direct seeding and has described needed site preparation. For some oak species, information is available on season (time of year) and methods of sowing and spacing to achieve desired stocking levels. Research on seed collection and storage provides the knowledge Page 298 of 404

to ensure that, at planting, the seeds are sound, robust, and physiologically capable of expressing their growth potential.

Research in seedling production has explored use of mycorrhizal fungal symbionts to enhance seedling growth, root growth potential, plant water relations, and the physiological condition of the seedling. Growing oak seedlings in containers is possible, and using container-grown seedlings has merit on stressful sites. Root systems colonized with specific mycorrhizal fungi and/or container-grown seedlings have not proved to have the widespread advantages once proposed for oak regeneration.

Research efforts have focused on identifying the seedling characteristics that are associated with survival and adequate growth and creating favorable site conditions. Efforts have centered on root morphology, developing an adequate number of lateral roots, creating the proper top:root ratio, and producing a physiologically superior seedling. Similarly, research has investigated the impacts on site (site preparation and weed control) and canopy manipulations (clearcutting, shelterwood, thinnings) on oak seedling survival and growth. Great strides have been made in these areas, but the large within-species and within-family genetic variability have resulted in non-uniformity of response to management treatments.

Even with all the uncertainties inherent in the procedure, artificial regeneration offers the best solutions to many oak regeneration problems in the future. Clearly, continued failures of natural regeneration make it imperative to press forward for breakthroughs in genetics, economics, and the biology of early growth. Today, land managers should proceed with the planting and seeding of oaks, but they should consider all the information provided in this symposium and proceed with caution.

LITERATURE CITED Baker, J. B.; Broadfoot, W. M. 1979. A practical field method of site evaluation for commercially important southern hardwoods. Gen. Tech. Rep. SO-26. New Orleans, LA: U. S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 51 pp.

- Barden, C. J.; Bowersox, T. W. 1989. The effect of root pruning treatments on red oak seedling root growth. In: Rink, G.; Budelsky, C.A., eds. Proceedings, seventh central hardwood forest conference; 1989 March 5-8; Carbondale, IL. Gen. Tech. Rep. NC-132. St.Paul, MN: U.S. Department of Agriculture, North Central Forest Experiment Station: 115-119.
- Beckjord, P. R.; Melhuish, J. H., Jr.; Hacskaylo, E. 1986. Journal of Environmental Horticulture. 4(4):127-129.
- Beckjord, P. R.; McIntosh. M. S. 1983. Growth and fungal retention by field-planted Quercus rubra seedlings inoculated with several ectomycorrhizal fungi. Bulletin Torrey Botany Club. 110(3):353-359.
- Buchschacher, G. L.; Tomlinson, P. T.; Johnson, P. S.; Isebrands, J. G. 1991. In: Coleman, S. S.; Neary, D.G., eds. Proceedings, sixth biennial southern silvicultural research conference; 1990 October 30-November 1; Memphis, TN. Gen. Tech. Rep. SE-70. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 126-130.
- Chambers, J. L.; Stuhlinger, H. C.; Clifton, R. G. P. 1987. Regeneration of bottomland hardwoo sites by pre-harvest planting. In: Phillip, D. R., ed. Proceedings, fourth biennial souther silvicultural research conference; 1986 November 4-6; Atlanta, GA. Gen. Tech. Rep. SE-4

Asheville, NC: U. S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 125-128.

- Crow, T. R.; Isebrands, J. G. 1986. Oak regeneration-an update. In: Integrated pest management symposium of northern forests. 1986 March 24-27; Madison, WI: Cooperative Extension Service, University of Wisconsin-Extension: 75-82.
- Dixon, R. K.; Wright, G. M.; Behrns, G. T.; Teskey, R. O.; Hinckley, T. M. 1980. Water deficits and root growth of ectomycorrhizal white oak seedlings. Canadian Journal of Forest Research. 10(4):545-548.
- Dixon, R. K.; Wright, G.M.; Garrett, H.E.; Cox, G.S.; Johnson, P.S.; Sander, I.L. 1981. Container- and nursery-grown black oak seedlings inoculated with Pisolithus tinctorius: growth and ectomycorrhizal development during seedling production period. Canadian Journal of Forest Research. 11(3):487-491.
- Dixon, R. K. 1989. Inoculation of northern red oak seedlings with the fungal symbiont Suillus luteus in a Michigan nursery. In: Rink, G.; Budelsky, C.A., eds. Proceedings, seventh central hardwood forest conference; 1989 March 5-8; Carbondale, IL. Gen. Tech. Rep. NC-132. St. Paul, MN: U. S Department of Agriculture, Forest Service, North Central Forest Experiment Station: 141-145.
- Farmer, R. E., Jr. 1975a. Dormancy and root regeneration of northern red oak. Canadian Journal of Forest Research. 5(2):176-185.
- Farmer, R. E., Jr. 1975b. Growth and assimilation rate of juvenile northern red oak: effects of light and temperature. Forest Science. 21(4):373-381.
- Francis, J. K.; Johnson, R. L. 1985. Direct-seeded sawtooth oaks (Quercus acutissima Carruth.) show rapid growth on diverse sites. Tree Planters Notes. 36(3):3-5.
- Gall, W. R.; Taft, K. A., Jr. 1973. Variation in height growth and flushing of northern red oak (Quercus rubra L.). In: Proceedings, twelfth southern forest tree improvement conference: 190-199.
- Garrett, H. E.; Cox, G. S.; Dixon, R. K.; Wright, G. M. 1979. Mycorrhiza and the artificial regeneration potential of oak. In: Holt, H. A.; Fischer, B.C., eds. Regenerating Oaks in Upland Forests. West Lafayette, IN: Department of Forestry and Natural Resources, Purdue University: 82-90.
- Gingrich, S. F. 1967. Measuring and evaluating stocking and stand density in upland hardwood forests of the central states. Forest Science. 13(1):38-53.
- Hilt, D. E. 1977. Introduction of black walnut and northern red oak seedlings in an upland hardwood forest in southeastern Ohio. Res. Note NE-241. U. S. Department of Agriculture, Forest Service. 5 pp.
- Hodges, J. D.; Janzen, G. 1987. Studies on the biology of cherrybark oak: Recommendations for regeneration. In: Phillips, D. R., ed. Proceedings, fourth biennial southern silvicultural research conference. 1986 November 4-6; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U. S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 133-139.
- Johnson, P. S. 1976. Eight-year performance of interplanted hardwoods in southern Wisconsin oak clearcuts. Res. Pap. NC-126. St. Paul, MN: U. S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 9 pp.
- Johnson, P. S. 1979. Growth potential and field performance of planted oaks. In: Holt, H. A.; Fischer, B.C., eds. Regenerating oaks in upland hardwood forests. West Lafayette, IN: Department Forestry and Natural Resources, Purdue University: 113-119.

- Johnson, P. S. 1980. Response to fertilization of five oak species eight years after planting. Tre-Planters Notes. 31(1):9-10.
- Johnson, P. S. 1984. Responses of planted northern red oak to three overstory treatments. Canadian Journal of Forest Research. 14:536-542.
- Johnson, P. S.; Rogers, R. 1982. Hardwood interplanting in the upper Mississippi valley. In Proceedings, hardwood regeneration conference. 1982 March 2-3; Winona, MN.: Available fron University of Minnesota, College of Forestry, Minneapolis, MN: 90-109.
- Johnson, P. S.; Dale, C. D.; Davidson, K. R.; Law, J. R. 1986. Planting northern red oak in the Missouri Ozarks: A prescription. Northern Journal of Applied Forestry. 3:66-68.
- Johnson, P. S. 1988. Undercutting alters root morphology and improves field performance o northern red oak. In: Proceedings, tenth north american forest biology workshop. 1988 July 19-22: Vancouver, B.C.: University of British Columbia: 316-323.
- Johnson, P. S. 1989. Growing hardwood nursery stock for planting on forest sites with specia reference to northern red oak. In: Proceedings, 1989 Northeastern area nurserymen's conference 1989 July 24-27. U. S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry: 46-61.
- Johnson, R. L. 1979. Adequate oak regeneration -- a problem without a solution. In: Management and Utilization of Oak. Proceedings, of the seventh annual hardwood symposium hardwood research council. Asheville, NC: 59-65.
- Johnson, R. L. 1983. Nuttall oak direct seeding still successful after 11 years. Res. Note SO-301. New Orleans, LA: U. S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 3 pp.
- Johnson, R. L. 1984. Direct-seeded cherrybark and shumard oaks battle natural regeneration through 10 years. Southern Journal of Applied Forestry. 8(4):226-230.
- Johnson, R. L.; Krinard, R. M. 1976. Hardwood regeneration after seed tree cutting. Res. Pap. SO-123. New Orleans, LA: U. S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 7 pp.
- Johnson, R. L.; Krinard, R. M. 1985. Regeneration of oaks by direct seeding. In: Proceedings, third symposium of Southeastern hardwoods. 1985 April 16-17; Dothan, AL. Atlanta, GA: U. S. Department of Agriculture, Forest Service, Southern Region: 56-65.
- Johnson, R. L.; Krinard, R. M. 1987. Direct seeding of southern oaks--a progress report. In Proceedings, fifteenth annual hardwood symposium of the hardwood research council. 1987 May 10-12; Memphis, TN. Broomall, PA: U. S. Department of Agriculture, Forest Service: 10-16
- Kaczmarek, D. J. 1991. Seedling morphology related to growth and survival in northern red oal (Quercus rubra L.). West Lafayette, IN. Purdue University. 176 pp. Thesis.
- Kaczmarek, D. J.; Pope, P. E. 1992. Covariate analysis of northern red oak seedling growth. In Proceedings, seventh biennial Southern silvicultural research conference. 1992 November 17-19 Mobile, AL. (Accepted).
- Kissee, K. K.; Garrett, H. E.; Pallardy, S. G.; Reid, R. K. 1989. Effects of shading and mycorrhizae on the growth and development of container-grown black and English oak seedlings. In: Miller, J. H., ed. Proceedings, fifth biennial Southern silvicultural research conference; 198: November 1-3; Memphis, TN. Gen. Tech. Rep. SO-74. New Orleans, LA: U. S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 63-66.
- Kormanik, P. P.; Ruehle, J. L.; Muse, H. D. 1989. Frequency distribution of lateral roots of 1-6 bare-root white oak seedlings. Res. Note SE-353. Asheville, NC: U. S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 5 pp. Department of A04

Page 301 of 404

- Lantagne, D. O.; Ramm, C. W.; Dickmann, D. F. 1990. Tree shelters increases heights of planted oaks in a Michigan clearcut. Northern Journal of Applied Forestry. 7(1)24-26.
- Lantagne, D. O. 1991. Tree shelters increase heights of planted northern red oaks. In: McCormick, L. H.; Gottschalk, K.W., eds. Proceedings, 8th central hardwood forest conference. 1991 March 3-6; Pennsylvania State University, University Park, PA. Gen. Tech. Rep. NE-148. Radnor, PA: U. S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 291-298.
- Malac, B. F.; Heeren, R. D. 1979. Hardwood plantation management. Southern Journal of Applied Forestry. 3(1):3-6.
- Maronek, D. M.; Hendrix, J. W. 1979. Growth acceleration of pin oak seedlings with a mycorrhizal fungus. Horticulture Science. 14(5):627-628.
- Maronek, D. M.; Hendrix, J. W.; Stevens, C. D. 1981. Fertility- mycorrhizal isolate interactions in production of containerized pin oak seedlings. Scientia Hortic. 15:283-289.
- Marx, D. H. 1979. Ectomycorrhizae: Rx for artificial regeneration of oak? Proceedings, seventh annual hardwood symposium of the hardwood research council; 1979 May; Cashiers, NC: 66-71.
- Marx, D. H. 1979a. Synthesis of Pisolithus ectomycorrhizae on white oak seedlings in fumigated nursery soil. Res. Note SE-280. Asheville, NC: U. S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 4 pp.
- Marx, D. H. 1979b. Synthesis of ectomycorrhizae by different fungi on northern red oak seedlings. Res. Note SE-282. Asheville, NC: U. S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 8 pp.
- McGee, C. E.; Loftis, D. L. 1986. Planted oaks perform poorly in North Carolina and Tennessee. Northern Journal of Applied Forestry. 3:114-115.
- Miller, J. H.; Burkhardt, E. C. 1987. A trial of herbicide treatments for enrichment plantings of cherrybark oak. In: Hay, R. L.; Woods, F.W.; DeSelm, H., eds. Proceedings, central hardwood forest conference, VI; 1987 February 24-26; Knoxville, TN. Knoxville, TN: University of Tennessee, Department of Forestry, Wildlife and Fisheries: 53-58.
- Moorhead, D. J. 1978. Effects of container size and growth media on the early growth and field survival of four southern red oak species. Starkville, MS: Mississippi State Univ. 75 pp. Thesis.
- Moorhead, D. J. 1981. Container size and growth media influence early growth and survival of southern oaks in Mississippi. In: Workshop on seedling physiology and growth problems in oak planting; 1979 November; Columbia, MO. Gen. Tech. Rep. NC-62. St. Paul, MN: U. S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 20 pp.
- Myers, R. K.; Fischer, B. C.; Wright, G. M. 1989. Survival and development of underplanted northern red oak seedlings: 6-year results. In: Rink, G.; Budelsky, C.A., eds. Proceedings, seventh central hardwood forest conference; 1989 March 5-8; Southern Illinois University, Carbondale, IL. Gen Tech. Rep. NC-132. St. Paul, MN: U. S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 150-155.
- Nix, L. E.; Cox, S. K. 1987. Cherrybark oak enrichment plantings appear successful after 7 years in South Carolina bottomlands. In: Phillips, D. R., ed. Proceedings, fourth biennial southern silvicultural research conference. 1986 November 4-6; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U. S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 129-132.
- Nix, L. E. 1989. Early release of bottomland oak enrichment plantings appears promising in South Carolina. In: Miller, J. H., ed. Proceedings, fifth biennial southern silvicultural research conference; 1988 November 1-3; Memphis, TN. Gen. Tech. Rep. SO-74. New Orleans, LA: U.
 S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 379-383.

- Pope, P. E.; Chaney, W. R. 1985. Influence of *Pisolithus tinctorius* and fertilization of development of container grown red oak seedlings. In: Shoulders, E., ed. Proceedings, biennial southern silvicultural research conference. 1984 November 7-8; Atlanta, GA. Gen. 7 Rep. SO-54. New Orleans, LA: U. S. Department of Agriculture, Forest Service, Sou Forest Experiment Station: 403-409.
- Pope, P. E. 1988. Pisolithus tinctorius increases the size of nursery grown red oak seedlings. Forests. 2:5-16.
- Potter, M. 1987. Advances in tree shelter research and design. Oxford Forestry Institute Occas papers No. 34. National Hardwoods Programme. Report of the Seventh Meetings. P.S. Savil 7 pp.
- Reber, R. T. 1991. Quercus rubra L. and ectomycorrhizae: Natural colonization in field plan and fertilization effects on specific fungal species and growth parameters in a greenhouse st West Lafayette: Purdue University. 100 pp. Thesis
- Ruehle, J. L.; Kormanik, P. P. 1986. Lateral root morphology: A potential indicator of see quality in northern red oak. Res. Note SE-344. New Orleans, LA: U. S. Departmer Agriculture, Forest Service, Southern Forest Experiment Station. 6 pp.
- Russell, T. E. 1971. Seeding and planting upland oaks. In: Oak Symposium Proceedings. August; Morgantown, WV. Upper Darby, PA: U. S. Department of Agriculture, Forest Ser-Northeastern Forest Experiment Station: 49-54.
- Schultz, R. C. 1992. Lateral root morphology and the growth of northern red oak. Pers Communication.
- Seifert, J. R.; Pope, P. E.; Fischer, B. C. 1985. The effects of three levels of site preparation planted swamp chestnut oak on a poorly drained site. In: Shoulders, E., ed. Proceedings, t biennial southern silvicultural research conference; 1984 November 7-8; Atlanta, GA. Gen. T Rep. SO-54. New Orleans, LA: U. S. Department of Agriculture, Forest Service, South Forest Experiment Station: 53-56.

Stroempl, G. 1985. Grading northern red oak planting stock. Tree Planters Notes. 36(1):15-18

- Tinus, R. W. 1979. Use of container-grown hardwoods for reforestation. In: Holt, H. A.; Fisc B.C., eds. Regenerating Oaks in Upland Forests. 1979 February 22-23; West Lafayette, Department of Forestry and Natural Resources, Purdue University: 120-125.
- Toliver, J. R.; Sparks, R. C.; Hansbrough, T. 1980. Effects of top and lateral root pruning survival and early growth on three bottomland hardwood species. Tree Planters No 31(3):13-15.
- Tworkoski, T. J.; Smith, D. W.; Parrish, D. J. 1986. Regeneration of red oak, white oak, and w pine by underplanting prior to canopy removal in the Virginia Piedmont. Southern Journa Applied Forestry. 10:206-210.
- Webb, C. D. 1969. Uniform seedling density is important in hardwood progeny test nurseries Proceedings, tenth southern conference on forest tree improvement; 1969 June 17-19; Hous TX. Publ. No. 30. So. For. Tree Improv. Comm.: 208-216.
- Wendel, G. W. 1980. Growth and survival of planted northern red oak seedlings in West Virg Southern Journal of Applied Forestry. 4(1):49-54.

Artificial Regeneration of Bottomland Oaks

Harvey E. Kennedy, Jr., Southern Forest Experiment Station, USDA Forest Service, Stoneville, MS 38776

ABSTRACT Research results from the Southern Hardwoods Laboratory, Stoneville, Mississippi, on direct-seeding and planting oaks is now being applied to establish commercial plantings. The Louisiana Department of Wildlife and Fisheries has successfully planted and direct-seeded over 4,000 acres during a 5-year period. Primary species were Nuttall, water, willow, and Shumard oaks. The Mississippi Department of Wildlife Conservation is well into a 10-year program of direct-seeding oaks on the Malmaison Game Management Area. Large-scale plantings (seedlings) and direct-seeding of oaks are being conducted under the Conservation Reserve Program in Louisiana, Arkansas, Mississippi, Tennessee, and Alabama.

Over 10 years, the growth and development of saplings of four oak species planted at five spacings in a minor stream bottom in southeast Arkansas showed significant differences among species and spacings. Spacing affected all tree-size and biomass variables but had no effect on survival. Water oak developed most rapidly; swamp chestnut oak most slowly.

Measurements made on a 27-year-old water oak plantation near Winnsboro, Louisiana, and a 20-year-old cherrybark oak plantation near Vicksburg, Mississippi, showed that oaks can be successfully grown in plantations.

Successful plantings from direct-seeded Nuttall, water, cherrybark, and Shumard oaks have been established.

The importance of site-species selection, site productivity, soil water, effects of flooding depth and duration, and site preparation is emphasized. Basics of planting and direct-seeding, including seedling and acorn quality, spacing, storage and handling, and hand versus machine planting and direct-seeding, are discussed.

INTRODUCTION

Bottomland hardwood forests provide sites for some of the most productive timber and wildlife habitat in the United States. Additional benefits are improved water quality from forests that act as filter zones along streams and restoration of deteriorated sites. Such sites support two to five times as many game animals as do upland pine sites. Hard-masted species, such as oak and pecans, provide extremely important resources for wildlife. Deer, turkey, rabbits, squirrels, migratory ducks, and wood ducks are taken in great numbers each year from hardwood forests (Allen and Kennedy 1989). Before the first European settlers arrived on the North American continent, de hardwood forests and cypress-tupelo forests covered the floodplains of major a minor stream bottoms, mountainous coves, and the swamps along coastal are More than 100 species were represented (Freeman and Frome 1968). T tremendous agricultural potential of the areas with their rich soils and abund moisture, especially those along rivers and streams, was realized alm immediately. Gradually, as flooding was brought under control and drain projects were established, increasingly more hardwood forests were cleared agriculture.

By the late 1930's, about half the original eastern hardwood forest had be cleared. Bottomland clearing continued at a rapid pace through the late 1970 especially in the lower Mississippi River Valley. It is now apparent that ma areas that were cleared would be better suited as forests because they are subj to flooding and can successfully produce an agricultural crop only in relatively c years (Allen and Kennedy 1989). These areas, and many other forests where oak component is not present, can only be reforested to oaks by using artific regeneration. Artificial regeneration of bottomland oaks can be accomplished the South using direct-seeding or planting methods developed at the Southe Hardwoods Laboratory in Stoneville, Mississippi. Landowners who wish reforest oaks artificially must make several decisions that reflect a high level commitment to meeting requirements.

PLANNING AN ARTIFICIAL REGENERATION PROJECT

Careful planning is the first and most critical step in a regeneration project (All and Kennedy 1989). Landowners must decide on location and site preparati needs for the planting or direct-seeding site, tree species, quantity of seedlings seeds needed to complete the project, and the mode of planting or seeding. The decisions must be made well in advance of the planting or direct-seeding da (preferably as much as a year) to ensure availability of planting or seeding stoc labor, and equipment.

Proper tree species selection greatly affects the probability of success of t project. Selection must be based on factors such as soil type, frequency a duration of flooding (table 1), flooding depth, site-species adaptability, availabil of planting stock, and objectives for regeneration. In many planting projects in t lower Mississippi Valley, emphasis has been placed on hard mast-producing speci such as oaks (*Quercus* spp.) and sweet pecan (*Carya illinoensis* (Wangenh.) Koch), for both timber production and wildlife. Light-seeded species such sweetgum (*Liquidambar styraciflua* L.), sycamore (*Platanus occidentalis* L.), a the ashes (*Fraxinus* spp.) usually become established on the site because their see are easily dispersed by wind and water. Planting technology for other hardwoo has also been developed.

The two major reforestation methods are direct-seeding and planting seedlings, bd of which can be done by hand or machine. Planting includes 1-0 seedlings and older, larger seedlings where fewer seedlings per acre are planted as enrichmer or supplemental plantings. Two large, well-planned seeding and planting program are those of the Louisiana Department of Wildlife and Fisheries and the Corps

- Continuous flooding Periodic flood		- Periodic flooding		
January-June	January-May	January-May	January-April	January-March
Cypress	Green ash	Sweetgum	Sawtooth oak	Shumard oak
Overcup oak	Nuttall oak	Water oak	Sycamore	Cherrybark oak
Water hickory	Persimmon	Willow oak	Cottonwood	Swamp chestnut
·		Nuttall oak	Sweet pecan	oak
		Green ash	Nuttall oak	Nuttall oak
			Green ash	Green ash

Table 1-Species tolerance in relation to flooding time and duration

Engineers Lake George project. The Louisiana group has direct-seeded and planted over 4,000 acres during a 5-year period in old fields in the Ouachita Wildlife Management Area near Monroe, Louisiana. The Lake George project is an 8,800acre area in west-central Mississippi that will be planted and direct-seeded over a 5-year period. Direct-seeding oaks is usually less expensive than planting oak seedlings, but costs depend on species, seed price, labor, availability of seedlings, and other factors. Direct-seeding costs per acre on old fields are estimated at about 40 percent those of planting seedlings (Bullard and others 1992). Also, directseeding has been successful only with hard-mast species such as oaks, pecan, and hickories (*Carya* spp.) (Johnson and Krinard 1987). Bare-root seedlings would have to be planted for the light-seeded species. Planted oak seedlings have faster establishment and growth rates through the first several years (Allen 1990, Wittwer 1991).

- **E PREPARATION** Site preparation may be necessary to create suitable growing conditions or to facilitate the use of mechanical planters or seeders. If the area has been in cultivation for a long time, site preparation should break up any plow pan or otherwise compacted soil and aid in weed control during the growing season after planting. Disking with a heavy disk at least twice late in the summer before planting is recommended. Disking should be to at least a 6-in. depth and preferably 8 to 15 in.
 - **SITE-SPECIES SELECTION** Growth of bottomland hardwoods depends on the physical condition of the soil, moisture availability during the growing season, nutrient availability, and aeration (Baker and Broadfoot 1979). Bottomland oaks grow best on moist, well-drained sites with good fertility and medium-textured soils. They will grow on the finer textured clays, but survival will probably be lower and growth will be slower. Growth is best in pH ranges of 6.0 to 7.0, but good survival and growth have been obtained with oaks in pH ranges as low as 5.0 to 5.4 (Kennedy 1985, Kennedy and others 1987). Soil pH above 7.5 limits growth and survival of several red oak species, although Shumard oak (*Q. shumardii* Buckl.) grows well at these high pH ranges (Kennedy 1984, Kennedy and Krinard 1985).

County soil surveys contain a wealth of information on soil characteristics (including degree of flooding and soil saturation) that affect the survival and growth Page 306 of 404

of trees, as well as information on the suitability of the soil type(s) on the site various tree species (Allen and Kennedy 1989).

Another good source of information is local expertise. Nearby landowners I know of changes in flooding patterns (due to drainage, etc.) that are not cove in the soil survey. They may also know of other planting or direct-seeding projuin the area and can relate any problems that were encountered.

DIRECT-SEEDING Oak seeding research at the Southern Hardwoods Laboratory was begun in early 1960's. Direct-seeding is becoming a widely used method of reforestati particularly in the lower Mississippi River Valley. Since 1981, some 5,000 10,000 acres of publicly and privately owned land have been direct-seeded v acorns or other heavy mast species.

Many early tests were with Nuttall oak (Q. nuttallii Palmer). The slow growth this species (and of most oaks) in the first 5 years may be discouraging beca competing sprouts and seedlings of other species outgrow the oaks. Howev within 10 to 15 years, enough oaks should be in a competitive crown positior ensure their presence in the future sawtimber stands (Johnson and Krinard 198 Although Nuttall has been the most popular red oak to direct-seed and has highest success rate, at least six other red oak species, four white oak species, i sweet pecan have been successfully direct-seeded (Johnson and Krinard 198 Species selection should be based on site quality and landowner preference.

Guidelines for direct-seeding Nuttall oak also apply to other hard mast spec however, lower germination rates can be expected with other oaks and pec Because white oaks germinate in the fall and the acorns are hard to store, aco should be direct-seeded as soon as possible after collection.

Collecting and Storing Seeds Acorn collecting is the biggest job in a direct-seeding program. In the Sou collections are normally done between October 1 and January 1. Fallen acorns quickly eaten by birds, deer, and rodents and the time when acorns drop varies species and individuals within a species, so one needs to keep checking the to (Johnson and Krinard 1987).

Early seeding eliminates long-term storage that requires cold space and monitor Freshly collected acorns seem to be less attractive to rodents if they are see immediately; unlike acorns that have been in cold storage, they do not exude oc that attract rodents (Johnson and Krinard 1987).

Many oaks produce an excellent seed crop only 1 or 2 years out of every 5. The for storable red oak acorns (Bonner 1973), extra collection may be required du good seed years. Storage may be expensive, but it is more reliable and product than collecting acorns during poor crop years. It allows for an annual see program even in years of total acorn crop failure.

Practical time limits for storing large quantities of red oak acorns may be less the 5 years, as indicated by research (Johnson and Krinard 1987). In Louis about 5,300 lbs. of Nuttall oak acorns were collected during fall-winter 198 Page 307 of 404 stored in a walk-in cooler at 35 °F to 40 °F, and used during the 1985, 1986, and 1987 planting seasons (about 30 months of storage). However, if they are not properly stored, acorns can lose viability after 1 year. In the Louisiana study, the acorns were stored in polyethylene bags with about 20 to 25 lbs. of seed per bag. This made handling easier and kept seeds from heating in the bag (Johnson and Krinard 1987). The bags were placed in cold storage on shelves to allow for air circulation. If weevils were present in the acorns, they would emerge, eat through the bag, and fall to the floor from the wire shelves, thus minimizing damage to the bags.

Collecting. As a general rule, acorns of white oak cannot be stored more than 3 to 4 months because they normally germinate in the fall. Rink and Williams (1984) suggest that acorns of Q. alba L. be stored in 1.75-mil polyethylene bags instead of the 4-mil bags suggested for red oak acorns because of lower germination after storage in 4-mil bags. Suspected causes of lower germination were higher concentrations of ethanol and carbon dioxide.

Labeling. It is important to label storage bags to identify species, collection date, and percent soundness.

eeding Operations Water (*Q. nigra* L.), cherrybark (*Q. falcata* var. *pagodifolia* Ell.), Shumard, and Nuttall oak acorns can be removed from cold storage and sown with good results even in the early summer. The least desirable months for planting in the South and Southeast are July through October, when the soil is normally hot and dry (Johnson and Krinard 1987). Seeds that germinate in cold storage will produce seedlings when field-sown even if their radicles are broken off (Bonner 1982). Acorns of site-suitable species can be sown in flood areas soon after the water recedes—usually in April, May, or June—and should germinate and produce healthy seedlings.

> Old fields have been hand- and machine-sown with and without site preparation, but seeding is easier and generally more successful on intensively prepared sites. Mechanical seeders cannot operate effectively in areas after a harvest cut because of the resulting slash and stumps.

> Machines used for seeding are actually modified one- and two-row soybean seeders. Some drop acorns automatically; others are ridden by operators who drop acorns at specified distances. Seeders that drop seeds automatically should be checked at the end of each row to make sure the hopper does not jam, leaving long distances unseeded. Seeds sown 1 to 6 in. deep will germinate and produce seedlings. Seeding at greater depths slows germination and emergence of the young shoot; about 2 in. is the recommended depth. Reported direct-seeding rates range from 30 to 40 acres per day with a three-person crew using machines, to 6 to 8 acres per person with hand seeding. Normal spacing is rows 10 to 12 ft. apart and acorns 3 to 5 ft. apart within rows, or 1,000 to 1,500 acorns per acre. Research at the Southern Hardwoods Laboratory has shown a high rate of success when direct-seeding is done in complete openings greater than 100 ft. on a side. Direct-seedings in smaller openings have generally failed as a result of rodent damage (Johnson and Krinard 1987).

Germination as high as 80 percent has been reported in some research t (Johnson and Krinard 1987). However, about 35-percent germination will be n typical for commercial sowings. Thus, 1,000 to 1,500 acorns per acre cal expected to produce 300 to 500 1-year-old trees, which would be a suffic number for most objectives.

On the basis of limited data (Johnson and Krinard 1987, Bullard and others 19 it appears that seedlings can be established 1 year after direct-seeding at a lo cost than planting seedlings. However, there are so many influencing factors reliable cost comparisons would involve a thorough economic analysis.

Some situations may not be conducive to direct-seeding. Failures may occur un adverse field environments (such as free water coupled with high temperature deadly environment for newly germinated acorns), extended periods of drou during the growing season, residual chemicals in old fields, presence of anir (especially raccoons), and defective acorns or improper storage techniques.

Post-Direct-Seeding Operations Most land managers have not attempted to control weeds around direct-seeded c in old fields (Johnson and Krinard 1987). In some research trials, bushhogg between rows of seedlings appeared to improve seedling survival and growth. one is willing to accept a little lower survival and growth, competition control r not be necessary in natural stand openings until free-to-grow oaks are 15- to 20 tall (Johnson 1981). Under normal conditions, a site with an 80- to 90-ft. index should produce 15- to 20-ft. tall trees in 10 to 15 years. At that time necessary, individual oaks can be released by deadening or by cutting compet trees.

PLANTING OAK SEEDLINGS Planting bottomland oak seedlings is a reliable method of reforestation. When d correctly, the chances for successful establishment of a forest plantation are l (Allen and Kennedy 1989, Kennedy and others 1987, Kennedy 1984). If planting is done incorrectly or carelessly, however, the project could be expensive failure. Keys to successful initial establishment of seedlings, in addi to selecting the appropriate species for the site, are obtaining good-qua seedlings, ensuring good preplanting care of seedlings, and using proper plan procedures.

> The primary sources of tree seedlings are State forestry organizations and pri nurseries. The seedlings available from most nurseries are bare-rooted—separ from the soil they were grown in at the nursery. Bare-rooted seedlings are expensive than container-grown seedlings, lighter, easier to transport, and ea to plant.

> In contrast to pine seedlings, which are usually planted while still quite small seedlings should have a top length of at least 18 in. and a root-collar diamete at least 3% in. Roots should be well developed and can be pruned to a leng about 8 in. to make planting easier.

Bare-rooted seedlings require careful handling; the roots are very tender and Seedlings generally come from the nursery wrapped in bundles of 50 to 200. If the seedlings are not to be planted immediately, they should be stored (still bundled) in a cool, dark place-ideally in a cold-storage unit. A barn, shed, or dense shade will do for a few days, as long as the roots do not freeze or dry vulnerable. out.

moist sphagnum or peat moss. They should never be carried in the hand while immediately or transferred in small groups to a bucket or planting bag containing planting; doing so will expose roots to the drying effects of the air and sun, and They should either be taken out of the nursery-supplied bundles and planted Only as many seedlings as can be planted in 1 day should be taken to the field. roots will dry very quickly. The best time to plant bare-rooted oak seedlings is while they are dormant and when the soil is moist-generally, from January through March in the South. Planting can be extended into May if the seedlings are kept in cold storage until planting.

However, if the goal is to reforest primarily for wildlife, spacings of 12 by 12, 15 Spacings of 10 by 10 ft. or 12 by 12 ft. are recommended for timber production. by 15, or even 20 by 20 ft. may meet the objective.

shovel. It is important that the roots be placed in the hole so they can spread out somewhat naturally-they should not be twisted, balled up, or bent. Moist soil Seedlings can be hand planted using a dibble bar, tile spade, or sharpshooter should be packed firmly around the roots.

collar just below the soil surface. Making holes too shallow for proper placement or, more commonly, not deep enough. Seedlings should be planted with the root of roots causes "J-rooting," which can result in malformed root systems. Another mistake is leaving an air pocket around the roots after closing the hole, which may Perhaps the most frequent planting mistake is planting the seedling either too deep allow the roots to dry out. experienced person can plant 600 to 800 seedlings per day if planting conditions are relatively good. This rate can vary substantially, however, for a number of reasons, including degree of site preparation, temperature, and the size of the seedlings being planted. One

Machine planting is usually much faster than hand planting. An experienced crew of two or three can plant 4,000 to 10,000 seedlings per day with a planting Often survival will be better than that achieved by hand planting. machine.

> Early Plantation Development

Outright failures and partially successful oak plantations occur occasionally. Seedling mortality and poor growth result from extended post-planting dry periods, flooding coupled with high temperatures, poor planting or seeding practices, residual herbicides or herbicide drift from nearby aerial applications, poor-quality seed or seedlings, or development of the new plantation. The landowner should closely monitor development of the Problems are likely to show up in the first 2 years after planting. animal depredation. Early weed control (by disking, mowing, or use of herbicides) will speed gree of seedlings during the first few years and perhaps slightly increase survival, the benefits may not justify the costs. Too much weed control might acture reduce the short-term value of the site for some types of wildlife that use the w for food or cover.

There is no cost-effective method to protect newly established seeds or seedl from animal predation. Domestic animals, birds, deer, rabbits, squirrels, raccobeavers, nutria, and mice may all destroy seeds or seedlings. Fencing to cor domestic animals and some wildlife, and good site preparation to clear grc cover, can reduce the number of rodents. Protection of individual seedling justified only where there is a high population of nutria. In such a case, chicl wire predator guards are essential. Some animal damage should be expected tolerated.

As long as the survival rate is satisfactory, landowners should not worry if the grow slowly at first (Allen and Kennedy 1989). Most species of oak grow onl to 2 ft. per year the first several years. After this stage, growth can increase to 4 ft. per year if seedlings are dominant and not overtopped. By the end of years, trees should be from 15 to 25 ft. tall or more and constitute a valual productive forest.

A number of successful oak plantations established by direct-seeding and plant seedlings have been reported. Johnson and Krinard (1987) reported results of oak plantation established by direct-seeding: after 10 to 15 years, oaks were 1. 20 ft. tall with about 250 trees per acre in the dominant/codominant position a free-to-grow condition. After 10 growing seasons, four oak species establis with 1-0 bareroot seedlings in a minor stream bottom in Arkansas ranged from in. d.b.h. and 12.9 ft. high for swamp chestnut oak to 3.1 in. d.b.h. and 23.9 high for water oak (Kennedy and others 1987). After 27 years, water plantings on an old-field loessial soil in Louisiana had per-acre stand values follows: number of trees, 356; average d.b.h., 6.6 in.; basal area, 86 ft²; t volume of trees ≥ 5.0 in. d.b.h., 2,017 ft²; average height of 50 tallest trees, ft. (Krinard and Johnson 1988). Planting cherrybark oak on abandoned field the loess hills appears to be a reliable method for establishing a valuable component (Krinard and Francis 1983). At age 20, cherrybark averaged 346 tr per acre. Diameters of the dominant and codominant trees ranged from 6.2 to in. d.b.h. and heights from 55 to 65 ft.

OUTLOOK The oak component in certain bottomland areas can be increased by direct-seed or planting. Oaks likely to be seeded or planted successfully include Nut water, cherrybark, Shumard, willow (*Q. phellos L.*), white (*Q. alba L.*), swamp chestnut (*Q. michauxii* Nutt.). Successful regeneration can be enhanced careful matching of species and site and rigorous attention to meeting preparation needs. If the procedures and techniques recommended in this pape not followed, costly failures are likely. The long-term outlook is for an increase of bottomland to be successfully regenerated to oak.

ERATURE CITED Allen, J. A. 1990. Establishment of bottomland oak plantations on the Yazoo National Wildlife Refuge Complex. Southern Journal of Applied Forestry. 14(4):206-210.

- Allen, J. A.; Kennedy, H. E., Jr. 1989. Bottomland hardwood reforestation in the Lower Mississippi Valley. Bull. Slidell, LA: U.S. Department of Interior, Fish and Wildlife Service, National Wetlands Research Center, and Stoneville, MS: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 28 pp.
- Baker, J. B.; Broadfoot, W. M. 1979. Site evaluation for commercially important southern hardwoods. Gen. Tech. Report SO-26. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 51 pp.
- Bonner, F. T. 1982. The effect of damaged radicles of presprouted red oak acorns on seedling production. Tree Planters Notes. 33(4):13-15.
- Bonner, F. T. 1973. Storing red oak acorns. Tree Planters Notes. 24(3):12-13
- Bullard, S.; Hodges, J. D.; Johnson, R. L.; Straka, T. J. 1992. Economics of direct-seeding and planting for establishing oak stands on old-field sites in the South. Southern Journal of Applied Forestry. 16(1):34-40.
- Freeman, O. L.; Frome, M. 1968. National Forests of the United States. New York: G.P. Putnam's Sons (eds.) and Waukesha, WI: Country Beautiful Foundation, Inc. 194 pp.
- Johnson, R. L.; Krinard, R. M. 1987. Direct-seeding of southern oaks-a progress report. In: Proceedings of 15th annual hardwood symposium of the hardwood research council; 1987 May 10-12; Memphis, TN. Memphis, TN: Hardwood Research Council: 10-16.
- Johnson, R. L. 1981. Oak seeding--it can work. Southern Journal of Applied Forestry. 5(1):28-33.
- Kennedy, H. E., Jr.; Krinard, R. M.; Schlaegel, B. E. 1987. Growth and development of four oaks through age 10 planted at five spacings in a minor stream bottom. In: Proceedings of the 9th annual southern forest biomass workshop; 1987 June 8-11; Biloxi, MS. Mississippi State, MS: Mississippi State University: 81-91.
- Kennedy, H. E., Jr. 1985. Cultural treatments influence hardwood growth and foliar nutrient concentration on a minor stream bottom site. Res. Pap. SO-215. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 6 pp.
- Kennedy, H. E., Jr.; Krinard, R. M. 1985. Shumard oaks successfully planted on high pH soils. Res. Note SO-321. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 3 pp.
- Kennedy, H. E., Jr. 1984. Hardwood growth and foliar nutrient concentrations best in clean cultivation treatments. Forest Ecology and Management. 8(1984):117-126.
- Krinard, R. M.; Johnson, R. L. 1988. Stand parameters of a 27-year-old water oak plantation on old field loessial soils. Res. Note SO-348. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 pp.
- Krinard, R. M.; Francis, J. K. 1983. Twenty-year results of planted cherrybark oak on old fields in brown loam bluffs. Tree Planters Notes. 34(4):20-22.
- Rink, G.; Williams, R. D. 1984. Storage technique affects white oak acorn viability. Tree Planters Notes. 35(1):3-5.
- Wittwer, R. F. 1991. Direct seeding of bottomland oaks in Oklahoma. Southern Journal of Applied Forestry. 15(1):17-22.

Artificial Regeneration of Oaks in the Uplan

Todd W. Bowersox, School of Forest Resources, Pennsylvania State University, University Park, PA 1

ABSTRACT Opportunities for planting and seeding oaks in upland sites have never been grea Land managers are making strong commitments to regenerating oaks, and artifi regeneration can be an important contribution to a successful operation. Di seeding and planting of oaks can be used to regenerate upland mixed oak sta that have had high mortality from gypsy moth and/or to augment nat regeneration in harvested stands. Results from northern red oak direct seeding planting studies at upland sites in Pennsylvania are summarized. Direct seedin recommended only if your are certain that there will be no predation by sr mammals. My current recommendation for planting seedlings on upland foree sites is to require appropriate planting procedures, acceptable seedling quality, freedom from overstory and understory competition.

INTRODUCTION Development of effective methods for establishing oak seedlings from artifi regeneration on upland forested sites has been hindered by many factors affect seed germination and seedling growth. The main factors are: (1) overstory t competition or interference; (2) understory vegetation interference; (3) predat by large and small omnivores; (4) physical environment; and (5) insects diseases. Each factor has an impact on the successful establishment of a seedli However, many of these are interrelated, and when working in concert cro conditions that are unfavorable to establishing oak seedlings in acceptable t frames.

> The literature on seed storage, seed germination, and seedling growth of oaks upland sites is abundant. A variety of regeneration methods or guidelines h been developed for regenerating oak stands. Some of these include planting acorns (Johnson and Krinard 1985) or seedlings (Johnson and others 1986), most assume natural regeneration (Gottschalk 1983, Hannah 1987, Roach Gingrich 1968, Sander 1988, Watt 1979). To optimize these methods, we nee understand acorn storage and germination requirements. Burns and Honkala (19 summarized the flowering, fruiting, seed production, dissemination, and seed development knowledge for the oaks. Cecich (1991) has summarized understanding of seed production in oaks. Schopmeyer (1974) provided abu general and oak-specific seed biology and development information.

McQuilkin (1983) concluded that the principal requirements for obtaining successful oak regeneration are moist, friable soils and acorns protected from desiccation and freezing. For acorns not buried in the soil, a cover of leaf litter is essential. Johnson (1970) and Shipman (1962) reported that depth of sowing influenced germination rate and success. Johnson (1967) also found lower germination success for Nuttall oak acorns planted in bare soil than those planted in litter covered soil. These investigators cited soil temperature and moisture as factors controlling germination rate and germination success. Swank and Vose (1988) reported on changes that cutting practices have on the micro-environment of the forest floor and soil. They concluded that the temporal changes in the physical, chemical, and biological properties of the forest floor and soil were important in controlling the success of hardwood regeneration. Little is known about how changes in the forest floor properties may affect the establishment and initial growth of artificially regenerated oaks.

Interference from competing vegetation is probably the most consistent factor contributing to oak regeneration failures in old fields (Bey and others 1975, Hilt 1977, von Althen 1977). In recent years, chemical weed control programs have been developed for establishing several valuable hardwood species on old field sites (Holt and others 1981). These control programs employ a variety of pre- and post-emergence herbicides to gain effective weed control. Typically these programs involve the elimination of existing annual and perennial vegetation before planting using broad-spectrum, foliar applied herbicides, such as glyphosate. Soil active pre-emergence herbicides may also be applied either before or after seedling planting to provide continued weed control throughout the growing season. Effective herbicides include simazine, oryzalin, diphenamid, napropamide, oxadiazon, oxyfluorfen, and metolachlor (Sam and others 1985) and sulfometuron methyl (McCormick and others 1991, Groninger and McCormick 1991, Horsley and others 1992). The specific herbicide or herbicide combinations used in a weed control program depends upon the sensitivity of the planted species to the herbicide, nature of the vegetation to be controlled, and soil-site conditions.

Stress of herbaceous vegetation on hardwood regeneration in forested areas is being recognized (Bowersox and McCormick 1987, Horsley 1982, Horsley 1988, Kolb and others 1989, Kolb and others 1990, McCormick and Bowersox 1989). Regenerating harvested forest lands with desirable timber species is frequently hindered by the presence of competing herbaceous vegetation. These competitors can assume a dominant role in areas where they were either absent or sparse before overstory removal. Some ferns and grasses can rapidly capitalize on changes in site conditions created by overstory removal. Unfortunately the actual factor(s) (i.e., increased light, nutrients, or moisture) responsible for establishing herbaceous species are not well understood. Equally uncertain is the degree that herbaceous vegetation will contribute to the survival and growth of oak seedlings.

There has been no great enthusiasm for artificially regenerating oak in Pennsylvania. Oak stands that have been the product of direct-seeding or planting are rare in Pennsylvania. Successful practices to direct-seed or plant oak seedlings developed for other locations (Johnson and Krinard 1985, Johnson and others 1986) could be used to revegetate or avoid natural regeneration failures. In mixed oak stands impacted by gypsy moth, oak seedlings accounted for 4 to 18 percent of the natural regeneration (Allen and Bowersox 1989). The Allegheny National Fc has a northern red oak planting program to restore oaks to about 2,000 acres have suffered high overstory mortality following gypsy moth defoliations (Crot 1992).

In 1984, Penn State started a research program to examine some of the fac controlling the establishment and growth of northern red oak (*Quercus rubra* We restricted our studies to mixed oak stands growing on better-than-average quality (oak site indices >70) which have little or no natural oak regenerat Our goal has been to determine the steps necessary to achieve seedlings that are m tall in 4 years or less.

This paper summarizes the results of our direct seeding and planting on upl sites in Pennsylvania. Recent information on germination and seedling gro from direct seeding, and tree improvement programs, nursery practices, plan procedures, seedling cultural practices, and methods will be summarized.

DIRECT-SEEDING Seeding acorns has been widely recommended as a way to regenerate oaks. Gra (1910) recommended planting acorns "in their permanent place" because of difficulty in transplanting oaks.

Direct-seeding of oaks has the attraction of requiring less time, effort, and expe than the production and planting of seedlings. Acorn storage, seed quality, tim of the seeding, and placement of the acorns are well known elements that influe establishing seedlings. Davidson (1988), Johnson and Krinard (1985) and Tec and Isebrands (1991) have provided recent summaries of artificially regenerat oaks. Our studies with seeded northern red oak acorns resulted in seedlings averaged 15 to 20 cm high after one growing season, and 20 to 50 cm after growing seasons (Kolb and others 1989, Kolb and others 1990). Prelimin results at the Harry's Valley and Sand Knob 1 locations (figure 1) have shown shade and competition from fern and grass communities can have a major affect seedling growth. We direct-seeded northern red oak to evaluate the effect different nursery and planting practices on seedling survival and growth (Zad and others 1991). The study was conducted in clearcut mixed oak stands at the locations: Michaux 1, Harry's Valley 2, and Potter (figure 1). Over all location the first-year average height of the standard 1-0 bareroot seedlings was about cm compared to about 20 cm for the direct-seeded seedlings. The bare seedlings averaged about 20 cm when planted. At the end of three grov seasons, there was no difference in the average height of seedlings from the dir seeding (65 cm) and those from the bareroot planting (70 cm).

Anyone who has tried to grow oaks from seed has experienced acorn losse small mammals. Small mammals can impact a seeding program by destroyin removing the acorns or by clipping the developing seedlings. Table 1 is a 1 the small mammals that have major impacts in Pennsylvania. Impact of mammals is difficult to predict and control. In regard to natural regene Healy (1988) discussed seed predation by birds and mammals and conclud losses are not sufficient to cause natural regeneration failures. He suggest over a long period of time, there should be sufficient supplies of acorns to meet the needs of wildlife and to accumulate adequate numbers of advanced oak seedlings to regenerate a stand. However, I believe that predation pressure on direct seeded acorns will be more intense than on natural supplies because of fewer acorns sown and availability of alternate acorns is low.




Figure 1-Locations of oak seeding and planting studies in Pennsylvania.

Table 1—Small mammals that can have major impacts on oak regeneration in upland sites of Pennsylvania

Common name	Scientific name	Туре
Mice		
White-footed mouse	(Peromyscus leucopus)	Omnivore
Voles		
Meadow vole	(Microtus pennsylvanicus)	Herbivore
Southern red-backed vole	(Clethrionomys gapperi)	Omnivore
Chipmunks		
Eastern chipmunk	(Tamias striatus)	Omnivore
Squirrels		
Fox squirrel	(Sciurus niger)	Omnivore
Gray squirrel	(Sciurus carolinensis)	Omnivore
Southern flying squirrel	(Glaucomys volans)	Omnivore

We experienced losses to small mammals that varied from none to complete disappearance of nearly all acorns planted. In one seeding attempt, we repeatedly experienced total losses overnight. In another study, we had greater variability in germination success among three locations than with whether or not the acorns were protected from small mammals (Zaczek and others 1991). Over all three locations, 59 percent of the unprotected acorns germinated, compared to 70 pe of the protected acorns. Only 29 percent of all acorns planted at a sou location (Michaux 1) germinated, whereas 75 and 91 percent of the a germinated at the central (Harry's Valley 2) and northern (Potter) locat respectively.

DeLong (1992) recently studied the predation on direct-seeded acorns. She observed the losses of direct-seeded acorns in relation to time of placement, c of placement, and overstory stand condition. This study with unprotected nort red oak acorns in mixed oak stands (Sand Knob 2, 3, and 4) has found that no all acorns seeded never develop into seedlings. Eight hundred seed spots located in uncut (100 percent stocked), improvement (70 percent stocked) shelterwood (20 percent stocked) treatment areas. Each of the 2,400 seed s were seeded in October and examined seven times between the seeding and of the following year. Acorns removed or destroyed were immediately repl with another sound acorn. In a year following low acorn production (1990) direct-seeding produced four seedlings by the end of one growing season. year following average acorn production (1991), the direct-seeding resulted in seedlings at the end of one growing season. The losses were similar for ac placed on the surface and those planted 2 cm into the soil. The highest losses v in the shelterwood treatment, with lower losses in the improvement treatment uncut stand. Based on the evidence at the seed spot, most of the acorns were to white-footed mice (Peromyscus leucopus).

Not willing to risk the success of our research efforts to the chance of preda by small mammals, we developed two methods for protecting acorns: A siprotector was designed for establishing a single seedling from spring seed (figure 2); a larger protector was developed for studies requiring minidisturbance to the litter and soil and for use with fall seedings (figure 3).

The single seedling protector was made from 2.5-cm diameter, schedule 40 I pipe that is cut lengthwise. The bisected pipe was cut into 12-15 cm sections, a cross-cut at the top and a 45° angle cut at the bottom. Angle cuts made it ea to insert the protector into the soil. The two halves have been secured at on two points with one and one-half wraps of masking tape. The protector inserted at least 10 cm into an augured hole. An acorn was added, covered suitable material to conserve acorn moisture and a 4-cm nail was place across top to prevent extraction of the acorn. This type of protector has been successfully with northern red oak, white ash, and white pine (Kolb and or 1990). Small mammals will excavate protectors that are placed < 10 cm into ground or protectors that are loosely inserted into the soil. This protector has susceptible to over-winter frost heaving in waterlogged soils.

The larger protector was made from 160 cm diameter ASTM 2729 PVC pip in to 12 to 15-cm lengths. The pipe was inserted at least 5 cm into the Insertion into rocky, root-laden soil has been accomplished by 160-cm rin to the desired depth, and then the pipe was pushed into the slit. After the p




Figure 2-Individual acorn protector used at the Pennsylvania State University.




Figure 3-Large acorn protector used at the Pennsylvania State University.

in place, the acorns were planted and the top was covered by 1.3-cm mesh that was attached to the top of the pipe at three or four points. Gaps between pipe lip and the mesh had to be <1.5 cm to exclude the mice-sized anin Finally, the protector was pinned to the soil by three to four wires, 20-25 cm le This protector has permitted us to conduct phytotoxicity tests on seeds germinating seedlings with soil-applied herbicides (McCormick and others 19 We are using this protector to evaluate the effects of varying amounts of li acorn placement, and overstory conditions on the storage, germination, and gro of northern red oak from fall plantings. This protector has been susceptibl winter frost heaving in waterlogged soils and has not protected the acorns f losses to raccoons, black bears, or humans.

We are trying to develop a degradable acorn protector. One device has wor well in a trial. The trial was conducted in the same stands as the acorn preda study (Sand Knob 2, 3, and 4) in the year with an average acorn crop. Degrada protectors containing northern red oak acorns were planted 5 cm deep in Octo 1990. In August 1991, each protector was evaluated. About 44 percent of protectors produced a seedling (table 2). This was lower than the >70 perc germination success expected from the PVC protectors but greater than the < percent success from the unprotected acorns in the predation study.

Fate	Number	Percent
Successful seedling	66	44
Protector removed	13	9
Germinated but no seedling	39	26
No apparent germination	32	21

 Table 2—Fate of northern red oak acorns planted in a prototype degradable seed protector¹

¹ Seeding was done in October and placed 2-3 cm below soil surface.

All small mammals listed in table 1 have the potential to feed on newly germina shoots. Meadow voles, southern red-backed voles, and white-footed mice are m likely to clip off new shoots than the other species. In some cases the shoots consumed, in other cases the shoots are simply clipped off and not eaten. likely that much of the small mammal clipping of newly emerged oak shoots been blamed on deer, insects, or disease. A careful examination shortly after clipping is necessary to determine the precise cause of a decapitation to a ne emerged oak shoot. Stormer (1968) studied the shoot clipping habits white-footed mice. He concluded that the clipping activity was more relate their propensity to gnaw and not related to their needs for food or water.

In November 1989, we direct-seeded northern red oak into shelterwood (20 p stocking) and improvement (70 percent stocking) treatments made in a mix stand (Sand Knob 2 and 3). Most of the acorn protectors were distur humans and then pilfered by small mammals. From a limited database for

a year of low acorn production, we observed frequent clipping of the emerging shoots by small mammals. In the 1990 growing season, 46 percent of the acorns that were going to germinate had germinated by June 1. Small mammals clipped off one-half of these seedlings. By June 21, 73 percent of the potential germinants had shoots and still about one-half of these were clipped off by small mammals. Clipping activity dropped sharply after July 1. The study was duplicated with November seedings in 1990. Small mammals clipped about 1 percent of the potential germinants in the improvement area and about 5 percent of the potential germinants in the shelterwood area. Most of the clipping activity was before July 1. All clipping action was a single event; we recorded no multiple clippings.

Studies examining the germination and growth of direct-seeded northern red oak at the Moshannon location were also conducted in the 1990 and 1991 growing seasons. These studies were in a mixed oak stand that had clearcut and uncut treatments. Germination success in 1990 averaged about 60 and 70 percent for the clearcut and uncut areas, respectively. Clipping by small mammals was <1percent in the clearcut area. In the uncut area, the June 25 inventories recorded about 25 percent of the emerged shoots were clipped by small mammals. Replicated direct-seedings into new treatment areas in 1991 had germination success of about 75 and 85 percent in the clearcut and uncut areas, respectively. Again there were <1 percent of the seedlings clipped off by small mammals in the clearcut areas. Clipping by small mammals dropped to <10 percent in the uncut area. Preliminary data for 2 years after seeding indicated there was no difference in survival or average height of the clipped and unclipped seedlings.

Small mammals can be a major factor in determining the success of oak directseeding on upland sites. From our experiences, small mammals will either consume or move most unprotected direct-seeded acorns. I am convinced that most direct-seeding of acorns in upland sites will fail to produce seedlings unless the acorns are protected from small mammals. In addition, preliminary information from our studies indicate that of those acorns that germinate some will be clipped off by small mammals.

The clipping has been limited to a one-time decapitation, usually early in the growing season. I do not believe this clipping activity will be a major problem unless there would be multiple clippings in the growing season.

PLANTING In 1984 we planted 1-0 bareroot northern red oak seedlings at three locations (Harry's Valley 1, Laurel Run, and Sand Knob 1). This project was designed to evaluate the effect of established ferns and grass communities on the survival and growth of oak seedlings. Overall survival and growth rates were encouraging (George and others 1991). Four-year survival was 58 percent for seedlings growing in the presence of ferns and grasses and 81 percent for those growing free from fern and grass interference. Average 4-year height was 1.1 and 1.3 m for the weedy and weed-free conditions.

Root system morphology and structure make it difficult to produce easily transplantable oak seedlings. Nursery practices of varying plant densities,

undercutting the root, and lateral root pruning have been used to reorganize t root systems to increase survival and growth rates. It is generally agreed th post-planting growth rates are dependent on the ability of the seedling to devel a root system that has established good contact with the soil water and nutric resources. Ruehle and Kormanik (1986) have found that lateral root systems vari among families, and the number of large first-order lateral roots were correlat with after-planting growth potential. Schultz and Thompson (1991) suggest that least six first-order lateral roots are needed for acceptable survival and growth oak seedlings planted in the prairie region of the Central States. Barden a Bowersox (1989, 1990) reported that clipping the newly emerged radical will crea a fourfold increase in the number of "taproots." We also found that new ro development was mainly from the cut ends of any of the larger root tissue Furthermore, Barden and Bowersox (1991) found that 1-0 seedlings that we established in the nursery with the radical clipped also performed better wh out-planted than seedlings with unclipped radicals.

We are concluding a study that evaluated a variety of nursery stock types at thr locations (Michaux, Harry's Valley 2, and Potter). Location had a strong effe on the 3-year-old height. Over all treatments, average height at the Harry's valle location of 1.3 m was greater than the average height at Michaux (0.6 m) at Potter (0.9 m). One half of the seedlings for all treatments were top-clipped to 10 cm height at planting. Clipping resulted in greater 3-year height growth that the unclipped seedlings, but there was essentially no difference in the total height at the end of three growing seasons. Over all treatments, 2-year-old stock had a average height of 1.2 m. This was 62 percent greater than the average 3-year-ol height of the 1-year-old stock. Undercutting in the nursery bed produced variab results but was most beneficial for the 2-year-old stock. Greatest 3-year heigh occurred in 2-0 container stock, which averaged 1.5 m tall. We concluded that the 2-0 bareroot seedlings would be an acceptable balance between seedling production planting limitations, and out-planting performance variables for planting norther red oak seedlings in the upland conditions of Pennsylvania. Annual undercuttir the seedlings will produce a more plantable root system and should improve the out-planted height growth.

Planting oak seedlings in upland sites in Pennsylvania has been an endurance te for planting crews. Most of the stands have between 10 and 90 percent of ti surface covered by rocks. The KBC bar has been the best choice for hand plantin and will create and close a 16-cm deep hole on most sites. However, this plantin method restricts the size and arrangement of the root systems that can be plante Roots longer than 16 cm and wider than 10 cm needed to be removed befo planting. Tree planters would discard seedlings with the larger, more develop root systems because they could not make a hole big enough to accommodate the root systems.

These soil conditions and the resulting planting procedures were seriously limit seedling characteristics. One-person augers have been used successfully planting trees in the hard, rocky soils in the inter-mountain region of the Ro Mountains. We also found these augers worked for planting oak in Pennsylvania uplands. We use a 10-cm, carbide-tipped auger powered chainsaw engine with a slow speed gearhead. We auger a 30-cm deep pl hole. Loose soil that falls back into a 30-cm hole is easily removed by hand before inserting the roots. One or two additional holes are needed to accumulate sufficient soil to plant the seedling. This method of planting is being used to plant northern red oak seedlings on about 2,000 acres in the Allegheny National Forest (Crothers 1991).

Pennsylvania's Bureau of Forestry and Pennsylvania State University have started a cooperative genetic improvement program for northern red oak (Bailey and others 1992). The objective of the program is to identify families that can achieve a 1.4m height in 3 years after out-planting. The study is being conducted at three locations (Michaux 2, Tuscarora, and Moshannon 2). All seedlings are plantings in recently clearcut mixed oak stands, maintained weed free, and fenced to exclude deer. The 2-year-old survival values for all families planted in 1990 were >96 Over all families, average height was greatest at percent at all locations. Tuscarora (0.8 cm), moderate at Michaux 2 (0.5 cm), and lowest at Moshannon 2 (0.4 cm). Family means ranged from 0.37 to 0.5 cm at Moshannon 2 to 0.6 to 1.0 cm at Tuscarora. These results were for seedlings planted as 1-0 bareroot stock, and 1991 was a very dry year in Pennsylvania. Progeny from about 40 parent trees have been planted as 2-0 stock in 1991 and 1992 and will be planted in 1993 and 1994. In total, we plant to screen about 180 parent trees.

SUMMARY AND CONCLUSIONS

Research programs have been aggressively studying direct seeding and planting of oak throughout the eastern hardwood region. This is particularly true for northern red oak. We have gained valuable foundational information on the elements affecting germination and seedling growth from direct seeding. To establish one oak seedling from direct-seeding, I recommend planting two to three viable acorns per spot—two acorns for spring seedings, three for fall seedings. Each acorn should be planted 25-50 cm deep in the soil. Overstory and understory competition needs to be reduced to a level that will permit the seedling to compete with the other ground-level vegetation. I do not recommend direct-seeding acorns in upland forest sites unless you are certain that there will be no small mammal predation or the acorns will be protected from predation.

Successful establishment and growth of oak seedlings from planting require appropriate planting procedures, acceptable seedling quality, and freedom from overstory and understory competition. There is considerable variability in size and root characteristics of oak seedlings that are available at any one time. Smaller seedlings being planted in unrestricted soils can effectively be planted with bars, whereas larger seedlings being planted in rocky soils are best planted with an auger. With all things being equal, I recommend planting seedlings that are at least 30 cm tall, have a root collar of 6 mm, and have a multiple-branched, 25-cm-long root system. Seedlings with these minimum characteristics are best planted with an auger, especially in rocky upland forest soils. Overstory and understory competition should be reduced to a level that the seedlings are capable of successfully competing with the understory vegetation. In most cases, this would be to reduce the overstory to <50 percent of crown cover and remove all over-topping understory vegetation. Tree improvement programs, nursery practices, and planting procedures provide additional information to have more successful planting programs. knowledge of the cultural requirements to gain acceptable post-planting see height growth rates is developing to the point were prescriptions will soo formulated. Refinements and adjustments will be needed for specialized situa and for individual species. We are ready to move to a higher leve understanding of the factors controlling successful artificial regenerating oal upland conditions. However, production-scale artificial regeneration trials needed to test the practical application of our current knowledge. Implement of the current technology is key to the future of developing practical artificial artificial artificial regeneration programs for oaks in the uplands of eastern North America.

LITERATURE CITED

- Allen, D.; Bowersox, T. W. 1989. Regeneration in oak stands following gypsy moth defoli Gen. Tech. Rep. NC-132. U. S. Department of Agriculture, Forest Service: 67-73.
- Barden, C. J.; Bowersox, T. W. 1989. The effect of root pruning treatments on red oak seedling growth capacity. In: Proceedings, 7th central hardwoods forest conference. Gen. Tech. Rep. 132. U. S. Department of Agriculture, Forest Service: 115-119.
- Barden, C. J.; Bowersox, T. W. 1990. Genotype and radical clipping influence northern red oal growth capacity. In: Abstracts of Proceedings, fourth workshop on seedling physiology growth problems in oak plantings. Gen. Tech. Rep. NC-139. U. S. Department of Agricu Forest Service. 2 pp.
- Barden, C. J.; Bowersox, T. W. 1991. Effects of radicle clipping on subsequent growth of rea seedlings in high and low moisture environments. In: Proceedings, 6th southern silvicu research conference. Gen. Tech. Rep. SE-70. U. S. Department of Agriculture, Forest Ser 131-137.
- Bailey, J. K.; Steiner, K. C.; Bowersox, T. W.; Zaczek, J. J. [in press]. 1992. In: Abstrac Proceedings, fifth workshop on seedling physiology and growth problems in oak planting;
- Bey, C. F.; Krajicek, J. E.; Williams, R. D.; Phares, R. E. 1975. Weed control in hard plantings. In: Proceedings, herbicides in forestry, J. S. Wright forest conference. West Lafa IN: Purdue University: 69-84.
- Bowersox, T. W.; McCormick, L. H. 1987. Herbaceous communities reduce the juvenile grov northern red oak, white ash and yellow- poplar but not white pine. In: Proceedings, c hardwood forest conference VI. Knoxville TN: University of Tennessee: 39-43.
- Burns, R. M.; Honkala, B. H., technical coordinators 1990. Silvics of North Americ Hardwoods. Agriculture Handbook 654. U. S. Department of Agriculture, Forest Servic pp.
- Cecich, R. A. 1991. Seed production in oak. In: Proceedings, the oak resource in the upper mi Implications for Management. Pub. Number NR-BU-5663-S. St. Paul, MN: Univer Minnesota: 125-131.

Crothers, C. 1991. Tree shelters help ANF protect oak stand. Pennsylvania Forest. 82(4):2

Davidson, W H. 1988. Potential for planting hardwoods in the Appalachians. In: Proce guidelines for regenerating Appalachian hardwood stands. Morgantown, WV: West University: 255-268.

- DeLong, C. A. 1992. Effects of small mammals on direct seeded northern red oak. MS thesis, in progress. The Pennsylvania State University, University Park, PA.
- George, D. W.; Bowersox, T. W.; McCormick, L. H. 1991. Effectiveness of electric deer fences to protect planted seedlings in Pennsylvania. In: Proceedings, 8th central hardwood conference. Gen. Tech. Rep. NE-148. U. S. Department of Agriculture, Forest Service: 395-401.
- Gottschalk, K. W. 1983. Management strategies for successful regeneration in oak-hickory. In: Proceedings, regenerating hardwood stands. University Park, PA: Pennsylvania State University: 190-213.
- Graves, H. S. 1911. Forest planting leaflet red oak (Quercus rubra). Circular 58. U.S. Department of Agriculture, Forest Service. 3 pp.
- Groninger, J. W.; McCormick, L. H. 1991. Effects of sulfometuron on hay-scented fern spore emergence. Canadian Journal of Forest Research. 21:942-943.
- Hannah, P. R. 1987. Regeneration methods for oaks. Northern Journal of Applied Forestry. 4:97-101.
- Healy, W. M. 1988. Effects of seed-eating birds and mammals on Appalachian hardwood regeneration. In: Proceedings, guidelines for regenerating Appalachian hardwood stands. Morgantown, WV: West Virginia University: 104-111.
- Hilt, D. E. 1977. Introduction of black walnut and northern red oak seedlings in an upland hardwood forest in southeastern Ohio. Res. Note NE-241. U. S. Department of Agriculture, Forest Service. 5 pp.
- Holt, H. A.; Fisher, B. C.; Siefert, J. R. 1981. Current and new technology for controlling weeds in central hardwood forests. In: Proceedings, weed control in forest management. J. S. Wright forest conference. West Lafayette, IN: Purdue University: 139-147.
- Horsley, S. B. 1982. Development of reproduction in Allegheny hardwood stands after herbicide-clearcuts and herbicide-shelterwood cuts. Res. Note NE-308. U. S. Department of Agriculture, Forest Service. 4 pp.
- Horsley, S. B. How vegetation can influence regeneration. 1988. In: Proceedings, guidelines for regenerating Appalachian hardwood stands. Morgantown, WV: West Virginia University: 38-55.
- Horsley, S. B.; McCormick, L. H.; Groninger, J. W. 1991. Effects of thinning of oust applications on survival of hardwood seedlings. Northern Journal of Applied Forestry. 22-27.
- Johnson, P. S.; Dale, C. D.; Davidson, K. R.; Law, J. R. 1986. Planting northern red oak in the Missouri Ozarks: a prescription. Northern Journal of Applied Forestry. 2:22-26.
- Johnson, R. L. 1967. Improved germination of Nuttall oak acorns. Res. Note SO-66. U. S. Department of Agriculture, Forest Service. 3 pp.
- Johnson, R. L. 1970. Field germination of Nuttall oak acorns. Res. Note SO-101. U. S. Department of Agriculture, Forest Service. 4 pp.
- Johnson, R. L. 1984. Direct-seeded cherrybark and Shumard oaks battle natural regeneration through 10 years. Southern Journal of Applied Forestry. 8:226-231.
- Johnson, R. L; Krinard, R. M. 1985. Regeneration of oaks by direct seeding. In: Proceedings, third symposium of Southeastern hardwoods. Dothan, AL and Atlanta, GA: U. S. Department of Agriculture, Forest Service, 56-65.

- Kolb, T. E.; Bowersox, T. T.; McCormick, L. H. 1990. Influences of light intensity weed-induced stresses of tree seedlings. Canadian Journal of Forest Research. 20:503-507
- Kolb, T. E.; Bowersox, T. W.; McCormick, L. H.; Steiner, K. C. 1989. Effects of shade herbaceous vegetation on first-year germination and growth of direct-seeded northern red white ash, white pine and yellow-poplar. In: Proceedings, seventh central hardwood confere Gen. Tech. Rep. NC-132. U. S. Department of Agriculture, Forest Service: 156-161.
- McCormick, L. H.; Bowersox, T. W. 1989. Artificial regeneration of oaks in Pennsylvan Problems and potential. In: Abstracts of Proceedings, fourth workshop on seedling physiology growth problems in oak plantings. Gen. Tech. Rep. NC-139. U. S. Department of Agricul Forest Service. 22 pp.
- McCormick, L. H.; Allen, D. H.; Groninger, J. W.; Bowersox, T. W. 1991. Effect sulfometuron on the emergence and growth of seeded northern red oak, white ash, and v pine. Northern Journal of Applied Forestry. 8(1):9-11.
- McQuilkin, R. A. 1983. Silvical factors affecting the regeneration of oaks and associated speci Pennsylvania. In: Proceedings, regenerating hardwood stands. University Park, PA: Pennsylv State University: 37-65.
- Roach, B. A.; Gingrich, S. F. 1968. Evenaged silviculture for upland central hardwoods. Hand 335. U. S. Department of Agriculture, Forest Service. 39 pp.
- Ruehle, J.L.; Kormanik, P. P. 1986. Lateral root morphology: a potential indicator of see quality in northern red oak. Res. Note SE-344. U.S. Department of Agriculture, Forest Ser 6 pp.
- Sam, A. W.; Bowersox, T. W.; McCormick, L. H. 1985. Preemergence herbicide performanforest soils. In: Proceedings, fifth central hardwood conference. Urbana-Champaign University of Illinois: 47-52.
- Sander, I. L. 1988. Guidelines for regenerating Appalachian oak stands. In: Proceedings, guide for regenerating Appalachian hardwood stands. Morgantown, WV: West Virginia Unive 189-198.
- Schopmeyer, C. S. 1974. Seeds of woody plants in the United States. Agric. Handbook 450. Department of Agriculture, Forest Service. 883 pp.
- Shipman, R. D. 1962. Nursery-seeded hardwoods--influenced by depth and density of sowing Planters Notes. 54:27-32.
- Stormer, F. A. 1968. Factors affecting the distribution and abundance of Peromyscus leucop Clethrionomys gapperi in mixed oak forests of central Pennsylvania. University Pennsylvania State University. 83 pp. Thesis.
- Schultz, R. C.; Thompson, J. R. 1991. The quality of oak seedlings needed for successful a regeneration in the central states. In: Proceedings, the oak resource in the upper m Implications for Management. Publication Number NR-BU-5663-S. St. Paul, MN: Unive Minnesota: 180-186.
- Swank, W. T.; Vose, J. M. 1988. Effects of cutting practices on microenvironment in rel hardwood regeneration. In: Proceedings, guidelines for regenerating Appalachian ha stands. Morgantown, WV: West Virginia University: 71-88.
- Teclaw, R. M.; Isbrands, J. G. 1991. Artificial regeneration of northern red oak in the Lak In: Proceedings, the oak resource in the upper midwest. Implications for Manageme Number NR-BU-5663-S. St. Paul, MN: University of Minnesota: 187-197.

von Althen, F. W. 1977. Hardwood planting in Ontario. Forest Chronicle. 53(4):1-6.

- Watt, R. F. 1979. The need for adequate advanced regeneration in oak stands. In: Proceedings, regenerating oaks in upland hardwood forest, J. S. Wright forest conference. West Lafayette, IN: Purdue University: 11-17.
- Zaczek, J. J.; Steiner, K. C.; Bowersox, T. W. 1991. Planting northern red oak: a comparison of stock types. In: Proceedings, 8th central hardwood conference. Gen. Tech. Rep. NE-148. U. S. Department of Agriculture, Forest Service: 604-605.

Oak Plantation Establishment Using Mechanica Burning, and Herbicide Treatmen

James H. Miller, Southern Forest Experiment Station, USDA Forest Ser Auburn University, Auburn, AL 30

ABSTRACT Mechanical methods, prescribed burning, and herbicide treatments for establish oak plantations are reviewed, with emphasis on herbicides. Integrated prescript for site preparation using these silvicultural tools are outlined for both clea forests and old field sites. The basic premise is that intensive cultural treatm will be required for repeated success in establishing oak plantations using ei seeding or planting, when advanced natural regeneration or adequate seed trees not present. Intensive treatments to promote rapid seedling growth appear cri to minimize the intensity and duration of animal predation on oak regenerat Labeled herbicides and appropriate application techniques for both woody herbaceous control are reviewed along with approaches for expanding exis labels and registering new herbicides for oak culture.

INTRODUCTION Much research attests to the fact that oak plantations probably cannot be establic consistently without intensive site preparation, herbaceous weed control, continued release from invading competitors (Malac and Heeren 1979, Ken 1981, Woodrum 1982, Nix 1989). Oak seedlings, like most plants, generally better in a competitor-free space, although not always as rapidly as other species (Sam and others 1986, Bowersox and McCormick 1987, George and c 1991). Why this is so is not completely clear, but it appears partly due t slower early growth habit of many oak species, and their propensity to all photosynthates into root storage and root growth instead of top growth. There been ample research demonstrating accelerated early growth of oaks vegetation control to warrant its use (Aust and others 1985, Krinard and Ke 1987, Wendel and Lamson 1987, Miller and Burkhardt 1987, Johnson and 1989). However, the investment-return outcome for such intensive culture unknown and will remain unknown for some time.

PRINCIPLES OF COMPETITION CONTROL

Over the past decade several principles of vegetation management for pla establishment have been discovered through extensive research, mainly on species. A summary of these principles that should apply to oak culture in

1. Vegetation control can increase tree growth by the greatest proporty years 1-3, especially when near complete control is achieved around large,

seedlings that have been properly planted, and are not injured by the vegetation control treatments.

2. Early growth gains from years 1-3 are maintained and amplified well into the rotation.

3. Threshold levels of competition reduction must be achieved before a positive growth response occurs, and a point of diminishing returns also exists where additional control gains no further growth response.

4. Plants immediately surrounding crop-tree seedlings are the strongest competitors and should be the main target for control.

5. A mixture of grasses, ferns, forbs, vines, and woody resprouts present severe competition to tree seedlings, but grasses and ferns are the most competitive in the early stages from years 1-5.

6. Woody plants become established in greatest numbers during the first year after clearcut harvesting, while some immigration is continuous.

7. The most severe woody competitors are tree species, not shrubs, that can grow equally with crop trees and maintain a position in the main canopy. Resprouting hardwoods are the most severe woody competitors to planted seedlings.

8. Following mechanical site preparation, woody competition starts exerting its influence in years 5-8, but woody control is most cost-effective prior to planting or in years 1-2.

9. Tree diameters respond proportionally more than heights to competition control.

10. Projected returns on investments in intensive site preparation treatments are usually justified only on sites of medium to high quality unless treatments are necessary to guarantee survival because of management objectives.

Besides competition control, it is known that oak seedlings benefit from soil tillage (both surface and subsoil) with assumed benefits in aiding rapid root development (Plass and Green 1963, Woodrum 1982, Seifert and others 1985, Nix 1989). It is also assumed that site preparation requirements will probably be more stringent for oak seeding than for seedling planting (Bullard and others 1992). It is recognized that these principles and concepts will require further verification for specific oak species as they interact with the multitude of sites and plant community situations.

GENERAL PLANTATION ESTABLISHMENT PRESCRIPTION

A proposed general prescription for oak plantation establishment is that large, healthy seedlings should be properly planted on tilled and appropriately subsoiled sites having minimum hardwood competition, with herbaceous vegetation control treatments applied annually for 2 to 3 years (Malac and Heeren 1979, von Althen 1987). Both woody and herbaceous control treatments must be very effective to realize a return on their investment. Because of the constant immigration of hardwoods into young stands (Cain and Yaussy 1984), woody control treatments should be performed at regular intervals as needed. Hannah (1987) summarized: "The key . . . is to control competing vegetation, keeping the oaks dominant and free to grow." Johnson and others (1989) summarized their understanding: "The abundance of red oak in the herbicide-treated clearcut suggests that the key to regenerating red oak may be competition control and not necessarily a long regeneration period."

Rapid early seedling growth is critical to oak establishment to minimize predation from deer, rabbits, and rodents. Only through enhanced early growth can the time spent in vulnerable stages be shortened. (However, rapid early growth begs the issue of eventual log quality as it is affected by a large juvenile wood core (Maeglin 1987).) At the same time, site preparation treatments must work in concert with the overall management plan and safeguard the multi-resource values of the forest, maintain or improve long-term soil productivity, and protect intrinsic site values.

SITE PREPARATION FOR CLEARCUT SITES

Intensive site preparation treatments on cutover forest sites often require the integrated use of mechanical, burning, and herbicide treatments.

Mechanical Treatments

The advantages of using mechanical site preparation treatments before planting o seeding are numerous. Logging slash can be pushed into piles or windrows using rootraking or processed with choppers or flailers. Rootraking is also used to dislodge woody plant roots, minimizing woody competition and increasing availabl nutrients and water for crop tree growth (Burger and Pritchett 1988). The clearing of slash then permits other soil amelioration treatments—disking, bedding, and subsoiling.

Morris and Lowery (1988) reviewed site preparation research in the South an concluded that only disking, bedding, and subsoiling are likely to have larg positive effects on tree growth. On upland and poorly drained sites, disking ha been shown to benefit oak establishment (Woodrum 1982, Seifert and others 1985). Nix 1989). On poorly drained sites, bedding has yielded short-term growt increases that may be maintained (Seifert and others 1985). Disking increases th rootability of the surface soil across the entire area, while bedding does the san and locally raises the soil above the winter water table (Morris and Lowery 1988 However, the volume of tilled soil is less with bedding compared to disking a root closure between rows may not occur as quickly as on disked sites.

The principal disadvantage of rootraking treatments occurs when valuable s nutrients are displaced into windrows and piles, and away from planting rows a seedling access (Morris and others 1983, Tuttle and others 1985). Further, rak or disked organic matter is made to decompose faster, perhaps before seedli roots can take full advantage of the ensuing nutrient release (Banker and oth 1983, Miller and Edwards 1985, Burger and Pritchett 1988). Soil disturbance a clears the organic mulch from the soil surface that aids in preventing evaporat moisture loss. At the same time an abundance of herbaceous plants can beco established on the bared soil and represent severe early competition. herbaceous communities also form excellent rodent habitat that encourages seed seedling predation. On some sites, revegetation after intensive mechan scarification is slowed or spotty leading to sheet and rill erosion and nutrient (Blackburn and Wood 1990). Only through careful and considerate applicatio mechanical treatments can these adverse effects be minimized or eliminated. Subsoiling and ripping are increasingly being used for pine establishment in the South after a history of mixed research results as far as improving both survival and growth (Berry 1979, Lantange and Burger 1983, Wittwer and others 1986). Subsoiling can result in both an increase in the amount of soil exploited by tree roots in the first few years and more uniform depth distribution of roots. Less is known about where and how to apply subsoiling than disking and bedding, but the benefits to deep-rooted species on restrictive soils is obvious. It appears that fracturing of dry pans is preferred to shearing of wet restrictive horizons (Parker and Amos 1982).

There is now a 3-in-1 blade plow manufactured by Symonds, an Australian company, that disks, subsoils (two directions by adding a wing to the ripping blade), and beds in one pass. The 3-in-1 plow has already been used operationally to establish oaks successfully in the South. Another mechanical site preparation tool that could be useful in oak establishment is the Bräcke Scarifer (Alm and others 1988). This scarifier makes discontinuous scarified-mounded planting sites across an area at regularly spaced intervals. About 10 percent of the area is scarified. Oak seeds and seedlings could be planted either in the scarified area or somewhere on the mound—edge to center. Because of the versatility of the hitching arrangement on the scarifier, this implement can be worked in both shelterwood stands and clearcut areas. Seeding and herbicide applications can be simultaneously performed during the operation with added attachments.

Resprouting vigor of hardwoods can be decreased significantly by performing felling or shearing operations after the spring growth flush has depleted root reserves (Zedaker and others 1987, McMinn and Nutter 1988). Performing mechanical treatments during the usually drier summer months should also be more efficient and effective. As the intensity of utilization increases in the future, the need for windrowing or piling treatments will probably decrease, while the need for soil tillage treatments may increase to ameliorate soil compaction caused by intensive harvesting operations.

Prescribed Burning Prescribed burning is commonly used after mechanical and chemical site preparation to further reduce logging slash and improve planter access. Prescribed fire can, with adequate fuel and proper burning conditions (timing), topkill woody plants that are less than 2-3 in. groundline diameter, but can increase the abundance of annual and biennial herbaceous plants (Langdon 1981, Miller 1982, Danielovich and others 1987, Sanders and others 1987, Yeiser 1992). Perennial grass clumps that are also partially consumed regrow with tender shoots, making them more susceptible to herbicide control. It is also known that repeated burns in a stand prior to harvest, spaced over 5 or more years, can reduce the size of woody plants. Fire can also be used after harvest to accomplish the same effect while reducing logging slash. But after harvest, and more so after methods of scarifying site preparation, fuels are discontinuous and ignition may be difficult and more costly with uneven hardwood control.

Soils are not usually degraded by one or more repeated burns, but developing research would suggest that macropores can be filled and porosity lowered with repeated burning of sandy loam soils, resulting in a depression of soil moistureholding capacity (Miller and Boyer 1991). Other research has shown th considerable nitrogen, approximately 60 percent of that in the fuel, can | volatilized from the site with a single burn (DeBell and Ralston 1970, Vose at Swank 1992). Long-term inputs of nitrogen from biological fixation at atmospheric inputs may replace the nitrogen lost from fire, but this is not assure (Boring and others 1991).

Herbicide Treatments Herbicide applications for site preparation before planting oaks can result in the most positive control of woody and herbaceous competition. Modern herbicide are both safe to the applicator and the environment when used according to lab instructions (Miller and Mitchell 1988). They can be applied broadcast, to plantin rows, or selectively to individual target stems (Newton and Knight 1981, Cantre 1985, Miller 1987, Miller and Mitchell 1988). It is necessary that the more ffective herbicide or tank mix be selected for both the target species on the si and also site constraints. The herbicide(s) should be applied at the most effective time using the correct application procedure.

Soil-active herbicides should be used with a full appreciation of the risk (nontarget plant damage due to residual carryover or application proximity. But should be recognized that herbicides having both foliar and soil activity are ofte the most effective. Those herbicides discussed in the following sections an described further in table 1.

		Active	Amount of a.i. or a		
Product	Manufacturer	ingredient(s)	in formulation ¹		
AAtrex 4L	Ciba-Geigy	atrazine	4 lb/gal		
Access ²	DowElanco	picloram + triclopyr	1 + 2 lb/gal		
Accord and Roundup	Monsanto	glyphosate	4 lb/gal		
Arsenal Ac	American Cyanamid	imazapyr	4 lb/gal		
Atrazine 4L	Du Pont	atrazine	4 lb/gal		
Chopper	American Cyanamid	imazapyr	2 lb/gal		
Chopper RTU	American Cyanamid	imazapyr	3.6%		
Escort	Du Pont	metsulfuron	60%		
Fusilade	ICI	fluazifop	1 lb/gal		
Garlon 3A	DowElanco	triclopyr amine	3 lb/gal		
Garlon 4	DowElanco	triclopyr ester	4 lb/gal		
Oust	Du Pont	sulfometuron	75%		
Pathway, Tordon 101R and RTU	DowElanco	2,4-D + picloram	¼ + 1 lb/gal		
Princep 4L ³	Ciba-Geigy	simazine	4 lb/gal		
Tordon K ²	DowElanco	picloram	2 lb/gal		
Tordon 101 ²	DowElanco	2,4-D + picloram	$\frac{1}{2} + 2 \ln/ga$		
Vantage (Poast)	BASF	sethoxydim	1 lb/gal		
Weedone 2,4-DP	Union Carbide	2,4-DP amine	4 lb/gal		
2,4-D ester	several	2,4-D ester	4 lb/gal		

 Table 1—Herbicides labeled or used for oak culture; manufacturers, active ingredients, ar concentrations in formulations

¹ a.i. = active ingredient; a.e. = acid equivalent.

² Restricted use herbicides that must be applied by a State certified applicator or permitted private lander (contact county agent for permit).

³ Other formulations of simazine are Princep Caliber 90, Princep 4G, and Princep 80W.

Herbicide Labels. The herbicide label is a legal document that specifies on what type of sites and how a herbicide can be applied. Herbicides legally used in forestry must be labeled for "forest sites," or in some instances for "noncrop areas" and "tree farms" when not broadcast for site preparation. In oak establishment, the use of herbicides that do not have one of these site specifications is unlawful and carries a prescribed fine and prison term for violations according to State laws. It is also unlawful to exceed labeled rates and use methods of application not outlined on the label.

Other specifications on a herbicide label, such as crop-tree species and target efficacy, are involved with product performance and manufacturer liability. It may be legal to use a herbicide for oak culture when the label states "for conifer release," but the manufacturer is not liable for poor performance or crop injury. Thus, the discussion of which herbicides are labeled for oak culture is not simple and will have to be qualified in the following discussions. Also, the interpretation of labels may vary by State. The direction and possible avenues of future herbicide development and registration for oak culture will be discussed at the end of this paper.

Timing of Herbicide Applications. Herbicides perform best when applied at times the target plants are most susceptible and/or the crop trees are most resistant to injury (Miller and Bishop 1989). Applying them before or after the correct time reduces or even eliminates their effectiveness and may damage the crop. As far as the most efficient timing of woody control treatments, herbicide applications should only be made after all resprouts have emerged following harvesting, burning, or mechanical disturbance. Woody rootstocks must have sprouts before herbicide activity and control can occur.

Woody Plant Control with Tree Injectors and Backpack Sprayers. The manually applied treatments for woody plant control that should have use for oak establishment are:

- tree injection
- stump sprays
- directed foliar sprays
- basal bark sprays

(Soil spot applications using Velpar L Herbicide by Du Pont appear to have limited use in hardwood culture because of the residual nature of Velpar L.) All sizes of trees and shrubs can be controlled by using the right treatment when the proper herbicide is applied at the correct time. Here are the sizes of woody plants that can be treated most effectively by manual application methods:

Method	Effective size of target stems controlled
Injection	sizes greater than 2 in. d.b.h.
Stump sprays	all sizes
Directed foliar sprays	up to 6 ft. tall
Full basal sprays	up to 6 in. d.b.h.
Streamline basal sprays	up to 2 in. d.b.h.

A combination of methods can be used on the same site when an array of targ stem sizes are to be treated. Often, on the same site, tree injection is used for t larger trees, basal or foliar sprays are applied to the smaller woody competitor and stumps of harvested trees are sprayed.

Tree Injection. Tree injection is the least costly of these herbicide treatments 1 controlling unwanted trees that are 2 in. in diameter and larger. This method highly versatile and can be used alone or in combination with other individual ste treatments for site preparation, hardwood release, timber stand improvement, sta conversion, and creating snags for birds and other wildlife. This physical demanding method requires applicators that can repeatedly and precisely chop in tree trunks deep enough to properly deliver herbicide for uptake in the sap flor Each cut must form a pocket into the sapwood, where the herbicide is placed f uptake. The herbicide should stay in the pocket and not seep out through any sp sides, because any herbicide on the bark is wasted. When treating sproutin clumps, each stem must be injected. The variable results that can occur with th method are partially caused by the inability of applicators to penetrate the ba sufficiently or to correctly place the herbicide for uptake.

Common methods of tree injection are:

- tubular tree injectors
- hypo-hatchets
- hack-and-squirt

Tubular tree injectors consist of a long metal tube fitted with a chisel-type blac that is used to cut through the tree bark into the sapwood near the base of the tre Several models are available. Units are equipped with a lever, handle, or wire th is pulled to deliver the herbicide (usually 1 mL) from the cylinder into the cu The delivery rate can be adjusted for accurate calibration. To calibrate: fill ar prime the injector; pull the handle or wire 10 times while collecting the herbicid in a container graduated in milliliters; if this is not 10 times the desired rate, adju the lock-nut and repeat the procedure until accurate calibration is achieve Frequent sharpening and maintenance of injection tools is needed for best result

The hypo-hatchet consists of a hatchet with an internal herbicide delivery syste that is connected by a hose to a herbicide container carried on the back or be When the hatchet strikes a tree, the blade must penetrate into the sapwood and impact drives a piston forward delivering 1 mL of herbicide into the cut. The r cannot be adjusted. Daily cleaning and lubrication of the impact piston is require maintenance, along with periodic replacement of rubber O-rings and seals. Saf glasses should always be worn when using a hypo-hatchet because of freque herbicide splashes.

Hack-and-squirt is a method that uses a narrow-bit ax, hatchet, or machete making the cut, along with a spray or squeeze bottle or oiler to deliver herbicide. A grinder can be used to narrow the bit of axes and hatchets for ea and better cuts. Most commercial spray bottles are set to deliver 1 mL with e trigger pull, but each must be checked prior to use. Safety glasses also shoul worn when using this method. Waist-high injections by the hypo-hatchet and t and-squirt methods are just as effective and fast to perform as basal inject With larger stems, more herbicide is applied by basal injections because of the larger groundline diameter compared to diameter at breast height.

The amount of herbicide per injection and the edge-to-edge spacing are specified on the herbicide label. Continuous edge-to-edge cuts should be used on hard-tocontrol species such as dogwood, maple, and hickory. Herbicides labeled for tree injection that have wide control spectrums are:

- Arsenal Ac
- Accord and Roundup
- Chopper
- Garlon 3A
- Pathway, Tordon RTU (Ready To Use), Tordon 101R, and Tordon 101
- 2,4-D

Some of these herbicides have the same active ingredients with new names or a slight difference in formulation (table 1). Accord is the same as Roundup without a surfactant, Chopper is half the concentration of Arsenal Ac, and Pathway is the new name for Tordon RTU and Tordon 101R. Accord will replace Roundup, and Pathway will replace Tordon formulations as the other names are phased out. Efficacy of these herbicides for selected species is presented in table 2.

Herbicide	Suscepti	ible	Moderate	Toleran
Arsenal Ac/Chopper	Sweetgum	Hickory		Pine
	Southern red oak	Dogwood		Elm
	Northern red oak	Ash		
	White oak	Beech		
	Post oak	Sourwood		
	Water oak	Blackgum		
	Chestnut oak	Red maple		
	Black cherry			
Accord/Roundup	Sweetgum		White oak	Ash
•	Southern red oak		Northern red oak	Hickory
	Post oak		Water oak	
	Blackgum		Red maple	
	Sourwood		Black cherry	
			Dogwood	
			Pine	
			Elm	
			Chestnut oak	
			Beech	
Garlon 3A	Sweetgum	Dogwood	Blackgum	
	v	Pine	Water oak	
	Northern red oak	Elm	Red maple	
	White oak	Chestnut oak	Black cherry	
	Post oak	Sourwood	Ash	
	Hickory		Beech	

Table 2-Species susceptibility to injection herbicides under ideal conditions and timing

Herbicide	Suscepti	ble	Moderate	Tolerant
Pathway/Tordon	Sweetgum Southern red oak	Beech Hickory	Red maple	
	Northern red oak			
	White oak	Dogwood		
	Post oak	Pine		
	Water oak	Elm		
	Chestnut oak	Ash		
	Black cherry	Blackgum		
2,4-D	Southern red oak		Sweetgum	Water oal
	White oak		Northern red oak	Red mapl
	Post oak		Black cherry	Ash
	Blackgum		Hickory	Chestnut o
	Dogwood		Pine	Beech
	Elm		Sourwood	

Table 2—Species susceptibility to injection herbicides under ideal conditions a timing—(Continued)

Garlon 3A and Accord (Roundup) have the advantage of no soil activity. Of the two, Garlon 3A is preferred because it is effective on more species, especial maple and hickory. Arsenal Ac, although soil active, has the broadest spectru of control of any of these herbicides and can be used at wider-spaced injection cut Garlon 3A and Arsenal Ac are usually applied diluted at 33-50 percent and 5-1 percent, respectively. All products can be applied year-round, except during time of heavy sap flow in the spring. Arsenal and Chopper are most effective whe injected from July to October.

Stump Spraying. Stump resprouting of many species can be prevented of decreased by a low-cost herbicide treatment following harvest and after partial cu for timber stand improvement. Stumps larger than 12 in. diameter do not usual resprout and do not need treating. Of course, other hardwoods may invade an ar with time. Hand clearing treatments for release or thinning can be enhanced I treating the stumps with herbicide. Stump spraying after bush-hogging is anoth treatment alternative. The same herbicides that are used for injection are label for stump treatments; also labeled are Chopper and Chopper RTU that have t same active ingredient as Arsenal Ac.

A backpack sprayer can be used that has a wand or spray gun equipped with straight stream, fan, or hollow-cone nozzle. A sawyer can carry herbicide in utility spray bottle for treating stumps after cutting. For small-diameter stump a wick applicator can be used.

Freshly cut stumps should be treated as soon as possible. Stump treatments wit 4 hours of cutting have been most effective—the sooner the better. For stun over 3 in. in diameter, the outer edge or cambial area must be completely wet with the herbicide. Smaller stumps are usually completely wetted. To successful, all small stumps should be treated. Thus it is best that the sawye companion applicator treats soon after felling so no stems are skipped. Cutting herbicide treatments can be performed during late winter and summer. Winter treatments are slightly less effective than growing-season treatments, but only a 60-80 percent control success should ever be expected with stump spraying (Zedaker and others 1987). Older cut stumps can be treated with the streamline basal stem mixture (see that section). The mixture is applied to the outer 1 in. edge of the stump until runoff and to the base of any sprouts.

Directed Foliar Sprays. Directed foliar sprays are more cost-effective than basal sprays for controlling woody competition that is less than 6 ft. tall (Thomas and others 1989). Directed foliar sprays are usually applied with a backpack sprayer fitted with a spray wand equipped with a full cone, flat fan, or adjustable cone spray tip. Spray guns attached to the backpack unit with narrow flat-fan tips are also used by some applicators. Backpack mistblowers are another means of applying foliar sprays (Garrett and others 1989). The spray is applied to the target foliage, being sure to cover the growing tips and wetting the leaf surfaces without drip.

Herbicides labeled for directed foliar sprays for site preparation, that are not restricted to "conifer" reforestation, are:

- Accord and Roundup
- Garlon 3A and Garlon 4
- Tordon 101 and Tordon K
- 2,4-D and Weedone 2,4-DP

Tank mixes of these products will usually be more effective when treating mixed species (Johnson 1987; Shiver and others 1990, 1991). The comparative efficacy of most of these products when applied singly as foliar sprays at various timings have been reported, except the Tordons and 2,4-D (Miller 1990b). In general, the most effective timing for most species was found to be from mid- to late-summer. Arsenal Ac was also tested and found to have the most broad-spectrum control, but it is only labeled for conifer site preparation. Care should be exercised that at least 6 months lapses between application and planting when using Tordon.

Basal Sprays. Basal sprays are more costly than directed foliar sprays for controlling the same sized woody plants (Thomas and others 1989), but basal sprays can control trees larger than 6 ft. tall and can be applied in the late dormant season resulting in less unsightly brownout than foliar sprays. Labeled herbicides that have broad-spectrum control are Garlon 4 with no soil activity, and Access, Chopper, and Chopper RTU with soil activity. Chopper RTU is applied undiluted while the other products are mixed with oil and/or a penetrant for bark applications.

Full Basal. Full basal treatments require that the lower 12 to 20 in. of target hardwood stems be completely wetted with the spray mixture on all sides. A backpack sprayer is used with a wand or spray gun fitted with a narrow angle flat fan, cone, or adjustable tip. Herbicides are used that are soluble in oil and mixed at percentages specified on labeled products, usually less than 10 percent.

Streamline Basal. Streamline basal treatments can control many woody plants including hardwoods up to 2 in. d.b.h. (Miller 1990a). Trees of susceptible species

up to 6 in. in diameter can be controlled. Treatment of small hardwoods that a less than 2 in. d.b.h. results in the most control.

To apply this treatment, a backpack sprayer is used with a spray gun and a lo flow (0.1-0.2 gallon per minute (gal/min)) straight-stream spray tip. Also, narrow-angled tip can be used, such as 15° and 0.1 gal/min. For controlli herbicide output to prevent waste, a pressure regulator is needed to mainta pressure below 30 lb per square inch (lb/in.²). At these pressures, an effecti reach of 9 ft. is possible while bark splash is minimized. Sprayers with diaphrag pumps will maintain about 30 lb/in.² with slow, steady pumping.

The most commonly used mixture for streamline application includes Garlon 4 20 percent, a penetrant at 10 percent, and a carrier such as diesel fuel or miner oil. This mixture is clear when made correctly, while a white cloudy liquid or g will form if even a small amount of water is present. No amount of water shou be in the sprayer or mixing container. Make sure that all water has been draine from the sprayer, the pump has been pumped dry, and the sprayer has been rinse and pumped with mineral oil or diesel before filling with the mixture.

For treating stems that are less than 2 in. d.b.h., apply the stream of spray up-and down single stems for about 6 to 8 in. or as a 2- to 3-in.-wide band across multip stems. Direct the spray stream at a point about 6 to 24 in. from the ground 1 smooth juvenile bark. Stems that are beyond the juvenile stage, thick barked, c near 3 in. in diameter require treatment on both sides, unless they are susceptibl species. Back and forth bands can also be sprayed on larger stems.

Applications are usually made in late winter and early spring when leaves do no hinder spraying the stem and the effectiveness for many species is maximur (Pancake and Miller 1990). The best application time will depend on the herbicide species, and location. Avoid applications on hot days if an ester formulation, suc as Garlon 4, is used because crop injury may occur from vapor drift.

Broadcast Applications by Helicopters and Ground Sprayers. Herbicide labeled for broadcast applications prior to planting hardwoods are:

- Accord and Roundup
- Garlon 3A and Garlon 4
- Tordon 101 and Tordon K
- Oust (herbaceous plant control)

The best control will be obtained by using mixtures of these products if there is mixture of target species on a site (Johnson 1987; Shiver and others 1990, 199 Seifert 1990a). Oust can also be mixed with these herbicides without decreasi their effectiveness while increasing herbaceous control (Jones and others 1986 However, the most effective mixture for controlling a specific species mix is s up for conjecture, requiring the user to consult local extension specialists a knowledgeable managers. When competition is essentially only one species, the one herbicide may be best. For controlling sugar maple (*Acer saccharum* Marsh Garrett and others (1989) reported that Tordon 101 as a 20-percent solution most effective of the 11 herbicides tested using a mistblower.

These herbicides can be applied by both aerial and ground sprayers. Aerial broadcast applications are commonly used on tracts of 50 acres or more, because of improved coverage and costs compared to ground applications. Professional aerial applicators generally use helicopters with spray systems equipped with microfoil booms, thru-valve booms, raindrop nozzles, or other devices to assure accurate applications with a minimum risk of off-site drift to neighboring lands. Managers should verify the accuracy of calibration before application. Herbicide mixing procedures, especially in batch trucks, should be examined and even tested for thoroughness, which can be performed by using electrical conductivity with Accord mixtures (Lautenschlager and Schaertl 1991). Pumping the mixture three times through the system can also assure adequate mixing. The land manager or owner also has responsibilities for preparing the site for aerial treatment, such as felling tall snags, heliport construction, and marking boundary lines that are visible from the air.

Broadcast site preparation treatments are also conducted with various types of tractor sprayers. Boomless cluster nozzles, manifold nozzles, and mistblowers mounted on either rubber-tired tractors and skidders or track-type tractors are used on sites with flat or gently rolling terrain. Applicators can now use computerized sprayer control systems on their tractors that automatically maintain the proper rate on terrain where ground speeds vary, but these require calibration that should be verified by the manager. For treatment of small regeneration areas, hose-reel sprayers mounted on trucks or tractors are also used.

Broadcast herbicide applications are often followed in 8-12 weeks with a prescribed burn to reduce standing and down woody material for better planter access. Burning does not always increase hardwood control (Minogue and Lauer 1992) and can volatilize site nitrogen, as previously discussed. Chopping is also being increasingly used after herbicide spraying in pine culture, with and without burning, to improve access and the ease of subsequent operations.

Control of Problem Plants During Site Preparation. Some of the problem weeds that can hinder or prevent plantation establishment are hayscented fern (*Dennstaedtia punctilobula* L.), New York fern (*Thelypteris noveboracensis* (L.) Nieuwland), japanese honeysuckle (*Lonicera japonica* Thunb.), kudzu (*Pueraria lobata* (Willd.) Ohwi), trumpetcreeper (*Campsis radicans* (L.) Seemann), grape (*Vitis* spp.), multiflora rose (*Rosa multiflora* Thunb.), privet (*Ligustrum* spp.), and eastern redcedar (*Juniperus virginiana* L.). A summary of herbicides for treatment includes:

Hayscented and New York ferns. Oust alone (1-2 oz/acre) or in a tank-mix with Roundup (1 qt/acre) applied between early July and early October (Horsley 1988, 1990). The timing of Oust for site preparation can influence the damage on other desirable hardwoods (Horsley and others 1992). McCormick and others (1991) in Pennsylvania on poorly drained soils found that fall applications of Oust at 2 and 4 oz/acre alone or in combination with Roundup at 1 qt/acre had no effect on germination of northern red oak (*Quercus rubra* L.), but did increase first-year mortality by 14-16 percent and reduced second-year height growth by 14-23 percent. Japanese honeysuckle. Escort at 1 oz/acre applied May to August or Roundup a 0.75-percent solution applied September to October (Edwards and Gonzalez 1986) Regeher and Frey 1988, Schmeckpeper and others 1987).

Kudzu. Tordon 101 at 1-2 gal/acre or Tordon K at 0.5-1 gal/acre applied June t September both in year 1 and repeated with half the rate in year 3, with spot clean up as needed and a 6-month wait before planting after last application (Mille 1988). Eradication on the site is needed before planting.

Trumpetcreeper. Roundup at 4 qt/acre applied July to September (Pyle an Krueger 1984).

Grape. Smith (1974) summarized screening trials and recommended basal bar sprays (lower 12-18 in.) in oil mixtures using 2,4-D ester (many brand names) and Weedone 2,4-DP at labeled rates and cut stump treatments using water mixtures o Tordon 101 (50-percent solution) and Roundup (20-percent solution) and undilute Pathway (Tordon RTU, 101R), all applied early March or mid-September.

Multiflora rose. Roundup in a 1-percent solution or Garlon 4 in a 0.5-percen solution using summer or winter applications (Bhowmik and Germond 1987).

Privet. Arsenal AC in a 1-percent solution and surfactant sprayed in mid-summer

Eastern redcedar. This species is not a basal sprouter, so cutting near groundline is effective. For large trees, foliar spray with Tordon K at 0.25-percent solution in a 1-percent diesel oil-water emulsion, wait 3-4 weeks, and ignite crown (Stritzke and others 1991).

These species must be nearly completely controlled or eradicated from a plantation site or the remaining plants will spread quickly.

SITE PREPARATION ON OLD FIELD SITES

The use of disking and subsoiling should be considered for improving so conditions and competition conditions before planting oaks on abandoned fields an pastures (Malac and Heeren 1979). Disking treatments will improve plantin operations if performed correctly and often promote annual herbaceous plants thare more effectively controlled with herbicides than are perennial plants. Diskir should be to a depth of 8 in. and should be done in strips along the contour reduce the chance of soil erosion. Unfortunately, disking can aggravate we weather planting the following spring. Subsoiling or ripping can be used to breaup plowpans that are common to these sites.

Part of the decision to use tillage treatments must consider whether the site designated as wetlands and whether the tillage practice would be considered "s busting" that might jeopardize participation in USDA-sponsored farm program

For controlling pasture grasses and/or forbs before planting, late-sumr applications of Roundup can be applied broadcast or in bands to form plan rows. Only Roundup appears specifically labeled for this situation. Rates of 3-5 qt/acre will be required to control established sod, and even then complete control cannot be expected. A prescribed burn in early summer, before the Roundup applications, can clear standing dead grass parts to improve herbicide efficiency. Also, mowing can be used to improve access and allows better spray coverage by reducing vegetation to a more uniform height. If mowing is done before a herbicide treatment, wait for 4-6 in. of weed regrowth before Roundup applications.

ERBACEOUS WEED ONTROL IN YOUNG OAK PLANTATIONS

Erdmann (1967) noted the benefits of herbaceous weed control on red oak establishment in old fields using Princep 4L at 1 gal/acre applied at the time of planting. Disking prior to spraying was found to enhance control. Erdmann, in the same study, and others (von Althen 1972, Wendel 1980, Nix 1989) have reported that black plastic and cardboard mulches are ineffective for herbaceous weed control with oak establishment and can harbor rodents. Erdmann reported that plowing and disking failed to stimulate rapid height growth of oak seedlings, but mechanical site preparation was a prerequisite to satisfactory weed control when using simazine (Princep). For sandy soils, he recommended applying 2-2.5 lb active ingredient/acre, although this use is no longer labeled.

Numerous screening studies have identified Oust, Princep, and Roundup (Accord) to be effective for herbaceous weed control when planting oaks, with the addition of atrazine having inconsistent results (Kosinski and Holt 1985; Wright and Holt 1985; Seifert and Holt 1985; Jones and others 1986; Seifert 1989a, 1989b, 1990b). Crop-tree injury usually increased with increased rates. Several of these studies also showed that Arsenal, as yet not labeled for hardwoods, held promise for herbaceous weed control with minimal oak injury. Seifert (1989b) after testing 11 herbicides/combinations applied for 2 consecutive years in southern Indiana, found that Oust provided the best weed control and least damage to northern red, white (Q. alba L.), bur (Q. macrocarpa Michx.), and black oak (Q. velutina Lam.) when applied at 1, 2, 4, and 8 oz/acre. Subsequent tests found that 4 and 8 oz/acre were too high and could result in stunting and mortality (personal communications with John Seifert). Wright (1986) studied pre-plant spring herbicide applications for site preparation for red and white oak plantings on old fields with forbs, grasses, and semiwoodies and found combinations of Roundup and Princep to be costly but effective while Oust was partially effective but released broomsedge. Unfortunately, resistant species (like broomsedge) or mid-summer annuals often recapture the treated area when effective herbaceous control is achieved, requiring the use of Roundup spot treatments with shielded spraying.

Two greenhouse studies have indicated that Oust inhibits the emergence and principally root growth of northern red oak (Sam and others 1986, Shipman and Prunty 1988), but the height growth of surviving seedlings was significantly increased. Similarly, Barnes and others (1990) have shown that Oust reduces root growth potential of loblolly pine (*Pinus taeda* L.), but it is still widely used for pine release because of eventual growth stimulation through competition control.

Sam and others (1986) found that of the preemergent herbicides tested, only Ou performed equally well on bare soil or forest litter. Therefore, applications to ba soil are not needed with Oust as has been required with Princep applications.

Two herbicides commonly used for establishing hardwoods have questionab registration at this time. In 1987, the use of Princep (simazine) for establishin forest plantations was removed from the label, although the product is st registered for Christmas trees, nurseries, and shelterbelts. Also, the use of atrazin is limited on labels to conifer establishment, which appears to make use hardwood establishment the responsibility of the user with no recourse for produ performance or crop damage. This interpretation may vary by State.

The use of repeated disking or mowing as a substitute for herbicide applications f herbaceous weed control has not been fully tested. Zutter and others (198 reported that repeated herbicide applications were more effective than repeated cultivation for sweetgum (Liquidambar styraciflua L.) and green ash (Fraxin pennsylvanica Michx.), because the inter-row vegetation was not controlled 1 cultivation. Malac and Heeren (1979) presented plantation establishment guidelin for hardwoods that stressed at least 2 years of frequent disking for competitic control. The cultivator disks were designed to cast a mound of soil against the ro of trees to smother the weeds within the row. Kennedy (1981) reported that years of repeated cross disking on a clay soil with severe herbaceous competitic enhanced diameter growth of Nuttall oak (O. nuttallii Palmer) 240 percent ov checks, while repeated mowing was not different from the check. When comparin cultivation with herbicide control, it is recognized that surface evaporation fro cultivated soil is greater than from soil covered with herbicide-controlled vegetatic acting as a mulch. However, preemergent applications of herbicides, because their timing, may not result in a mulch of dead vegetation either. Also, the ba ground caused by either cultivation or herbicides will soon be revegetated to son degree.

Herbicides for Herbaceous Weed Control

Oust. At present, Oust is labeled for herbaceous weed control at 1-2 oz/acre aft transplanting (not seeding), specifically for northern red oak, white oak, a chestnut oak (Q. prinus L.). The lower rate is used on coarse-textured soil. T efficacy of Oust on certain species is presented in table 3.

1. Oust is best applied as a preemergent herbicide (before weeds emerge).

2. Oust is not recommended for use on poorly drained or marshy sites, bu may be used where hardwoods have been planted in beds.

3. Application should be made at the time of tree planting or within 2 weeks tree planting, in the spring before planted seedlings leaf out.

4. Seedling injury can occur if the planting slit is not fully closed and if I break has occurred. Injury also can occur if heavy rainfall occurs after applicat and root growth has started.

Simultaneous seedling planting and spraying can be used with a simple sp attachment to the planting machine. Three years of herbaceous weed control h been recommended for hardwood establishment on old fields in Indiana by Wr and Holt (1985). Even though mixtures of atrazine and simazine have been in for many years, they are no longer labeled specifically for hardwood establishment

Susceptible	Moderate	Tolerant
Ragweed	Panic grasses	Broomsedge
Fescue	Goldenrod	Bermudagrass
Horseweed	Dogfennel	Nutsedge
Ferns ¹	Bahiagrass	Morningglory
Burnweed	Johnsongrass ¹	Woolly croton
Boneset	Pokeweed	Tropic croton
Sunflower		Trumpetcreeper
Poorjoe		Sicklepod
Dewberry ¹		Cocklebur
Vetch		Lespedeza
Geranium		Wiregrass
Golden weed		Plumegrass
Sweet clover		Ū.
Crabgrass		
Brome		

 Table 3—Weed species usually susceptible to preemergent application of

 Oust at 2 oz active ingredient/acid equivalent

¹ Controlled only partly in preemergent or early postemergent applications to seedling plants, not established perennials.

Fusilade and Vantage (formerly Poast). Two herbicides that can be used for controlling only grasses are Fusilade and Vantage. (Vantage is a ready-to-use, weaker formulation of the herbicide Poast.) They are expensive products to use and both require up to two applications for controlling perennial grasses. Neither control broadleaf forbs or nutsedge. Fusilade is generally considered safe for over-the-top applications on all hardwood and conifer seedlings, while the phytotoxic effects of Vantage are assumed to be the same. Grasses must be small and tender for over-the-top applications. Fusilade is applied at 32-48 oz/acre with a nonionic surfactant and Vantage is applied at 35-61 oz/acre. Both can be applied only by ground application, using label-specified equipment.

Application Methods for Herbaceous Control Herbaceous Control Herbaceous Control Herbaceous Herbaceous weed control can be accomplished using broadcast, band, or planting spot applications. Band applications are possible only when planting rows are well defined and wind conditions permit spraying well-defined bands. Banded or spot applications cost considerably less than broadcast treatments since only a part of the area is treated, while only small losses in pine growth and survival occur when bands or spots exceed 3 ft. in width or diameter relative to broadcast applications (Dougherty and Lowery 1991, Yeiser 1992). However, growth response increases in proportion to the area treated and reinvasion may be quicker with spots and bands (Dougherty and Lowery 1991). Initial growth retardation in the first year due to toxicity by some herbicides is overcome, resulting in a positive growth response by age 2-3 for both oaks and pines (Wright and Holt 1985, Yeiser 1992).

> Broadcast applications can be made by backpack sprayers, tractor sprayers, or helicopters (Oust only). Banded treatments are applied using backpack, all-terrain vehicle (ATV), or tractor sprayers. Backpack sprayers treat one row at a time

while machine sprayers can treat two or more rows using a boom with space nozzles. Bands usually range from 3 to 6 ft. wide. Precise mixing and application are essential for successful treatments.

To apply banded treatments, the boom or spray wand is fitted with a wide-angle flat-fan tip, such as 80° or 110°, with flow rates of about 0.2 to 0.3 gal/min o two narrow-angle tips per row. When using a single wide-angle tip per row, a special flat fan tip that is "even" will increase control on the edges, compared to a regular flat fan that applies less herbicide on the edges. Also, flood tips are commonly used for banded applications, because one tip can apply a 3 to 5 ft band. Wide angle or multiple tips are used to minimize the wind influence by having the tip close to the ground. Tip arrangements should be selected and arranged to assure even distribution across the band while minimizing rates around the planted seeds or seedlings. Thus, two tips can be positioned on either side of the seedling where the overlap is minimal to lessen the rate applied directly to the seedling.

Pressure regulation and a constant nozzle height and ground speed are needed to maintain uniform application rates. Use a pressure of 5 to 15 lb/in.² to give large droplets with reduced drift. The low flow rates from low pressure operation also permit more acreage to be covered per fill-up, adding to the efficiency of the operation. Special "extended range" or "low pressure" tips are designed especially for low-pressure applications. A pressure regulator is necessary to maintain low pressures with backpack sprayers as well. Some backpack sprayers can be set at the desired pressure while others rely on the installation of regulators in line or on the wand.

Many types of sprayers can be modified to apply banded herbaceous weed control treatments in plantations. One increasingly common machine is a four-wheel, ATV equipped with a sprayer. ATV and tractor sprayers can be equipped with sprayer control systems, with ground speed sensing. Sprayer control systems can maintain a constant application rate over a wide range of operating speeds (Miller and Mitchell 1988).

Spot applications must be used when planting rows are not well defined. Spot applications can be made with the above banding procedure using a backpack sprayer with some change. Mainly, the sprayer is turned off between seedlings as the applicator follows the rows. The sprayer is turned on about 1.5 to 2.5 ft before the seedling and shut off at the same distance past the seedling. Thus the same tips, pressure, and calibration can be used. The applicator must still maintair a constant walking speed while spraying each seedling if the rate is to be constant

Full cone tips, with flow rates from 0.2 to 0.4 gal/min, can be used to apply circular, tree-centered spots using a backpack sprayer or spotgun. However, ful cone tips with these low flow rates produce very fine droplets that are easily blown by wind. An adjustable cone nozzle produces large droplets, but these droplets ar too large to ensure uniform coverage of preemergent herbicides. Thus, it i difficult to achieve good results with full cone tips.

RELEASE AND THINNING Inevitably release treatments will be required in oak plantations as partially controlled woody plants and new immigrants appear. Although research has begun to find effective over-the-top selective release treatments, it may be some time before successful treatments are identified and developed (Pham 1987).

Mixed results for oak release have been reported, but the variable results are probably due to ineffective control and the initial vigor of the released oaks. Wendel and Lamson (1987) concluded that rapid resprouting of cut-only release treatments nullify any benefit. Russell (1974) studied planted northern red oaks treated at 6-7 years in the Cumberland Plateau and found that woody competition clearing, once or continuously for 3 years, showed improved diameter growth, but not height growth after 3 years. Nix and Cox (1987) studied cherrybark oak (Q. falcata var. pagodifolia Ell.) plantings (2-0) and found that seedlings released in 3-4 ft. radius spots using Roundup directed sprays late in the second year did not grow better than unreleased trees. In a subsequent study, Nix (1989) reported significant release response to the same treatment on the same crop species, but released in the first year. Woodrum (1982) studied cherrybark oak in South Carolina and found that glyphosate release spraying caused decreased height growth (suspected drift) but increased diameter growth in the growing season of treatment.

As summarized in the introduction, diameters are influenced more than heights by competition reduction treatments, as also reported in these release studies. Also, release treatments must be applied at a very early stand age to ensure a response. Release should be applied in year 2 and repeated as needed to control invading arborescent hardwoods. Often, past the second year, the costs of control increase geometrically. Release treatments that treat greater than a 3 to 5 ft. diameter spot around the seedling should also be more effective but obviously will cost more. Sprouting clumps and tree species with rapid juvenile growth should be the primary control targets.

Roundup is the only herbicide that is currently labeled for "postdirected sprays" on "silvicultural sites." A 2-percent solution should be sprayed to cover the foliage of target woody competitors while preventing spray solution on crop foliage. To minimize crop seedling damage, a shield cover can be placed over the nozzle to help prevent drift. Also, applications should only be made during times of low wind. Drift control agents can be added to the spray mixture, but their addition can possibly lessen control effectiveness. Herbicide damage to crop trees can nullify any release response.

Release by directed sprays will only be feasible when target plants are less than 6 ft. tall and are species susceptible to Roundup. Garlon 3A, Garlon 4, and Arsenal AC could legally be used for directed sprays, but since the label specifies such treatments only for conifer release, the user would accept all liabilities of oak seedling damage.

Basal sprays with Garlon 4 in oil carriers can be used for release treatments, but crop-tree damage can occur from volatility when treating on warm days. Thus, winter applications in January and February are preferred. For larger competitors, cutting and stump treatments with Roundup and Garlon 3A are labeled uses, but crop-tree mortality may occur due to root grafts and root exudation and uptake. Thinning can be important in the production of quality hardwood logs, and thus herbicides may play a critical role for precommercial thinnings and preventing resprouting of commercially thinned oaks. Both directed foliar and basal sprays could be used for precommercial thinnings. The systemic activity of herbicides will have to be tested to identify herbicides suitable for thinning sprouting clumps so as not to injure the selected crop sprout.

FUTURE REGISTRATION OF HERBICIDES FOR HARDWOOD MANAGEMENT

All herbicides used in the United States must be registered by the Environmental Protection Agency (EPA) as specified in the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and its amendments. To develop a new herbicide requires years of extensive scientific testing on the product's chemical properties, efficacy, and mammalian and environmental toxicology, after formulation development has been achieved. Costs of developing, testing, and registering a new herbicide presently exceed \$30 million.

A unique requirement for pesticides registered for "forest use" is the environmental fate study, whereby the active ingredient and byproducts (metabolites) are tracked through a watershed application until dissipation below detection limits is documented in plants, soil, and water. This study alone now costs about \$500,000 for a single test. Herbicide companies must then plan to recapture this investment plus profit, before embarking on a full "forest use" registration. Obviously, for products with good conifer tolerance and broad-spectrum control, this investment is justified, and over the past 15 years there have been numerous new herbicides registered accordingly. Unfortunately most of these herbicides are for controlling hardwoods, not establishing them.

Some herbicides, such as Fusilade and Vantage, are being used in forestry, especially for hardwood culture, under the categories of "noncrop areas" and "tree farm" without having an environmental fate study. It has been verbally conveyed by EPA that this procedure will be permitted as long as the herbicide is not used for broadcast applications in site preparation. Perhaps other herbicides presently registered for other crops can gain use in hardwood culture in a similar manner.

The registration process offers three other avenues for gaining labeled herbicides required for hardwood culture. First, herbicide companies could be requested to extend their existing forest use registrations of "site preparation herbicides for conifer culture" to include uses in hardwood culture. Tests of the residual activity of site preparation herbicides, like Arsenal Ac, would have to be performed and appropriate waiting periods between application and planting or seeding be specified on the label, to manage the liability risk. Second, shielded applications could be specified for nonselective herbicides for use as hardwood release treatments. This is underway with the Accord label, with promises to have shielded applications specified on the next label edition.

Perhaps the quickest way to gain registration is though the provisions in FIFRA for Special Local Need registration. Under Section 24(c) of FIFRA, a State may register any federally registered pesticide to satisfy special local needs, provide that (1) registration for such use has not previously been denied or canceled b EPA, and (2) a food tolerance (safe levels in food), if required, has been established for the proposed use. In forestry the food tolerance would not be required. This "24(c)" registration process has been used for forestry herbicides, the most notable being for Oust in Indiana that led to a Federal supplemental label.

Going further back in the development process, some herbicide manufacturers have begun greenhouse screening on woody plants soon after newly synthesized molecules show promise on major crops and weeds. In the recent past, products suitable for forestry were only identified by happenstance during field screenings performed by company researchers in the "specialty products" areas—forestry being one of these. Still, it is most often the responsibility of "minor use" industries and government agency researchers to identify and develop the use of herbicides from those labeled for general agriculture or rights-of-way. Perhaps herbicide manufacturers can in the future be encouraged to identify those test products that demonstrate selectivity among hardwood species during these early screenings. This will probably be the only way that broadcast oak release herbicides can be developed. Of course, specific herbicides will only be developed when hardwood plantings are being made on a sizable enough acreage to justify this intensity of initial screening.

PUTTING IT ALL TOGETHER Only with effective cultural tools can intensive oak culture become an affordable reality. Much greater research effort is required to develop these tools. Another effort will be required in sharing what works and what does not work—thus the importance of these Proceedings. The tools must be reasonably priced, which means that the market must be large enough for economy of scale in manufacturing, registering, and marketing. This will take time in the early stages as market size grows with continued success as well as good communication of cultural needs to entrepreneurs interested in this market. The implementation of new tools will require an added degree of technical sophistication by managers to be able to use new methods properly, such as modern herbicides. Integrated establishment systems, combining these tools, will need development and testing. The demand for quality oak wood will drive this development, but timely contributions from research, extension, and manufacturing will all be required.

LITERATURE CITED

- Alm, Alvin A.; Long, Steven E.; Eggen, David L. 1988. Integration of forest site preparation with the Bräcke Scarifier. Northern Journal of Applied Forestry. 5:12-13.
 - Aust, W. Michael; Hodges, John D.; Johnson, Robert L. 1985. The origin, growth and development of natural, pure, even-aged stands of bottomland oak. In: Proceedings of the 3rd biennial southern silvicultural research conference; 1984 November 7-8; Atlanta, GA. Gen. Tech. Rep. SO-54. New Orleans, LA: U.S. Department of Agriculture, Forest Service: 163-170.
 - Banker, R.E.; Miller, J.H.; Davis, D.E. 1983. First-year effects of rootraking on available nutrients in Piedmont Plateau soils. In: Proceedings of the 2nd biennial southern silvicultural research conference; 1982 November 4-5; Atlanta, GA. Gen. Tech. Rep. SE-24. Asheville, NC: U.S. Department of Agriculture, Forest Service: 23-25.
 - Barnes, Andrew D.; Zedaker, Shepard M.; Feret, Peter P.; Seiler, John R. 1990. The effects of sulfometuron on the root growth of loblolly pine. New Forests. 3:289-295.

- Berry, Charles R. 1979. Subsoiling improves growth of pine on a Georgia Piedmont site. Res. Note SE-284. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 3 pp.
- Bhowmik, Prasanta C.; Germond, Bart J., Sr. 1987. Multiflora rose control in 1983, 1984 and 1985 [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1986 December 2-4; Milwaukee, WI. Champaign, IL: North Central Weed Science Society: 41:241.
- Blackburn, W.H.; Wood, J.C. 1990. Nutrient export in stormflow following forest harvesting and site-preparation in East Texas. Journal of Environmental Quality. 19:402-408.
- Boring, Lindsay R.; Hendricks, Joseph J.; Edwards, M. Boyd. 1991. Loss, retention, and replacement of nitrogen associated with site preparation burning in southern pine-hardwood forests. In: Nodvin, Stephen C.; Waldrop, Thomas A., eds. Fire and the environment: ecological and cultural perspectives: Proceedings of an international symposium; 1990 March 20-24; Knoxville, TN. Gen. Tech. Rep. SE-69. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 145-153.
- Bowersox, T.W.; McCormick, L.H. 1987. Herbaceous communities reduce the juvenile growth of northern red oak, white ash, yellow poplar, but not white pine. In: Hay, R.L.; Woods, F.W.; DeSelm, H., eds. Proceedings of the central hardwood conference VI; 1987 February 24-26; Knoxville, TN: 39-43.
- Bullard, Steven; Hodges, John D.; Johnson, Robert L.; Straka, Thomas J. 1992. Economics of direct seeding and planting for establishing oak stands on old-field sites in the South. Southern Journal of Applied Forestry. 16:34-40.
- Burger, J.A.; Pritchett, W.L. 1988. Site preparation effects on soil moisture and available nutrients in a pine plantation in the Florida flatwoods. Forest Science. 34:77-87.
- Cain, M.D.; Yaussy, D.A. 1984. Can hardwoods be eradicated from pine sites? Southern Journal of Applied Forestry. 8:7-13.
- Cantrell, Rick L. 1985. A guide to silvicultural herbicide use in the Southern United States. Alabama Agricultural Experiment Station and Auburn University School of Forestry. 592 pp.
- Danielovich, S.J.; Van Lear, D.H.; Cox, S.K.; Augspurger, M.K. 1987. Burning in southern Appalachian logging slash - Effects of residual vegetation and regrowth. In: Hay, R.L.; Woods, F.W.; DeSelm, H., eds. Proceedings of the central hardwood conference VI; 1987 February 24-26; Knoxville, TN: 91-98.
- DeBell, D.S.; Ralston, C.W. 1970. Release of nitrogen by burning light forest fuels. Proceedings of the Soil Science Society of America. 34:936-938.
- Dougherty, Phillip M.; Lowery, Robert F. 1991. Spot-size of herbaceous control impacts loblolly pine seedling survival and growth. Southern Journal of Applied Forestry. 15:193-199.
- Edwards, M.B.; Gonzalez, F.E. 1986. Forestry herbicide control of kudzu and japanese honeysuckle in loblolly pine sites in central Georgia. In: Proceedings of the Southern Weed Science Society annual meeting; 1986 January 20-22; Nashville, TN. Champaign, IL: Southern Weed Science Society: 272-275.
- Erdmann, Gayne G. 1967. Chemical weed control increases survival and growth in hardwood plantings. Res. Note NC-34. St. Paul, MN: U.S. Department of Agriculture, Forest Service North Central Forest Experiment Station. 4 pp.
- Garrett, H.E.; Thomas, M.W.; Pallardy, S.G. 1989. Susceptibility of sugar maple and oak to eleve foliar-applied herbicides. In: Rink, George; Budelsky, Carl A., eds. Proceedings of the centr hardwood forest conference; 1989 March 5-8; Carbondale, IL. Gen. Tech. Rep. NC-132. S

Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 81-85.

- George, David W.; Bowersox, Todd W.; McCormick, Larry H. 1991. Effectiveness of electric deer fences to protect planted seedlings in Pennsylvania. In: McCormick, Larry H.; Gottschalk, Kurt W., eds. Proceedings of the 8th central hardwood forest conference; 1991 March 4-6; University Park, PA. Radnor, PA: School of Forest Resources, Pennsylvania State University: 395-401.
- Hannah, Peter R. 1987. Regeneration methods for oaks. Northern Journal of Applied Forestry. 4:97-101.
- Horsley, S.B. 1988. Control of understory vegetation in Allegheny hardwood stands with Oust. Northern Journal of Applied Forestry. 5:261-262.
- Horsley, S.B. 1990. Control of grass and sedge in Allegheny hardwood stands with Roundup-residual herbicide tank mixes. Northern Journal of Applied Forestry. 7:124-129.
- Horsley, Stephen B.; McCormick, Larry H.; Groninger, John W. 1992. Effects of timing of Oust application on survival of hardwood seedlings. Northern Journal of Applied Forestry. 9:22-27.
- Johnson, James E. 1987. Evaluation of herbicides for site preparation in central Wisconsin. Northern Journal of Applied Forestry. 4:139-142.
- Johnson, Paul S.; Jacobs, Rodney D.; Martin, A. Jeff; Godel, Edwin D. 1989. Regenerating northern red oak: three successful case histories. Northern Journal of Applied Forestry. 6:174-178.
- Jones, Brad S.; Georges, Gary T.; Iverson, Richard D.; Butler-Fastelund, Marna C. 1986. Effects of sulfometuron on the silvicultural efficacy of common site preparation herbicides [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1986 December 2-4; Milwaukee, WI. Champaign, IL: North Central Weed Science Society: 106.
- Kennedy, Harvey E., Jr. 1981. Foliar nutrient concentrations and hardwood growth influenced by cultural treatments. Plant and Soil. 63:307-316.
- Kosinski, William G.; Holt, Harvey A. 1985. Herbicide tolerance for first-year weed control in woody species [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1985 December 10-12; St. Louis, MO. Champaign, IL: North Central Weed Science Society: 29.
- Krinard, Roger M.; Kennedy, Harvey E., Jr. 1987. Fift.een-year growth of six planted hardwood species on sharkey clay soil. Res. Note SO-336. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 pp.
- Langdon, O.G. 1981. Some effects of prescribed fire on understory vegetation in loblolly pine stands.
 In: Wood, G.W., ed. Prescribed fire and wildlife southern forests: proceedings of a symposium;
 1981 April 6-8; Myrtle Beach, SC. Georgetown, SC: Clemson University, Belle W. Baruch Forest Science Institute: 143-153.
- Lantange, D.O.; Burger, J.A. 1983. First-year survival and growth of loblolly pine (*Pinus taeda L.*) as affected by site preparation on the South Carolina and Georgia Piedmont. In: Proceedings of the 2nd biennial southern silvicultural research conference; 1984 November 4-5; Atlanta, GA. Gen. Tech. Rep. SE-24. Asheville, NC: U.S. Department of Agriculture, Forest Service: 5-10.
- Lautenschlager, R.A.; Schaertl, G. Richard. 1991. Electrical conductivity of five concentrations of two glyphosate-containing herbicides. Southern Journal of Applied Forestry. 15:85-88.
- Maeglin, Robert R. 1987. Juvenile wood, tension wood, and growth stress effects on processing hardwoods. In: Proceedings fift.eenth annual hardwood symposium of the Hardwood Research Council; 1987 May 10-12; Memphis, TN: 100-108.

286 Oak Regeneration Symposium Proceedings . . . September 1992

- Malac, Barry F.; Heeren, Robert D. 1979. Hardwood plantation management. Southern Journal of Applied Forestry. 3:3-6.
- McCormick, Larry H.; Allen, David H.; Groninger, John W.; Bowersox, Todd W. 1991. Effects of sulfometuron on the emergence and growth of seeded northern red oak, white ash, and white pine. Northern Journal of Applied Forestry. 8:9-11.
- McMinn, James H.; Nutter, Wade L. 1988. Season and intensity of whole-tree harvesting influence regeneration in the oak-pine type. Canadian Journal of Forest Research. 18:669-674.
- Miller, J.H. 1982. Hardwood control using pelleted herbicides and burning. In: Proceedings of the Southern Weed Science Society annual meeting; 1982 January 19-21; Atlanta, GA. Champaign, IL: Southern Weed Science Society: 210-215.
- Miller, James H. 1987. The use of herbicides in hardwood forestry. In: Proceedings of the Hardwood Research Council annual meeting; 1987 May 11-13; Memphis, TN: 31-52.
- Miller, James H. 1988. Guidelines for kudzu eradication treatments. In: Miller, J.H.; Mitchell, R.J., eds. A manual on ground applications of forestry herbicides. U.S. Department of Agriculture, Forest Service, Southern Region, Manage. Bull. R8-MB 21. Chapter 6.
- Miller, James H. 1990a. Streamline basal application of herbicide for small-stem hardwood control. Southern Journal of Applied Forestry. 14:161-165.
- Miller, James H. 1990b. Directed foliar sprays of forestry herbicides for loblolly pine release. Southern Journal of Applied Forestry. 14:199-206.
- Miller, James H.; Edwards, M. Boyd. 1985. Impacts of various intensities of site preparation on Piedmont soils aft.er 2 years. In: Proceedings of the 3rd biennial southern silvicultural research conference; 1984 November 7-8; Atlanta, GA. Gen. Tech. Rep. SO-54. New Orleans, LA: U.S. Department of Agriculture, Forest Service: 65-73.
- Miller, James H.; Burkhardt, E.C. 1987. A trial of herbicide treatments for enrichment plantings for cherrybark oak. In: Hay, R.L.; Woods, F.W.; DeSelm, H., eds. Proceedings of the central hardwood conference VI; 1987 February 24-26; Knoxville, TN: 53-58.
- Miller, James H.; Mitchell, Robert J., eds. 1988. A manual on ground applications of forestry herbicides. Manage. Bull. R8-MB 21. Atlanta, GA: U.S. Department of Agriculture, Forest Service. 316 pp.
- Miller, James H.; Bishop, Larry. 1989. Optimum timing for ground-applied forestry herbicides in the South. Manage. Bull. R8-MB 28. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Cooperative Forestry Technology Update. 2 pp.
- Miller, J.H.; Boyer, W.D. 1991. Effect of repeated biennial burns on soil properties and foliar nutrients in longleaf pine stands. In: Agronomy Abstracts, 1991 annual meeting; 1991 October 27-November 1; Denver, CO: American Society of Agronomy: 353.
- Minogue, P.J.; Lauer, D.K. 1992. Impact of broadcast burning on pine growth and hardwood density six years following site preparation with imazapyr. In: Proceedings of the Southern Weed Science Society annual meeting; 1992 January 20-22; Little Rock, AR. Champaign, IL: Southern Weed Science Society.
- Morris, Lawrence A.; Pritchett, William L.; Swindel, Benee F. 1983. Displacement of nutrients int windrows during site preparation of a flatwood forest. Soil Science Society of America Journa 47:591-594.
- Morris, Lawrence A.; Lowery, Robert F. 1988. Influence of site preparation on soil conditic affecting stand establishment and tree growth. Southern Journal of Applied Forestry. 12:170-1

- Newton, Michael; Knight, Fred B. 1981. Handbook of weed and insect control chemicals for forest resource managers. Beaverton, OR: Timber Press. 213 pp.
- Nix, L.E.; Cox, S.K. 1987. Cherrybark oak enrichment plantings appear successful aft.er seven years in South Carolina bottomlands. In: Proceedings of the 4th biennial southern silvicultural research conference; 1986 November 4-6; Asheville, NC. Gen. Tech. Rep. SE-42. Atlanta, GA: U.S. Department of Agriculture, Forest Service: 592-596.
- Nix, Larry E. 1989. Early release of bottomland oak enrichment plantings appears promising in South Carolina. In: Proceedings of the 5th biennial southern silvicultural research conference; 1988 November 1-3; Memphis, TN. Gen. Tech. Rep. SO-74. New Orleans, LA: U.S. Department of Agriculture, Forest Service: 379-383.
- Pancake, Dale C.; Miller, James H. 1990. Streamline rate and timing with triclopyr and imazapyr on seven woody species [Abstract]. In: Proceedings of the Southern Weed Science Society annual meeting; 1990 January 15-17; Atlanta, GA. Champaign, IL: Southern Weed Science Society: 222.
- Parker, J.C.; Amos, D.F. 1982. Subsoiling effects on four pine seed orchard soils. Soil Tillage Research. 2:53-60.
- Pham, C.H. 1987. Thinning young, natural hardwood regeneration with broadcast herbicide application. In: Hay, Ronald K., Dr.; Woods, Frank W., Dr.; DeSelm, Hal, Dr., eds. Proceedings of the central hardwood forest conference VI; 1987 February 24-26; Knoxville, TN. Knoxville, TN: University of Tennessee: 205-215.
- Plass, William T.; Green, Alan W. 1963. Preplanting treatments for brush old fields in southern Illinois. Res. Pap. CS-1. Columbus, OH: U.S. Department of Agriculture, Forest Service, Central States Forest Experiment Station. 8 pp.
- Pyle, S.L.; Krueger, W.A. 1984. Trumpetcreeper control with selected herbicides used in noncropped areas [Abstract]. In: Proceedings of the Southern Weed Science Society annual meeting; 1984 January 17-19; Hotsprings, AR. Champaign, IL: Southern Weed Science Society: 141.
- Regeher, David L.; Frey, David R. 1988. Selective control of japanese honeysuckle (Lonicera japonica). Weed Technology. 2:139-143.
- Russell, T.E. 1974. Weeding northern red oak plantations on the Cumberland Plateau. Res. Note SO-184. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 2 pp.
- Sam, Alan W.; Bowersox, Todd W.; McCormick, Larry H. 1986. Preemergence herbicide performance on forest soil. In: Dowson, J.O.; Majerus, K.A., eds. Proceedings of the central hardwood forest conference V; 1985 April 16-17; Urbana, IL: Univ. of IL. 47-52.
- Sanders, D.L.; Van Lear, D.H.; Guynn, D.C. 1987. Prescribed burning in mature pine-hardwood stands--effects on hardwoods and small mammals. In: Proceedings of the 4th biennial southern silvicultural research conference; 1986 November 4-6; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service: 93-96.
- Schmeckpeper, E.J.; Lea, R.; Phillips, D.; Jervis, L. 1987. Piedmont bottomland hardwood regeneration responds to preharvest japanese honeysuckle control. In: Proceedings of the 4th biennial southern silvicultural research conference; 1986 November 4-6; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service: 592-596.
- Seifert, John R. 1989a. The effects of chemical weed control on thirteen tree species at four testing sites in Indiana [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1989 December 5-7; Lexington, KY. Champaign, IL: North Central Weed Science Society: 62.

- Seifert, John R. 1989b. Two-year results of chemical weed control on nine hardwood tree species [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1989 December 5-7; Lexington, KY. Champaign, IL: North Central Weed Science Society: 63.
- Seifert, John R. 1990a. Site preparation results in the central hardwood region [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1990 December 11-13; Des Moines, IA. Champaign, IL: North Central Weed Science Society: 68.
- Seifert, John R. 1990b. Three year results of chemical and weed control on nine central hardwood species [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1990 December 11-13; Des Moines, IA. Champaign, IL: North Central Weed Science Society: 68.
- Seifert, J.R.; Pope, P.E.; Fischer, B.C. 1985. The effects of three levels of site preparation on planted swamp chestnut oak on a poorly drained site. In: Proceedings of the 3rd biennial southern silvicultural research conference; 1984 November 7-8; Atlanta, GA. Gen. Tech. Rep. SO-54. New Orleans, LA: U.S. Department of Agriculture, Forest Service: 53-56.
- Seifert, John R.; Holt, Harvey A. 1985. Herbicide test on 1-0 white ash, tulip poplar and red oak [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1985 December 10-12; St. Louis, MO. Champaign, IL: North Central Weed Science Society: 31.
- Shipman, R.D.; Prunty, T.J. 1988. Effects of herbicide residues on germination and early survival of red oak acorns. In: Proceedings, Northeastern Weed Science Society annual meeting; 1988 January 6-8; Hartford, CN. Champaign, IL: Northeastern Weed Science Society: 86-91.
- Shiver, Barry D.; Knowe, Steven A.; Kline, William N. 1990. Comparison of chemical site preparation treatments in the Georgia piedmont. Southern Journal of Applied Forestry. 14:24-32.
- Shiver, Barry D.; Knowe, Steven A.; Edwards, M. Boyd. 1991. Comparison of herbicide treatments for controlling common coastal plain flatwoods species. Southern Journal of Applied Forestry. 15:187-193.
- Smith, H. Clay. 1974. Forest management guidelines for controlling wild grapevines. Res. Pap. NE-548. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 15 pp.
- Stritzke, J.F.; Engle, D.M.; Bidwell, T.G. 1991. Paraquat plus aerial ignition for crown fire control of juniper [Abstract]. In: Proceedings of the Southern Weed Science Society annual meeting; 1991 January 14-16; San Antonio, TX. Champaign, IL: Southern Weed Science Society: 199.
- Thomas, M.W.; Miller, J.H.; Mitchell, R.J.; Minogue, P.J.; Gjerstad, D.M. 1989. Biological and economic comparison of mechanical and chemical alternatives for brush control in young pine stands-second year results [Abstract]. In: Proceedings of the Southern Weed Science Society annual meeting; 1989 January 16-18; Nashville, TN. Champaign, IL: Southern Weed Science Society: 247.
- Tuttle, C.L.; Golden, M.S.; Meldahl, R.S. 1985. Surface soil removal and herbicide treatment: effects on soil properties and loblolly pine early growth. Soil Science Society of America Journal. 49:1558-1562.
- von Althen, F.W. 1972. Stock age, planting method and weed control in red oak afforestation. Forestry Chronicle. 48:35-36.
- von Althen, F.W. 1987. Site preparation and weed control in hardwood afforestation in Ontario. In Hay, R.L.; Woods, F.W.; DeSelm, H., eds. Proceedings of the central hardwood conference VI 1987 February 24-26; Knoxville, TN: 75-80.

- Vose, James M.; Swank, Wayne T. 1992. Impacts of site preparation burning on aboveground biomass, nitrogen, and carbon pools in southern appalachian mixed pine/hardwood ecosystems [Abstract]. In: International conference on forest vegetation management "ecology, practice and policy" program and abstracts; 1992 April 27-May 1; Auburn University, AL. Auburn University, AL: School of Forestry and International Union of Forestry Research Organizations: 97.
- Wendel, G.W. 1980. Growth and survival of planted northern red oak seedlings, in West Virginia. Southern Journal of Applied Forestry. 4:49-54.
- Wendel, G.W.; Lamson, Neil I. 1987. Effects of herbicide release on the growth of 8- to 12-year-old hardwood crop trees. Res. Report 598. Radnor, PA: U.S. Department of Agriculture, Northeastern Forest Experiment Station. 6 pp.
- Wittwer, R.F.; Dougherty, P.M.; Cosby, D. 1986. Effects of ripping and herbicide site preparation treatments on loblolly pine seedling growth and survival. Southern Journal of Applied Forestry. 10:253-257.
- Woodrum, Walter Grover III. 1982. Response of bottomland-planted cherrybark oak seedlings to natural, mechanical, and chemical methods of competition control. Clemson, SC: Clemson University. 63 pp. Thesis.
- Wright, Galen M. 1986. Impact of chemical site preparation on survival and growth of five hardwood species. In: Dowson, J.O.; Majerus, K.A., eds. Proceedings of the central hardwood forest conference V; 1985 April 16-17; Urbana, IL: University of Illinois. 40-46.
- Wright, Galen M.; Holt, Harvey A. 1985. Effects of chemical weed control on hardwood plantation establishment [Abstract]. In: Proceedings of the North Central Weed Science Society annual meeting; 1985 December 10-12; St. Louis, MO. Champaign, IL: North Central Weed Science Society: 68.
- Yeiser, J.L. 1992. Post-establishment weed control for shortleaf pine. In: Proceedings of the shortleaf pine regeneration workshop; 1991 October 29-31; Little Rock, AR. New Orleans, LA: U.S. Department of Agriculture, Southern Forest Experiment Station: 77-100.
- Zedaker, Shepard M.; Lewis, James B.; Smith, David Wm.; Kreh, Richard E. 1987. Impact of season of harvest and site quality on cut-stump treatment of piedmont hardwoods. Southern Journal of Applied Forestry. 11:46-49.
- Zutter, Bruce R.; Nelson, Larry R.; Minogue, Patrick J.; Gjerstad, Dean H. 1987. Hardwood plantation growth following weed control using herbicides and cultivation. Southern Journal of Applied Forestry. 11:134-138.

Collection and Care of Acorns

F. T. Bonner, Southern Forest Experiment Station, USDA Forest Service, Starkville, MS 39759

ABSTRACT The biological characteristics of acorns present acute problems in the collection and care of these single-seeded fruits. Indices of maturity are not complicated, but they must be rigorously followed in collection operations. Acorns collected too soon before physiological maturity will seldom germinate normally. Another critical factor is the high natural moisture content of acorns and extreme sensitivity to desiccation that they exhibit when dried only slightly below these high moisture levels. Maintaining this high natural moisture content is the key to maintaining good acorn quality, both in transport and in storage. High moisture contents prohibit sub-freezing storage and ensure a rapid metabolism that requires adequate aeration. For species that exhibit dormancy, the recommended storage conditions of near-freezing temperatures and maximum moisture contents will substitute for stratification in most cases. For the most vigorous germination, however, these conditions must be balanced with good aeration.

INTRODUCTION The fruits of the genus *Quercus* are single-seeded nuts (Olson 1974). The biological characteristics of these nuts are different from those of most seeds, and these differences present acute problems in the collection and care of these fruits. Oaks of the United States belong to a group of species called "temperate recalcitrants." They are "temperate" because they grow primarily in the temperate zone; they are "recalcitrant" because their seeds do not tolerate desiccation below a critical moisture content (approximately 25 to 35 percent).

Seeds of other species, which can be desiccated to below 10 percent, are called "orthodox," which means that they behave nicely when desiccated and survive long periods of storage under proper conditions. The "orthodox" group contains all temperate zone conifers, and many important hardwoods, such as ash (*Fraxinus* spp.), black cherry (*Prunus serotina* Ehrh.), yellow-poplar (*Liriodendron tulipifera* L.), sweetgum (*Liquidambar styraciflua* L.), and sycamore (*Platanus occidentalis* L.).

The purpose of this paper is to review the crucial steps in collection and care of acorns, and to present the best current recommendations on how to collect good acorns and maintain their quality. Every step of this process is heavily influenced by the moisture relations and recalcitrant nature of acorns.

MATURITY INDICES Acorns should be collected when they are fully mature and not before. Unlike multi-seeded fruits, such as pine or yellow-poplar, single-seeded fruits generally will not complete maturation after separation from the tree. The best maturity indices for acorns are (Bonner and Vozzo 1987):

- 1. Color of the pericarp.
- 2. Ease of separation of acorns from cups.
- 3. Cup scar color.
- 4. Cotyledon color.

In red oaks the pericarps should have lost their green color and be primarily dark brown or black before collection. An occasional exception to this rule can be made for southern red (Q. falcata Michx.) and cherrybark (Q. falcata var. pagodifolia Ell.) oaks. Individual trees of these species may produce mature acorns with a greenish tint to their pericarps. In white oaks brown and black are also good mature pericarp indicator colors, but again there are exceptions. Acorns from certain trees of white (Q. alba L.) and swamp chestnut (Q. michauxii Nutt.) oaks may be fully mature when pericarps are still yellow or even a mottled yellow and green.

These color changes are related to moisture loss with maturation. In Mississippi, white oak acorn moisture peaks at about 65 percent in early September, then drops to between 50 and 55 percent at maturity (Bonner 1976). Water oak (Q. nigra L.) acorns in the same region exhibit their maximum moisture content in August (65 to 70 percent), which then decreases to 35 or 40 percent at maturity (Bonner 1974c).

When acorns are mature, their cups come away cleanly with only slight pressure. If attempts to remove the cups cause them to break apart and leave pieces attached to the acorn, then the acorns are not yet mature. This is a simple test to carry out when collecting from branches. Overcup oak (Q. lyrata Walt.) is an exception to this rule, as these acorns are disseminated with their enclosing cups attached. The cup tissue is full of small air spaces, which apparently allow the acorns to float and be spread by moving water.

In red oaks, the cup scars on mature acorns are "bright" in color. On acorns of southern red and cherrybark oaks the scars may be bright pink or orange when first exposed. These colors fade within a few days of cup loss, however, so many good acorns collected from the ground may not show these bright colors. This index is most useful in checking maturity of acorns still attached to trees.

The last test for maturity is to examine a cross-section of acorn for cotyledon color. Species with naturally high fat contents, such as water oak, should have dark yellow to orange cotyledons. A pale yellow or white cotyledon indicates immaturity. Species with low fat content, such as Shumard oak (Q. shumardii Buckl.) and all white oaks, should have creamy white or light yellow cotyledons. Immature coloration in these acorns is almost the same as that of mature acorns. The higher the fat content, the deeper the orange color of the cotyledons. Mature overcup oak acorns are 50 percent carbohydrate and less than 1 percent fat (Bonner 1974a), and have almost white cotyledons. Cutting acorns in half also provides an opportunity to assess insect damage in the field. If insect larvae are found in more

than 25 percent of the acorns, then collection crews should realize that additional acorns may be needed.

Because of the recalcitrant nature of acorns, much acorn quality is often lost POST-HARVEST between collection and storage. Acorns must be kept moist to maintain good seed CARE quality. They should be collected and transported in plastic bags or in containers that can be covered to reduce moisture loss, especially if extended travel in the back of trucks is required. The steps that are taken to inhibit moisture loss also can lead to the problem of overheating (Gosling 1989). Overheating must be avoided, especially when plastic bags are used. Acorns should be kept in the shade while awaiting transport, and during transport, trucks should be parked in the shade when not moving. If the weather is warm, dry, and/or windy, spray the acorns with water. These are small things, but they can help maintain seed quality. In our experience, a loss of 5 percent moisture can be tolerated, but additional desiccation can lower acorn quality. If acorns are dried too much during collection and transport, moisture can be easily replaced by immersing the acorns in water at room or cold-storage temperature (Gosling 1989). Immersion is, in most cases, a good practice, and it leads logically to cleaning, the next step in acorn care.

CLEANING As soon as possible after collection, all acorns should be immersed in water. This procedure serves two functions. First, it allows removal of leaves, cups, other trash, and insect-damaged acorns that float. Sound, healthy acorns typically sink in water. The exception to this rule is overcup oak, whose acorns always float with their large cups full of air spaces. Second, immersion helps maintain that all-important high seed moisture.

If conditions are extremely dry when acorns are collected from the ground, many good acorns will float initially. Under such conditions, acorns should be kept in the water for up to 24 hours to elevate their moisture contents and allow sound acorns to sink. Acorns collected from wet conditions should separate easily at initial floating. One should always cut samples of "floaters" and "sinkers" to determine the effectiveness of floation to remove bad acorns.

After flotation and removal of trash, the water should be drained away prior to storage. Insect control measures should be taken at this time. The two common methods of control are immersion in hot water (120 °F) for 40 minutes, and fumigation with methyl bromide or other recommended chemicals (Olson 1974). Both of these methods present considerable risk to acorns, and the best alternative may be to do nothing. Most infested acorns will be removed in flotation. The larvae do not attack intact acorns during storage, so infestation does not increase. Damage is further decreased when acorns are put into cold storage. The temperature change encourages larvae to emerge from the acorns to pupate, and they die in the bottom of the container. Additional larval emergence can be encouraged by moving the acorns from cold storage to room temperatures and bacl again. Larval feeding must destroy the embryonic axis to prevent germination, s damage solely to cotyledon tissue does not prevent germination and development of a normal seedling.

If acorns are to be sized, the separations should be done at this time. Most nurserymen do not size acorns, but increasing use of mechanical planting, both in nurseries and in direct seeding, may lead to a wider adoption of the practice. Round-hole screens that are used in air-screen cleaners may be used, especially for small acorns. Sizing can have advantages in nurseries through its effect on seedling uniformity. A positive correlation between acorn size and seedling size (height or leaf area) has been reported for (*Q. rubra* L.) (Farmer 1980), *Q. robur* L., and *Q. petraea* (Mattushka) Lieblein. (Kleinschmit and Svalba 1979).

STORAGE N

Most seed managers do not like to store acorns over long periods, because their size requires large refrigerated storage space and because viability declines each year. Acorns of most red oaks can be stored for 3 years without critical losses in viability (Bonner 1973), while most white oaks can be stored only 6 months without complete loss of viability. One solution to the problem is to plant acorns in the fall immediately after collection and avoid storage. For some nurseries and conditions, this option is a good one for next year's crop. Short-term storage under good conditions between collection and sowing is essential to maintain good acorn quality, however, and many managers would like to store extra acorns for use 1 or 2 years later. Since good storage practices for both purposes require the same facilities and procedures, the recommendations are the same.

Acorns of red oak species should be stored with their moisture contents at 30 percent or higher in temperatures near, but above, freezing (34 °F to 40 °F). Airtight storage is lethal, so containers must allow some gas exchange with the atmosphere while maintaining high acorn moisture levels (Bonner 1973). Polyethylene bags with a wall thickness of 4 to 10 mils are good. For large quantities of acorns, storage can be in drums, cans, or boxes with polyethylene bag liners. Container tops and liners should not be completely closed; this will allow sufficient gas exchange. If water collects in the bottoms of storage containers, it should be drained from acorns intended for storage longer than over winter.

With proper care, many southern red oaks should maintain good viability for at least 3 years (table 1). We have had good success in our laboratory with water, cherrybark, and Nuttall oaks (Q. nuttallii Palmer), but less success with Shumard and willow (Q. phellos L.) oaks. Similar methods were used by Farmer (1975) for successful storage of northern red and scarlet (Q. coccinea Muenchh.) oaks, and by Suszka and Tylkowski (1982) for northern red oak in Poland.

With few exceptions, white oak acorns cannot be stored longer than over winter (4 to 6 months) without complete loss of viability. For over-winter storage, the same methods outlined for red oak storage should generally be used. Thinner polyethylene (1.75 mil) or cloth bags may be advantageous for white oaks because of a need for greater aeration (Rink and Williams 1984). Schroeder and Walker (1987) reported excellent results in storage of bur oak (Q. macrocarpa Michx.) for 6 months at 34 °F and 44 percent acorn moisture in sealed plastic bags. No information was given on the thickness of the bags. Any reduction in acorn moisture significantly decreased germination capacity and rate. Tests in our laboratory (table 2) have provided some rare successes with storing white oaks species.

Moisture content remains a crucial factor throughout storage. With acorn moisture levels above 30 percent and temperatures above freezing, respiration proceeds at a rapid rate. This process gradually decreases acorn dry weight, causing small increases in the percentage of moisture over time (table 1). Schroeder and Walker (1987) found no increase in bur oak moisture content over 6 months of storage, but Gosling (1989) reported that English oak acorn moisture contents increased as much as 5 percent over 6 months in storage. The loss in dry weight is why a static state of equilibrium between internal acorn moisture and the storage atmosphere, such as we find in orthodox species, is never reached for acorns. Approximate equilibrium moisture contents have been determined for a few species (table 3), but these probably change over long storage periods. Note that white oak has much higher equilibrium levels than the two red oak species. This is because starch, the major storage food in white oak, is more hygroscopic than lipid, the major storage food in red oaks.

Original moisture content and storage	Germination 3 °C 8 °C		Final moisture		
period			content 3 °C 8 °C		
	5.0	0 C	50	00	
	Percent				
24 percent moisture					
6 months	80	76	27	28	
18 months	9	0	30	34	
30 months	25	24	32	33	
31 percent moisture					
6 months	99	99	34	34	
18 months	99	96	35	35	
30 months	81	71	36	36	
33 percent moisture					
6 months	100	98	34	32	
18 months	93	95	31	36	
30 months	94	34	37	40	

Table 1—Germination and moisture contents of cherrybark oak acorns stored in polyethylene bags at 3 $^{\circ}$ C and 8 $^{\circ}$ C¹

¹ Bonner (1973).

Table 2—Viability retention of various southern white oak acorns stored at 2 °C, high moisture content, and in polyethylene bags¹

Species		Germina	Germination	
	Original	6 mo.	1 yr.	2 yrs.
	Percent			
Q. alba, white	90+	7.0		
Q. virginiana, live	96.0		60.7	17.6
Q. muhlenbergii, chinkapin	91.3		39.0	2.0
Q. michauxii, swamp chestnut	86.1		65.1	2.0
Q. lyrata, overcup			95.8	

¹ Bonner and Vozzo 1987.

	Storage conditions			
Species	40-55% relative humidity 4-5 °C	95% relative humidity 4-5 ℃		
	Percent			
Shumard oak	13	32		
Water oak	17	29		
White oak	37	50		

Table 3—Equilibrium moisture contents of acorns for three southern oaks stored under two conditions of temperature and humidity¹

¹ Bonner and Vozzo 1987.

Between 70 and 75 percent of total acorn moisture in water and Shumard oaks is in the cotyledons and embryonic axes, while the comparable total for white oak is only 58 percent (Bonner 1974b). White oak pericarps are thicker than those of red oaks, and they retain more moisture. Cup scar vascular openings are major conduits for moisture uptake (Bonner 1968). Recent experiments in our laboratory on acorn desiccation indicate that these openings comprise the key pathway for moisture loss also. As acorns dry, the proximal end of the cotyledons (just beneath the cup scar) loses moisture first. Unless the pericarp splits, as it does at radicle emergence, the embryonic axis and the cotyledon tissue surrounding it (distal end) is the last area to be desiccated.

One method tested successfully for storing Nuttall oak acorns solved the drying problem nicely. Johnson (1979) stored Nuttall acorns for 6 months in drums of water maintained at 34 °F to 40° F. Similar results were obtained for water and cherrybark oak acorns in our laboratory for 5 months, but longer storage periods of 17 and 29 months led to considerable loss in viability.

Germination during storage has always been a problem for acorns, although not as great a problem as some might think. Southern white oaks have so little dormancy that they will germinate on the tree in extremely wet falls, so it is no wonder that they germinate profusely in storage. There seems to be an inverse relationship between degree of dormancy and germination in storage among red oak species. The conditions recommended for storage are the same ones normally prescribed for pretreatment (stratification) to overcome dormancy. Epicotyls usually do not appear, but radicles emerge and can grow as much as 8 in. in storage.

Microorganisms kill many radicle tips in storage, and many more are broken when sowing takes place. Secondary radicle development occurs in oaks, however, and other radicles should develop. These secondary radicles may even form sort of a multiple taproot system. Broken radicles did not adversely affect seedling production in nursery beds of cherrybark and Shumard oaks in Mississippi (Bonner 1982). Barden and Bowersox (1990) obtained similar results with northern red oak in Pennsylvania, but there were strong family differences. The common and proper response by seed managers is to do nothing to prevent this early germination. Decreasing acorn moisture slightly during storage of California black oak (*Q. kelloggii*) is reported to reduce the sprouting, yet not harm acorn quality (Tim Plumb, personal comm.). This approach should be studied for southern oaks. **CONCLUSIONS** Acorns are the most difficult of all temperate zone American seeds to collect and care for properly. Most of the problems are related to the naturally high moisture content of these seeds and the need to maintain those moisture levels to maximize seed quality. Desiccation during collection and transport must be avoided; a 5 percent loss of moisture can harm acorn quality. Acorns of many species can maintain viability for up to 3 years if they are stored a few degrees above freezing with high moisture contents and some gas exchange allowed. If seed managers recognize the moisture considerations and plan for their impact, loss of acorn quality can be minimized.

LITERATURE CITED Barden, Charles J.; Bowersox, Todd W. 1990. Genotype and radicle clipping influence northern red oak root growth capacity. [Abstract]. In: Fourth workshop on seedling physiology and growth problems in oak plantings; 1989 March 1-2; Columbus, OH. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 2 pp.

Bonner, F.T. 1968. Water uptake and germination of red oak acorns. Botanical Gazette. 129:83-85.

Bonner, F.T. 1973. Storing red oak acorns. Tree Planters Notes. 24(3): 12-13.

- Bonner, F.T. 1974a. Chemical components of some southern fruits and seeds. Res. Note SO-183. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 3 pp.
- Bonner, F.T. 1974b. Determining seed moisture in *Quercus*. Seed Science and Technology. 2:399-405.
- Bonner, F.T. 1974c. Maturation of acorns of cherrybark, water, and willow oaks. Forest Science. 20:238-242.
- Bonner, F.T. 1976. Maturation of Shumard and white oak acorns. Forest Science. 22:149-154.
- Bonner, F.T. 1982. The effect of damaged radicles of presprouted red oak acorns on seedling production. Tree Planters Notes. 33(4):13-15.
- Bonner, F.T.; Vozzo, J.A. 1987. Seed biology and technology of *Quercus*. Gen. Tech. Rep. SO-66. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 6 pp.
- Farmer, R.E., Jr. 1975. Long term storage of northern red and scarlet oak seed. The Plant Propagator. 21(1):11-14.
- Farmer, Robert E., Jr. 1980. Comparative analysis of 1st year growth in six deciduous tree species. Canadian Journal of Forest Research. 10:35-41.
- Gosling, Peter G. 1989. The effect of drying *Quercus robur* acorns to different moisture contents, followed by storage, either with or without imbibition. Forestry. 62(1):41-50.
- Johnson, R.L. 1979. A new method of storing Nuttall oak acorns over winter. Tree Planters Notes. 30(2):6-8.
- Kleinschmit, J.; Svolba, J. 1979. Moglichkeiten der zuchterischen Verbesserung von Steilund Traubeneichen (*Quercus robur* und *Quercus petraea*). III. Nachkommenschaftsprufung von Eichenzuchtbaumen. Allge. Forest-Jagdzeitung 150(6): 111-120: [Seed Abstracts 3(7): 1986 1980].

- Olson, David F., Jr. 1974. *Quercus* L. Oak. In: Seeds of woody plants in the United States. Agric. Handb. 450. Washington, DC: U.S. Department of Agriculture, Forest Service: 692-703.
- Rink, George; Williams, Robert D. 1984. Storage technique affects white oak acorn viability. Tree Planters Notes. 35(1):3-5.
- Suszka, Boleslaw; Tylkowski, T. 1982. Storage of acorns of the northern red oak (*Quercus borealis* Michx. = *Q. rubra* L.) over 1-5 winters. Arboretum Kornickie. 26:253-306.
- Schroeder, W.R.; Walker, D.S. 1987. Effects of moisture content and storage temperatures on germination of *Quercus macrocarpa* acorns. Journal of Environmental Horticulture. 5(1):22-24.

The Use of Tree Shelters and Underplanting for Oak Regeneration

James L. McConnell, Southern Region, USDA Forest Service, Atlanta, GA 30367

ABSTRACT Supplemental planting of oaks has been a success in many cases. It has also been a dismal failure in many other cases. Since successful stands of oak regeneration are often difficult to obtain, supplemental planting could be a method for increasing the desirable component.

A new tool recently surfaced in the United States is called the British "tree shelter." Tree shelters have been used in Great Britain for several years with good success. Initial trials in this country show that tree shelters contribute to better survival and increased initial growth. They will not, however, make a tree grow on a rock. Learning how to use tree shelters to the best advantage is still evolving in this country.

INTRODUCTION

Oak regeneration is seldom easy. Forest Service silviculturists are concerned because in many stands the oak component is being lost, especially when the stand is cut (George and Fischer 1991). Also, unusually large losses of northern red oak have been recorded in certain areas (Tainter and others 1984). The lack of adequate regeneration is a serious concern along the Appalachian Mountains in the eastern part of the region and especially to the west in the Ozark and Ouachita Highlands, where hardwoods, especially the oaks, are a major component of many stands (van Hees 1980, Beck 1983).

Even-age management by clearcutting has been the treatment of choice for many Forest Service silviculturists during the past several decades. Clearcutting without planting often did not result in a desirable species composition. Planting oaks, until recently, also did not result in notable success. Pine and hardwood species, other than oak, usually are more aggressive and invade good quality upland sites. All of this, despite concerned efforts, has resulted in a net loss of oak as a component of most upland hardwood stands.

Successes with clearcutting (Zobel and Davey, no year), seed tree cutting (Johnson 1976), and shelterwood methods (Loftis 1983) have been recorded. Others have tried the different regeneration systems on the same site (Johnson 1984). In most cases, however, when natural regeneration has been successful it has failed to increase the amount of oak in the new stand (Clark and Watt 1971). The authors also suggest that oak regeneration is more likely to fail on good sites than on poor sites.

The purpose of this paper is to discuss underplanting and tree shelters as means of increasing oak regeneration. Information in this area is limited. The current ban on clearcutting and the "New Perspectives" concept of forestry has added emphasis to the technique of underplanting hardwoods, especially oak, on National Forests.

Successful planting of oaks in an understory has been accomplished in the Missouri Ozarks (Johnson and others 1986). The prescription is a four-step process:

- 1. Prepare the site by controlling undesirable vegetation.
- 2. Create a shelterwood.
- 3. Plant.
- 4. Remove the shelterwood.

Step one reduces competition from woody plants. Unwanted woody plants less than 5.08 cm d.b.h. are controlled by the application of registered herbicides. The techniques used can be basal spray, injection, or sprayed cut stumps.

The second step is the creation of a shelterwood with about 55 to 65 percent stocking based on Gingrich's (1967) stocking relations. If the stand is already understocked, little need be done. If it consists of cull trees that are not merchantable, this step can be combined with step 1. Johnson states: "The purpose of the shelterwood is to provide a uniform tree cover that permits a moderate amount of light to reach the forest floor without greatly stimulating the growth of woody and herbaceous competition." Some undesirable stock may need to be left to achieve the desired uniform shade condition.

The third step is very important. Large nursery seedlings, 1.27 cm diameter or more measured 2.54 cm above the root collar, should be used. Most nurseries do not grow a seedling this large in 1 year. However, 1-0, 2-0, or 1-1 seedlings can be used as long as the desired size is obtained. Root pruning, 20.32 cm below the root collar and top pruning 20.32 cm above the root collar, is recommended. The top pruning should be done within 2 weeks of planting to allow root growth promoting substances in buds to be translocated to the roots.

The fourth step requires the complete removal of the shelterwood and must be accomplished in the dormant season 3 years after planting. If necessary, stumps of undesirable species in the shelterwood cut should be treated after harvest.

This technique is applicable in other areas but the number of trees planted per acre, time of planting, and competition control will vary and must be adjusted to meet local conditions.

Johnson (1984) also compared the responses of planted northern red oak in clearcuts and plots thinned to 60-percent stocking. The study was conducted in the Missouri Ozarks. Planting stock was divided into four size classes: (1) large 1-0 seedlings (>60 cm tall); (2) small 1-0 seedlings (30 to 60 cm tall); (3) 1 - 1 transplant seedlings (>30 cm tall); and (4) 1-year-old container grown seedlings (>30 cm tall).

Shoots were clipped on one half of the trees in each size class. After three field growing seasons, the overstory was removed on half of the underplanted plots. After 5 years, average survival was 84 percent.

Johnson states that, "Overall, planted oaks grew faster in clearcuts than under partial cuts. But because competitors also grow rapidly after clearcutting, planted trees must promptly initiate rapid shoot growth after clearcutting. Clipped planting stock in the underplant-release treatment demonstrated that capacity during the two years after clearcutting; trees planted directly into clearcuts did not." The reason for this may, in part, be due to the greater initial root growth per unit leaf area of transplants over seedlings.

After 5 years in the field, the most successful trees were clipped 1-1 seedlings with initial shoot diameters (2 cm above the root collar) of 10 mm or more that were underplanted and subsequently released. However, undercut seedlings may perform as well as 1-1 transplants (Johnson 1989).

It is important that additional studies have also been installed and are in progress. Results have not been reported. The Ozark National Forest has installed an administrative study designed to help that forest determine the optimum amount of overstory density to maintain above planted northern red oak seedlings (Smith and others 1984). "The objectives of the study are to evaluate the growth of underplanted 2-0 northern red oak that are released from one of three overstory densities (40, 60, or 80 percent stocking) 3 years after planting. In addition, the effects of the following three levels of understory competition control will be evaluated: (1) none, (2) one pre-planting application of herbicide, and (3) one pre-planting plus one post-planting mechanical control treatment of undesirable plants. Planting stock will include fall- and spring-lifted trees. Within both classes of stock, the effects of shoot removal 15 cm above the root collar will be evaluated by comparing spring removal, fall removal, and no removal. Seed source will also be a variable in the study. Four local sources will be used to assure and account for some genetic diversity."

The technique of underplanting hardwoods, especially oak, is not widely used by silviculturists. When I queried the Forest Service silviculturists in Region 8, the response was negative. Over the years several forests and districts have tried underplanting oaks of various species with little success. Most referred me to the work being done by Paul Johnson in Missouri.

In the present climate of limited clearcutting on the National Forests in Region 8, underplanting certainly has a role to play. If the oak component is declining, underplanting may be the only method available to maintain the species in many stands. Silviculturists would be well advised to learn more about the technique.

TREE SHELTERS Tree shelters are a relatively new tool in the United States. They were developed in Great Britain in 1979; therefore, the use in field situations has been short-term. Their use with hardwood trees is intended to increase survival, aid in establishment, and increase early growth. Tree shelters also provide protection from animal damage, herbicide applications, and offer some protection against mechanical and environmental damage—sun scald, wind, and ice (Potter 1991).

Tree shelters are transparent or translucent polypropylene plastic tubes, eithe round or square, of varying diameter and height, color, and design. They ar generally secured to the ground with a wood or metal stake of appropriate lengtl Page 363 of 404

Several types of tree shelters are commercially available from both U.S. and European manufacturers. Tree shelters should not be confused with tree guards which are made from metal or plastic mesh and are only intended to provide protection from various animals.

The plastic shelter creates a mini greenhouse around the tree. Temperature ranges, humidity, and CO_2 levels are increased while light levels are lowered (Rendle 1985). This effect promotes survival and rapid juvenile growth for the oaks. Under normal unsheltered conditions, oak seedlings usually develop very slowly.

Tree shelters also can serve an important function in planting oaks and other hardwoods in urban areas. On these incredibly harsh sites they protect young trees from both human traffic and groundcare equipment (American Forestry Association 1989).

On extremely arid sites tree shelters have been used to protect against high winds and blowing sand. Moisture stress from drying winds was also reduced. Bainbridge (1990) reports, with mesquite in the Colorado desert, that after 90 days seedling survival increased from 0 percent in plastic mesh to over 80 percent in tree shelters. Average height growth after 90 days was 0 cm using plastic mesh and 40 cm using tree shelters.

Increased growth and survival, however, seem to be larger in Great Britain than in the United States. Tree species and climate apparently influence the potential of hardwood seedlings in a tree shelter (Windell 1991). Nevertheless, the impressive results from both survival and growth in Great Britain and initial tests in the United States cannot be ignored.

Early Results After 3 years the mean height growth, in Great Britain, of sessile oak transplants in shelters was 142 cm compared with 45 cm in a mesh guard and 27 for unprotected trees. The average stem volume was 118 cm³, 37 cm³, and 19 cm³, respectively. There was no difference in growth in a range of sizes of conical- and cylindrical-shaped shelters. An 8-cm diameter cylinder is large enough for sessile oak (Tully 1985).

In another study in Great Britain, Rendle (1985) states: "The results of a three year investigation into the effects of tube shelters on microclimate and the growth of oak (Q. rubur) compared with field conditions, showed the environment of tubes has an increased temperature range, lower light levels and increased humidity when the tree is in leaf. Trees grow faster in height in tubes compared with the open but do not differ in terms of total dry weight production. However, the distribution of dry weight between stems, branches, and roots does differ."

At this time, results of studies in the United States are limited, but there is sufficient evidence to proceed quickly in determining just what role tree shelters should play in hardwood regeneration and management.

Manchester and others (1988) report that, after 2 years in North Carolina, chestnut oak grew to an average height of 47.5 cm in shelters and 50.8 cm without; white oak was 46.3 cm tall in shelters, 27.9 cm without; and northern red oak was 62.2

cm tall in shelters, 12.7 cm without. The survival rate after 2 years for trees without shelters was 31 percent; for trees with shelters, survival was 94 percent for all species. Chemical control of vegetation was applied in a 0.6-meter circle around each seedling 2 months following planting.

A test in Michigan using tree shelters shows that, after 2 years in a clearcut, northern red oak grew 42.9 cm in height with shelters and 23.4 cm with no shelter (Lantangne 1990). After planting, seedlings were clipped 17.8 cm above the groundline. Tree shelters were then installed. In this test northern red oak seedlings planted in tree shelters were 40 percent taller than unsheltered seedlings after one growing season and 42 percent taller after two growing seasons. The average increment in total height due to sheltering trees declined from a factor of 2.4 times during the first growing season to a factor of 1.8 times during the second growing season. No mention is made of survival for the planted trees.

Early test results from the United States are impressive though not as dramatic as those from Great Britain. The benefit of increased survival of planted hardwoods, especially oaks, seems to be as important as the increased juvenile growth.

Other studies are being installed throughout the United States. More and more companies are manufacturing and selling shelters of different designs. As a planting tool, they are likely to be with us for awhile; but, like the old snake oil drummer harking loudly that his potion will make hair grow on a billiard ball, tree shelters will not make a tree grow on a rock. Their useful function in this country remains to be determined.

SUMMARY Clear or white shelters increase success in planted shelterwoods or when used in an understory situation. Green or brown shelters, however, seem to be more environmentally acceptable to the public.

Tree shelters must be installed properly. They should be secured to a wood or metal stake which has been driven into the ground. The tree shelter, itself, should be pushed or driven into the loose soil about 1 in. deep at the bottom. If it isn't, the shelter will act as a chimney and cause the seedling to dry out. Tree shelters come in various lengths. An appropriate length should be used.

Public acceptance of tree shelters is still an open question. They are more acceptable if they blend into the background and are not planted in geometric patterns or straight lines. The plastic residue of the shelters, as they break down and become a part of the forest environment, may be a long-term problem. Some workers have proposed that the residue be picked up and hauled out of the forested setting. This would be an added cost unless different and more acceptable materials are found to construct the tree shelters.

The cost of tree shelters along with the cost of installation is relatively high per unit. Manchester (1991) reports of \$2 to \$3 per tree for total establishment on the Andrew Pickens Ranger District in South Carolina. Initial cost of the tree shelters may come down as more and more are manufactured and sold in this country. Also, the per-acre cost may be acceptable because less hardwood trees need to be planted if survival increases substantially as indicated in early tests. Page 365 of 404

Tree shelters definitely protect trees against deer and other browsing animals. The cost will probably be less than that of constructing deer-proof fence. Bears have destroyed small areas planted with tree shelters by knocking them down and tearing them into large pieces. Their presence seems to offend bears. Bluebirds and buntings may get into the tops of shelters and become entrapped. One manufacturer has devised a flexible polypropylene netting to place over the top of the tube.

Mulch mats and chemical weed control are more beneficial to seedling growth than fertilization alone when used in conjunction with tree shelters.

Pressure-treated and hardwood stakes can be used with tree shelters. The pressuretreated stakes will last longer in moist climates in the South.

LITERATURE CITED American Forestry Association. 1989. Tree shelters for rural releaf. Urban Forestry Forum, Box 2000, Washington, D.C., July/August.

Bainbridge, David A. 1990. Tubex treeshelters for tree establishment on extremely arid sites. Systems Ecology Research Group. San Diego State University, CA: 1-3.

Beck, Donald E. 1983. Planned hardwood regeneration. Forest Farmer. 43(1):18-19.

- Clark, F. Bryan; Watt, Richard F. 1971. Silvicultural methods for regenerating oaks. In: Doolittle, Warren T., Oak symposium proceedings; 1971 August 16-20; Morgantown, WV: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 37-43.
- Evans, J. 1984. Silviculture of Broadleaved Woodland. Forestry Commission Bulletin 62. London. Her Majesty's Stationary Office. 225 pp.
- George, David W.; Fischer, Burnell C. 1991. The occurrence of oak reproduction after clearcut harvesting on the Hoosier National Forest. Northern Journal of Applied Forestry. 8(4):144-146.
- Gingrich, S.F. 1967. Measuring and evaluating stocking and stand density in upland hardwood forests in the Central States. Forest Science. 21:38-53.
- Harris, Richard H. 1989. Arboriculture: world glimpses and ideas. Journal of Arboriculture. 15(3):62-66.
- Higgins, K.B.; Manders, P.T.; Lamb, A.J. 1989. The efficacy of microclimate shelter in improving seedling survival in re- establishment of the Clanwilliam cedar. South African Forestry Journal. 151(December): 1-5.
- Johnson, Paul S. 1979. Growth potential and field performance of planted oaks. Holt, Harvey A.; Fischer, Burnell C., Proceedings, regenerating oaks in upland hardwood forests; 1979 February 22-23; West Lafayette, IN: Purdue University. 113-119.
- Johnson, Paul S. 1984. Responses of planted northern red oak to three overstory treatments. Canadian Journal of Forest Research. 14(4):536-542.
- Johnson, Paul S.; Dale, Charles D.; Davidson, Kenneth R.; Law, Jay R. 1986. Planting northern red oak in the Missouri Ozarks: a prescription. Northern Journal of Applied Forestry. 3:66-68.
- Johnson, Paul S. 1989. Growing hardwood nursery stock for planting on forest sites with special reference to northern red oak. Northeastern Area Nurserymen's Conference. July 24-27. Peoria, IL: Northeastern State, Federal and Provincial Nursery Association: 46-62. (Available from Division of Forest Resources, Mason State Nursery, RR1, Box 235, Topeka, IL 61567).

- Johnson, R.L.; Krinard, R.M. 1976. Hardwood regeneration after seed tree cutting. Res. Pap. SO-123. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 9 pp.
- Kelty, Matthew J.; Kittredge, David B. 1986. Potential applications of British tree shelters to hardwood regeneration in the northeastern United States. Northern Journal of Applied Forestry. 3:173-174.
- Lantagne, Douglas O.; Ramm, Carl W.; Dickerson, Donald I. 1990. Tree shelters increase heights of planted oaks in a Michigan clearcut. Northern Journal of Applied Forestry. 7(1):24-26.
- Loftis, David L. 1983. Regenerating Southern Appalachian mixed hardwood stands with the shelterwood method. Southern Journal of Applied Forestry. 7(4):212-217.
- Manchester, Edwin H.; Roland, Felicia G.; Sims, Daniel H. 1988. Tree shelters show promise for oak regeneration. Management Bulletin R8-MB 25. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 2 pp.
- Manchester, Edwin. 1991. Using British Tree Shelters. Forest Management Update. Number 12. Morgantown, WV: U.S. Department of Agriculture, Forest Service, Northeastern Area, State and Private Forestry: 10-12.
- McGee, Charles E.; Loftis, David L. 1986. Planted oaks perform poorly in North Carolina and Tennessee. Northern Journal of Applied Forestry. 3:114-115.
- Morrow, Dan. 1989. Tree shelters: silvicultural tool of the future for hardwood regeneration in the northeast? Forest Management Update. Number 9. Morgantown, WV: U.S. Department of Agriculture, Forest Service, Northeastern Area, State and Private Forestry: 10-12.
- Olson, David F., Jr. 1972. Northern red oak plantings survive well in southern Appalachians. Tree Planters Notes. 23(1):16-18.
- Potter, Mark. 1986. Major innovation mark the 1986 tree-shelter scene. Forestry and British Timber. 15(10):19-23.
- Potter, M.J. 1991. Treeshelters. Forestry Commission Handbook 7. London, Her Majesty's Stationary Office Publications Centre. 48 pp.
- Rendle, Elizabeth L. 1985. The influence of tube shelters on microclimate and the growth of oak (Quercus rubur). In proceedings, 6th meeting of the National Hardwoods Program, 1985. Oxford Forestry Institute, Department of Plant Sciences, University of Oxford, Oxford (Oxfordshire): 8-16.
- Sandor, Ivan L.; Johnson, Paul S.; Rogers, Robert. 1984. Evaluating oak advance reproduction in the Missouri Ozarks. Res. Pap. NC-251. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 16 pp.
- Smith, O.D.; Johnson, Paul S.; Graney, David L. 1984. Methods of planting northern red oak on the Ozark National Forest. Unplublished administrative study: Russellville, AR: U.S. Deaprtment of Agriculture, Forest Service, Forest Supervisor, Ozark- St. Francis National Forest.
- Tainter, F.H.; Fraedrich, S.W.; Benson, D.M. 1984. The effect of climate on growth, decline, and death of northern red oaks in the western North Carolina Nanatahala Mountains. Castanea. 49:127-137.
- Trimble, G.R., Jr. 1972. Reproduction 7 years after seed-tree harvest cutting in Appalachian hardwoods. Res. Pap. NE-223. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 19 pp.
- Tully, Graham, 1983. Shelters improve the growth of young trees in the forest. Quarterly Journal of Forestry. 77:77-87.

- Tully, Graham, 1984. Tree shelter take the greenhouse to the tree. Forestry and British Timber. 13(10):17-22.
- Tully, Graham, 1985. The growth of young oak trees in shelters. Forestry. 58(2):180-195.
- van Hees, Willem W.S. 1980. Arkansas forests. Resource Bulletin SO-77. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 32 pp.
- Wendel, G.W. 1980. Growth and survival of planted northern red oak seedling in West Virginia. Southern Journal of Applied Forestry. 4(1):49-54.
- Windell, Keith; 1919. Tree shelters for seedling protection. 1E12L88-Seedling Protection. Missoula, MT: U.S. Department of Agriculture, Forest Service, Technology and Developemnt Center. 142 pp.
- Zobel, B.J.; Davey, C.B. No year. A simplified guide to hardwood management in the southeast. Raleigh, NC: School of Forest Resources, North Carolina State University. 21 pp.

Summary Papers

Oak Regeneration—Where Do We Go from Here?

R. C. Kellison, Hardwood Research Cooperative, North Carolina State University, Raleigh, NC 27695

ABSTRACT Care has to be exercised when referring to the regeneration problems of oak. Special problems are encountered with the regeneration of northern red oak in the Southern Appalachians and the Central States, but many red oaks regenerate without limitation.

On upland mesic sites where yellow-poplar and red maple are strong competitors, procedures are available to determine whether the desired reproduction will be obtained or not. Advance oak reproduction (greater than 4 ft. tall) must be present at the time of the regeneration treatment for the desired oak component to be competitive. Control of the undesired vegetation is required if the smaller oak reproduction is to gain dominance.

There will be a continuing demand for quality oak timber; thus, the question is "who will be producing that timber?" I do not foresee forest industry contributing significantly to the resource base because of their commitment to fiber production. Also, I do not foresee the nonindustrial private owner making a conscious effort to regenerate the species due to the long rotation ages needed to produce quality timber. It, therefore, will likely fall the responsibility of the public sector to husband northern red oak to keep it from becoming the "California Condor" of the Eastern deciduous forest in future centuries.

INTRODUCTION Fresh with a Ph.D. in forest genetics I was asked, some years ago, to prepare a paper on the genetics of the southern pines. Setting to the task, I did what I thought was a good job. Eager for a reviewer's comment, I asked the principal geneticist of Britian's Forestry Commission—who was visiting us at the time—for his assessment. Back came the reply: "You write a good paper but treat the subject as if the southern pines are the only pines in the world. You need to spend time outside the southern United States to learn the reproductive biology and silviculture of other pine ecosystems." That constructive criticism stung but, upon reflection, I set out to find a place to accommodate me. The choice was New Zealand, where I was employed by the New Zealand Forest Research Institute for 1 year. My job in that distant land was to increase seed production of Monterey pine (*P. radiata*) seed orchards.

Since New Zealand was (and still is) almost totally dependent upon exotic tree species for establishment of commercial plantations, and since the ink on my Ph.D. thesis was still wet, I arrived there with the hidden agenda of convincing the authorities to do away with monoculture, the top three species of which consisted of Monterey pine, Monterey pine, and Monterey pine. When I departed 1 year later it was I who had been converted. Even if catastrophe were to strike, the New Zealanders had determined that the benefits from intensively culturing Monterey pine were sufficiently greater over the second- and third-best exotics to warrant the extensive monoculture.

I cite this experience to suggest that the majority of the researchers and practitioners of oak management attending this symposium look beyond the Southern Appalachians and Central States to experience for themselves that the oak regeneration "problem" is not everywhere a problem. Research results obtained through the N.C. State-Industry Hardwood Research Cooperative¹ show that adequate oak reproduction has been obtained from east-central Mississippi (Bowling and Kellison 1983) to Florida (Leach and Ryan 1987) (Table 1), to Virginia (Heeren 1992, personal communication²). Despite the fact that the oaks identified in these research trials are predominately *Quercus laurifolia*, *Q. nigra*, and *Q. phellos*, they are oaks. They are highly desired as fiber species, and their fruit provides nutrition for a wide array of birds and animals, not to mention insects.

Treatment	Oaks	Gums	MCS ²	Elm	NCS ³
		Ster	ns per Acre ((no.)	
Clean cut	1,296	141	50	35	613
Without residual control	664	11	2	37	375
Residuals excluded	639	2	2	36	374
		Vo	lume per Acr	re (ft³)	
Clean cut	1,221	134	29	35	169
Without residual control	978	213	12	67	124
Residuals excluded	710	17	12	4	94

Table 1—Regeneration at 18 years following clean cut of a mature stand, with and without residual control, on a black-water river (Escambia River, Escambia County, Florida)¹

¹ Data courtesy of Champion International, Cantonment, FL.

² Miscellaneous Commercial Species.

³ Non-Commercial Species.

Oak regeneration, primarily of sprout origin, is also common in the high Piedmont Plateau and the foothills of the mountains on land of site quality less than 70 (50-year base) (Brenneman and Boyette 1978). There the species are primarily black, (Q. velutina), scarlet (Q. coccinea) and white (Q. alba) oak. In west-central Tennessee, chestnut oak (Q. montana) of site index 70 (50-year-base) dominates whole ecosystems.

¹ The Hardwood Research Cooperative consists of 11 industrial and three public agencies with land holdings in 10 Southern States.

² Hardwood Research Forester, Union Camp Corporation, Franklin, VA.

Similarly whole ecosystems of oak, primarily of black, northern red, and white oak, dominate the forests of the Boston Mountains where site index only occasionally exceeds 80 (50-year base) (Graney 1983). Even though an objective of this symposium is to identify the scope of the problem, I caution us not to be so quick to identify oak regeneration as everywhere a problem.

Regionally, the lack of oak regeneration, especially northern red oak (Q. rubra) is a problem, as identified by Loftis (1990a, 1990b). However, those problem areas occur on high oak site-index (more than 70 feet, 50-year base) land, especially where yellow-poplar (*Liriodendron tulipifera*) and red maple (*Acer rubrum*) are competitors. It is to these sites that the remainder of this paper is committed.

Quality red oak lumber has been in demand for furniture, barrel staves, building The Demand beams, farm implements, wood fuel, and other uses since the time of the European settlers. Following World War II, oak strip flooring became highly popular and great demand was again made on the natural resource. That fad lost momentum in the late 1950's, as a result of wall-to-wall carpet that covered the floor of lowgrade lumber or plywood. Another large drain on northern red oak began with the bicentennial celebration of the United States in 1976. Oak furniture became the item of choice, succeeding from pecan (Carya spp.) in the previous decades. In actuality, production of wood furniture inclusive of oak has been relatively constant over the past two decades, increasing from a base of 100.0 in 1972 to only 107.1 in 1990 (Nolley 1992). More significant, however, is the increase in the value of wood furniture imports during that period. Using the 1982 United States dollar as the base and adjusted for inflation, the value of imports increased from \$288 million in 1972 to \$1,949 million in 1990. In contrast, the value of domestic wood furniture shipments increased from \$6,029 million in 1972 to only \$6,455 million in 1990. These values correspond to an increase in the ratio of imports to domestic production from .05 in 1972 to .30 in 1990.

> The conclusion from these statistics might be that the tropical rain forests have contributed greatly to the wood furniture industry. Until recently, when a hue-andcry developed over destruction of the tropical rain forest, that conclusion would have been acceptable. However, the concept carries less validity when it is realized that the export of red oak logs from the United States has increased from 16,549 thousand board feet (mbf) in 1978 to 54,396 mbf in 1988. During that time, the price of Grade 1, 16-ft. logs increased from \$161/mbf in 1975 to a high of \$648/mbf in 1990, f.o.b. mill. The fact is that much of the log-price increase and much of the increase in wood-furniture imports are coming from logs that were exported from our shores.

> The trends during the past two centuries and especially during the past two decades suggest that there will continue to be a demand for northern red oak, for quality timber and wildlife values as well as to ensure that the venerable species remains a component of the Appalachian, Allegheny, Northern Hardwood, Central Hardwood, and Boston Mountain forests. The question remains: "Who will grow quality hardwoods and especially who will grow northern red oak?"

The three major categories of landowners of the Eastern deciduous forest-nonindustrial private, industrial private, and public-have holdings that

approximate a 70:20:10 ratio. Even with the high log prices I have cited, I do not see the nonindustrial private owner managing for northern red oak on any significant scale. Even if he is interested in regenerating the species, there will be reluctance to cut the timber until a family emergency arises or until the property changes ownership. And, then, there is always the question of economics. Not many of us, in the absence of an owned manufacturing facility, are willing to invest money in a project rife with potential disaster from fires, winds, insects, and diseases, that will mature in 60 to 120 years. Treasury bonds, even with the huge debt of the U.S. Government, are more attractive investments. Similarly, I do not see forest industry—especially those with pulp and paper as their objective-managing for the relatively long rotations required for the production of quality northern red oak timber. They are primarily interested in fiber to keep the huge pulp mills running, costing in excess of \$500 million. If the private sector were solely responsible for ensuring the supply of northern red oak, this valuable tree species might become the "California Condor" of the Eastern deciduous forest (Fralish and others 1991).

That leaves the public sector with its 10 percent of the Eastern deciduous forest. Twenty years ago most of us would have concluded that the future supply of domestic quality hardwoods, inclusive of northern red oak, would be coming from public lands, primarily from the National Forests. Today, that assumption is very much in question. The demand being made by the public on the National Forests for purposes other than timber production, and the New Perspectives approach being taken by the USDA Forest Service (Salwasser 1991), will pretty much exclude that source of timber. If those concerns are not enough to stop the timber harvest on public lands, an endangered species such as the northern spotted owl (Strix occidentalis) or another ploy governed by legislation will be used to accomplish the purpose. At best, only selective forestry will be practiced on the National Forests. Evaluation of the land then will be measured by social values rather than by Faustmann's land rent values (Davis and Johnson 1987). As long as the timber can be obtained elsewhere, we will become a net importer of quality hardwoods and products from which they are made. Western Europe has been in this mode since at least 1970. One wonders what the alternative will be when other countries also decide that their timber is more important to them for production and social value than it is for export.

In the Meantime Research has shown that northern red oak can be regenerated and made part of the succeeding stand on the high quality sites (site index 90, 50-year base for yellowpoplar) to which it is adapted in the Southern Appalachians (Loftis 1990a, 1990b). However, the success comes at great expense from control of yellow-poplar and red maple on sites where these species are common. The cause for these two species suppressing northern red oak, even when all were of equal size at time of regeneration or release, is unknown. However, it is speculated that the absence of ground fires and cattle grazing have contributed to the problem. Logic would suggest that many occurrences of fire, at periodic intervals, would be needed to control the earlier successional species of yellow-poplar and red maple while favoring northern red oak. Another probable cause for the demise of northern red oak at the expense of yellow-poplar and red maple is the loss of American chestnut (Castanea dentata) to chestnut blight (Endothia parisitica). Evidence shows that yellow-poplar was at a considerably lower frequency level at the time of the Page 374 of 404

predominance of American chestnut. Whether yellow-poplar was held in check by an allelopath or another competitive effect of American chestnut, or whether yellow-poplar is an opportunistic species that has only filled the void created by the death of American chestnut, is a matter of speculation.

Many of us at this symposium are addressing the idiosyncracies of oak regeneration and subsequent stand development as if they were new phenomena. Not so!! One of the most noted naturalists in the history of our country, Henry David Thoreau, made detailed observations and notes about the difficulties of obtaining desired oak reproduction (primarily northern red and scarlet) following removal of the parent stand in which good quantities of oak were present. He also concluded: "The time will soon come when we shall have to take special pains to secure and encourage the growth of white oaks, as we already must that of chestnuts for the most part. The oaks will be so scattered that there will not be enough to seed the ground rapidly and completely" (Torrey and Allen 1962, page 1698). The time was in the late 1850's, and the area was the environs of Walden's Pond, Massachusetts.

The naturalist observed that (red) oak regeneration under pine stands (*P. strobus* and *P. rigida*) was greatly superior to that under oak stands. The seedling oaks under overstory oaks were fewer than under pines; roots of these were old and decaying, and the shoots slender, feeble, and more or less prostrate under the leaves. The roots were not as fusiform shaped, and only one-tenth as many would develop into a second-growth stand as would develop under a pine stand. He further concluded: "If you expect oaks to succeed a dense and purely oak wood, you must depend almost entirely on sprouts, but they will succeed abundantly to pine where there is not an oak stump for them to sprout from" (page 1709).

Going further, the naturalist concluded: "Methinks you do not see numerous oaks of all ages and sizes in an old oak wood, but commonly large trees of about the same age and little ones like huckleberry bushes under your feet; and so commonly with pine woods. In either case, if the woods are well grown and dense, all the trees in them appear to have been planted at the same time. For aught I know, I would much rather have a young oak wood that had succeeded from pines than one that had succeeded from oaks, for they will make better trees, not only because the soil is new to them, but because they are all seedlings, while in the other case the greater part are sprouts."

I conclude from these remarks, from personal experience, and from the results of numerous researchers (Beck 1980, Carvell and Tryon 1961, Loftis 1990a, Sander and Clark 1971) that regenerating an oak stand following removal of the parent oak stand on upland mesic sites can be done, but it will be time-consuming and costly. The only logical chance of immediate success is to favor those stands with advance reproduction (Loftis 1990a) or those with high stump-sprouting potential (Johnson 1977). Models have been developed to serve as guidelines for successfully regenerating oaks in the Southern Appalachians and the Central States by the above-named scientists, and in the Mississippi Delta by Robert L. Johnson, former Project Leader of the USDA Forest Service, Southern Hardwoods Laboratory. These and other models are in need of refinement which can only be satisfactorily accomplished from input of research results being regenerated by the likes of the scientists represented at this symposium.

I also conclude that efforts to cause oaks to succeed themselves on upland mesic sites is unnatural, and that a reduction in tree and stand quality, and in site degradation, as suggested by Henry David Thoreau (Torrey and Allen 1962) could be a result on marginal sites. The total or partial harvests used today for timber removal and stand regeneration are greatly different from what would have been encountered in the original stands where repeated low intensity fires and grazing were common phenomena, or where conflagrations or wind would have likely completely destroyed the forest cover. In those situations, species succession would likely have played a major role, preventing one species from succeeding itself to the exclusion of other species. The benefits from species succession are the prevention of site degradation and pest build-up. The forest manager can learn a lesson in this respect from the agronomist who refrains from planting the same crop on a given soil for more than 2 successive years even when the option exists to replenish nutrients that are removed with each annual crop, and even when integrated pest management could conceivably hold epidemic levels of pests at bay. Still, the value of regenerating a valuable crop such as northern red oak, for biodiversity, timber, wildlife, aesthetic, watershed, and recreational purposes, may more than offset the potential site degradation and pest build-up problems. Northern red oak could conceivably become to the Southern Appalachians what Monterey pine is to New Zealand!

Confusing Terminology

I have used the term "forest succession" in the above paragraph. The purist would take exception to that term, equating it to the concept of *relay floristics* as proposed by Clements (1916). *Relay floristics* presupposes that plant formations follow a progressive succession, from pioneering species to a climax or stable community. Unfortunately, it also presupposes that the pioneer community modifies the environment to allow for the succeeding vegetation stage (sere), and that sere modifies the environment to allow for the next sere, etc., until the climax sere is reached. In my simplistic way of looking at plant succession, I do not presuppose that one plant community modifies the environment for the second sere, etc. I do hold to the principle that one sere succeeds another, but all of the species of the different seres could either be present from the beginning (*initial composition floristics*, as proposed by Egler, 1954) or they could be located there by plant migration, over time, through various dispersal mechanisms and the resiliency of seed banks and vegetative propagules (Gleason 1926).

I am of the opinion that we are confusing ourselves, to say nothing of the public, by our adamancy for labeling forest ecology events. The public only wants to know that we are practicing good forestry, and that should also be our first priority. Let us quit being so sensitive, for example, about the neophyte advocating the use of selective harvesting which, to us, equates to high grading. The neophyte does not have high grading in mind; he only wants the forest to be harvested (if it is to be harvested!) in a way that does not destroy his vista. We have the options to sometimes avoid those confrontations. Let us exercise those options, where appropriate, and call it good forestry without confusing our potential adversaries with terminology we *ourselves* do not fully comprehend.

CONCLUSION	Northern red oak is a struggling species. If it is to be maintained, special effort,
	at a high cost, will be required to ensure that either advance reproduction is present
	before the canopy of the parent stand is opened to any significant degree, or that
	competiting species such as yellow-poplar and red maple are controlled enough to
	allow development of subadvance reproduction. Numerous procedures have been
	detailed at this symposium to determine the conditions on which northern red oak
	can be successfully managed. In addition to the regeneration, the best silvicultural
	practices will be required to suppress the competition, including the use of fire,
	herbicides, and hand weeding. Because of the cost of the exercise, the practice
	will largely be confined to public lands. Additional research is needed to
	determine the cause of poor early development of northern red oak when compared
	to other species.

LITERATURE CITED Beck, D. E. 1980. Natural regeneration of southern hardwoods. In: McElwee, R.L.; Bell, N.H., eds. Regenerating the southern forest, tenth forest wildlife forum. Blacksburg, VA: Virginia Polytechnic Institute & State University: 132-142.

- Bowling, D.; Kellison, R. C. 1983. Bottomland hardwood stand development following clearcutting. Southern Jornal of Applied Forestry. 7(3):110-116.
- Boyette, W. G.; Brenneman, D. L. 1978. Five-year thinning results from a six-year-old upland mixed hardwood stand. NCFS Res. Note 34. North Carolina Forest Service, Raleigh. 12 pp.
- Carvell, K. L.; Tryon, E. H. 1961. The effect of environmental factors on the abundance of oak regeneration beneath mature oak stands. Forest Science. 7:98-105.
- Clements, F. E. 1916. Plant succession: An analysis of the development of vegetation. Washington Publ. No. 242. Carnegie Institute: 3-4.
- Davis, Lawrence S.; Johnson, K. Norman. 1987. Forest Management. New York: McGraw-Hill Book Company. 790 pp.
- Egler, F. E. 1954. Vegetation science concepts. 1. Initial floristic composition--a factor in old-field vegetation development. Vegetation 4:412-417.
- Fralish, J. S.; Crooks, F. B.; Chambers, J. L.; Harty, F. M. 1991. Comparison of presettlement, second-growth and old-growth forests on six site types in the Illinois Shawnee Hills. The American Midland Naturalists. 125(2):294-309.
- Gleason, H. A. 1926. The individualistic concept of the plant association. Bulletin of the Torrey Botanical Club. 53:7-26.
- Graney, David L. 1983. Intermediate oak stand management in the Boston Mountains. In: Proceedings, eleventh annual hardwood symposium of the hardwood research council. Cashiers, NC: 44-56.
- Johnson, P. S. 1977. Predicting oak stump sprouting and sprout development in the Missouri Ozarks. Res. Pap. NC-149. U. S. Department of Agriculture, Forest Service. 11 pp.
- Leach, Gregory N.; Ryan, Patrick P. 1987. Natural hardwood regeneration in the Escambia River bottom following logging. Res. Note GS-87-03. Cantonment, FL: Research and Development, Forest Products, Champion International Corporation. 5 pp.
- Loftis, David L. 1990a. Predicting post-harvest performance of advance red oak reproduction in the southern Appalachians. Forest Science. 36(4):908-916.

- Loftis, David L. 1990b. A shelterwood method for regenerating red oak in the Southern Appalachians. Forest Science. 36(4):917-920.
- Nolley, Jean W. 1992. Bulletin of Hardwood Market Statistics: Fall 1991. Res. Note NE-348. Radnor, PA: U. S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 47 pp.
- Salwasser, H. 1990. Gaining perspective: Forestry for the future. Journal of Forestry. 88(11):32-38,22.
- Sander, I. L. 1971. Height growth of new oak sprouts depends on size of advance reproduction. Res. Pap. NC-79. Journal of Foestry. 6 pp.
- Torrey, Bradford; Allen, Francis H., eds. 1962. The Journal of Henry D. Thoreau. Vol II (Chap. VIII XIV). New York: Dover Publications, Inc.

Oak Regeneration: A Summary

C. E. McGee, Center for Oak Studies, Department of Forestry, Wildlife, and Fisheries, University of Tennessee, Knoxville, TN 37996

David L. Loftis, Southeastern Forest Experiment Station, USDA Forest Service, Asheville, NC 28806

A FINAL WORD The oaks are a paradox. Under many conditions they reproduce themselves naturally in profusion. The genus has also been widely planted as an ornamental with great success. Yet, these Proceedings document that oak regeneration can be difficult to obtain. The purpose of the Symposium was to define trends and problems and to present the best available technology on oak regeneration. The purpose of this summary paper is to react to some of the observations, conclusions, and recommendations presented in the Proceedings. This paper will recognize consensus but will also point out exceptions and inconsistencies. Gaps in knowledge and lack of reliable procedures will also be briefly discussed.

The oak regeneration problem is recognized and described under two categories, natural regeneration and artificial regeneration. Natural stands of oaks have been observed under certain site conditions not to be regenerating. These failures in regeneration have been observed following an array of regeneration methods. The regeneration failures have ranged from almost complete loss of the oak component to a reduction in the relative dominance in the stand or the loss of one species of oak, when compared to the composition of the previous stand. The regeneration problem with oak plantations has been mostly one of slow growth of the oak seedlings following planting. Low early survival has also been an occasional problem. The general complaint, however, has been that we can get the planted oaks to live but we cannot get them to grow.

The oak regeneration problem has affected most of the oak species in the East and occurs throughout the entire Eastern oak forest. While the problem is reported to be widely distributed, both geographically and by species, the intensity of the problem is strongly associated with the quality of the site. The consensus in these Proceedings is that the difficulty of obtaining successful oak regeneration increases as the quality of the site increases. While some oak regeneration problems do occur on poor sites, e.g., plantation survival, the most severe problems have been reported on the very best sites.

The potential causes of the oak regeneration problem are apparent and are readily identified and described. However, it is much more difficult to determine and describe the specific actual cause of a particular oak regeneration failure. Causes of oak regeneration problems can usefully be separated into aggravating factors and limiting factors. A relatively long list of aggravating factors has been mentioned. These aggravating causes include flower production, acorn supply, seedling density, climate, insects, disease, animal depredation, fire, and flooding. However, there is consensus that the primary cause of general oak regeneration failure is the slow growth of oak reproduction relative to competing species. Even in plantations when competition has been eliminated, juvenile growth has usually been slow. The biological explanation as to why oak reproduction grows slowly even when released is an important subject covered in these Proceedings. A key statement suggests that oaks are not flexible and do not adapt well to a rapidly changing environment. Thus, increased supplies of light, water, materials, and space cannot be rapidly utilized. Similarly, saplings may not adapt to the adverse effects of sudden exposure to light and temperature extremes. Lack of flexibility, however, does not explain slow growth in plantations. The best explanation of poor plantation performance is that the planted oaks have had shoot/root ratios that were too high.

In any discussion of oak regeneration and oak regeneration problems, the potential sources of oak regeneration must be considered. By consensus, three categories of oak regeneration are recognized:

- (1) Seedlings that germinate and develop following a harvest or disturbance.
- (2) Sprouts growing from a stump.
- (3) Oak stems of various sizes present at the time of harvest.

The authors tend to discount the first category, seedlings that become established after harvest, as playing an important role in the regeneration of current oak stands. General agreement among authors is reached when the potential sources of future oak regenerations are considered. Seedlings established after disturbances are discounted because they do not compete with faster growing species and/or oak stump sprouts. Stump sprouts are frequently unimportant as a major or primary source of regeneration because large stumps usually do not sprout, and managed stands on medium-to-good sites will contain few small oak trees. Therefore, the consensus is that the regeneration of oak stands in the future is dependent upon developing a population of oak stems that can compete with other species following harvest or other disturbance, and this population will consists primarily of large advance reproduction. In the long term, however, successful oak regeneration requires flowering, acorn production, seedling establishment and development, and response to release.

A major purpose of the oak symposium was to provide the best available technical recommendations on how to regenerate oaks. Evaluation of existing oak regeneration potential in the existing stand is essential. Several authors discussed models and methods for evaluation of oak regeneration potential. The authors point out that the development of evaluation models is a complex, time-consuming task. The authors generally agree that, while broad evaluation principles apply, models and methods must be developed to fit local conditions and requirements. Most practitioners can find a method, with some modifications and adaptations, to describe the available sources of regeneration and their expected contribution to the next stand. The practitioner is, in many cases, left with making the decision as to whether existing regeneration potential is adequate or inadequate.

Once a stand or site has been evaluated for regeneration potential, harvest recommendations can be suited to the particular conditions. The authors generally agree that if and when regeneration potential is adequate, the overstory can be removed in one or more cuts. Several authors point out that single-tree selection will not provide for oak regeneration. Thus, the forester who wishes to regenerate oaks will ultimately use a regeneration method that would be classified as evenaged or as group selection.

When a stand or site to be regenerated to oak is judged to have inadequate regeneration potential, several options are available. Harvest could be postponed until the stand evaluation becomes favorable. However, without overt action the regeneration potential may not improve. Therefore, several authors suggest taking steps that will increase the oak regeneration potential. It is suggested that reduction of the understory and intermediate trees under a mature forest canopy will allow the development of existing oak seedlings. While this reduction in low competition will enhance the size of existing oaks, the procedure does not necessarily increase the number of oak seedlings present. It is generally concluded that, as the component of large advance oak reproduction increases, the potential of the stand to regenerate to oaks increases dramatically. Once the potential is judged adequate the overstory should be removed in one or more cuts.

Several papers in the Proceedings deal with the opportunities and problems related to the planting of seedlings and acorns as a means of establishing oak forests. The consensus is that the opportunity and outlook for successful oak planting and seeding is good. While most workers are optimistic about the future of oak planting and seeding, there remain serious words of caution. It is clear that great variation exists in the requirements for planting and seeding success between species, geographic areas, and topographic conditions. It is generally conceded that for the best chance of success the planting and seeding effort must be relatively intense.

Each of the authors emphasizes the particular requirement for an area or species. The following are general observations concerning planting or seeding oaks:

• Carefully match species and site.

• Provide adequate site preparation. Control competition and improve soil condition.

• Use only seedlings that meet known physical and physiological requirements.

• Plant or seed with care. Protect seedlings. Ensure adequate collection, storage, and treatment of seed and seedlings.

• Provide adequate follow-up competition control. Mow, cultivate, or treat with herbicides as needed.

• Anticipate slow early growth and be prepared to provide the treatment needed to keep the seedlings free to grow.

• Consider use of fertilizers, soil amendments, and tree shelters when local conditions are conducive to their use.

Even with careful attention to the above suggestions, planting or direct-seeding remains uncertain. But even with all the uncertainties and the expenses inherent

in the procedure, artificial regeneration may offer the best solution to many current and future oak regeneration problems.

The authors of these Proceedings have presented an impressive array of biological and silvicultural information and fused it into practical recommendations. An interested reader must be impressed by the complexity of the subject of oak regeneration and the skill and perserverance that has been needed to reach current levels of understanding. However, the observant reader will quickly perceive that there are many unanswered questions relevant to how oaks regenerated in the past and how best to regenerate them in the future. The following comments will highlight some of the major gaps in current knowledge of oaks:

- How did current stands of oak regenerate?
- What was the role of fire in establishment of existing oak stands?
- Is there a role for prescribed fire in oak regeneration?
- What factors influence oak flowering and fruit maturation?
- Can flower and acorn production be predicted?
- Why do oak seedlings grow so slowly?

• Why are direct seeding and planting efforts succeeding better in the bottomlands than in the uplands?

• Can natural regeneration methods and artificial regeneration methods be combined?

• What are the economic realities related to artificial regeneration of oaks?

Even as the foregoing questions indicate gaps in the knowledge of oak regeneration, it is clear from these Proceedings that much is known about the regeneration requirements of the oak genus. Continual efforts by forest scientists will provide answers to biological questions so that it is likely that oak regeneration in the future will be limited by economic and social constraints.

The objective of these Proceedings was to document trends, problems, and recommendations relevant to oak regeneration. The authors were asked to emphasize proven relationships and to make practical recommendations; they have accomplished their objective and fulfilled their commitment with an excellent aggregation of papers. We, the editors, commend them for their scholarship and thank them for their efforts.




The Forest Service, U.S. Department of Agriculture,

is dedicated to the principle of multiple use management of the Nation's forest resources for sustained yields of wood, water, forage, wildlife, and recreation. Through forestry research, cooperation with the States and private forest owners, and management of the National Forests and National Grasslands, it strives – as directed by Congress – to provide increasingly greater service to a growing Nation.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-5881 (voice) or (202) 720-7808 (TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, DC 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.



ALABAMA FORESTRY COMMISSION

INTRODUCTION

Quality and successful tree planting does not happen by chance. Success requires thought, planning, and attention to detail. From the first seed planted at the nursery to the last seedling planted in the field, all persons involved must accept responsibility for tree planting success or failure - negative impacts of poor seedling care and handling accumulate and cannot be reversed. Professional planting can be ruined by improper handling before the first seedling is placed in the ground. Likewise, fool-proof care and handling can be ruined by improper planting techniques. In either event, time, effort, and money is wasted.

<u>Seedling Care and Reforestation Standards</u> is intended to help landowners and vendors achieve quality and successful planting. So doing, newly planted seedlings will be prepared to capitalize on favorable growing conditions and better prepared to deal with growing conditions that are adverse.

TABLE OF CONTENTS

l.	Seedling Care and Handling Before Planting A. Transportation B. Storage 1) Refrigerated 2) Non-Refrigerated	1 1 1 1 1 1
II.	Evaluating Seedling Quality A. Physical Features I) Pine 2) Hardwood B. Seedling Condition	2 2 2 3 4
111.	Seedling Care and Handling During Planting A. On-Site Storage B. Time of Planting I) Planting Season 2) Weather Considerations	
IV.	 Proper Planting Techniques A. Method of Planting B. Equipment C. Seedling Care and Root Protection D. Culling Seedlings Before Planting E. Seedling Placement and Packing 	
V.	 Minimum Standards for Cost-Share Tree Plantings A. Tree Planting Inspections B. Defining a Correctly Planted Seedling C. Inspection Procedures 	8
VI.	Additional Steps to Insure Successful Planting A. Landowner Considerations B. Vendor Considerations	11 11 11
VII.	Appendix A. Proper Hand Planting Diagram B. Machine Planting Notes C. Four Days in the Life of a Mishandled Seedling	

I. Seedling Care and Handling Before Planting

Planting trees that will live starts with transportation and storage of seedtings between the nursery and planting site. Just like apples, oranges, and milk, **SEEDLINGS ARE PERISHABLE!** If they are mishandled during transportation and storage, **THEY WILL SPOIL AND DIE!** Follow the recommendations below to help insure your seedlings are at their best when planted.

Advice on Transportation

Use refrigerated transportation - especially when transportation will exceed 2-4 hours. Refrigerate at 33-38 F and 85.95% relative humidity.

Do not stack seedlings more than 3 bundles high. Use spacers to provide sufficient air space between stacks of bundles to reduce heat build-up.

Do not transport seedlings in truck beds containing fertilizer, chemicals, or fuel residues. These substances can be lethal to seedlings.

If seedlings must be hauled in open truck beds, use a tarpaulin to cover the seedlings. Leave sufficient air space between the top of the seedlings and the tarpaulin so air can circulate and reduce heat build-up.

Park in the shade when you stop to limit sun exposure on seedlings and reduce the chance of heat build-up.

Unload and properly store seedlings as soon as you reach your destination.

Advice on Storage

Use refrigerated storage, especially for storage in excess of 3-5 days.

store at 33-38 Fand maintain 85-95% humidity

do not stack seedlings more than 2 bundles high and use spacers to provide sufficient air space between, stacks of bundles and reduce heat build-up

for best results, plant as soon as possible within 4 weeks

store no longer than 8-12 weeks

From lifting to landowner pick-up, the Alabama Forestry Commission provides refrigerated transportation and storage for <u>all</u> their bareroot seedlings. With cooler facilities at E.A. Hauss Nurserv In Atmore. refrigerated vans, and local cooler facilities et various Alabama Forestry Commission District Offices across the state, landowners and tree planting vendors can pick up refrigerated seedlings within a two hour drive from anywhere 17 Alabama.

If non-refrigerated storage must be used:

store in a protected area such as a shed to avoid freezing, wind, and heat build-up provided that the area is not prone to overheating during direct sunlight

do not stack seedlings more than two bundles high and use spacers to provide sufficient air space between stacks of bundles to reduce heat build-up

for best results, plant as soon as possible

store no longer than 2-3 weeks if storage temperatures range from 38-5 0 F Bareroot longleaf pine seedlings do not store as well as other seedlings. Extra effort should be made to arrange for planting to occur within 10 days of lifting at the nursery.

store no longer than 3-5 days if storage temperatures range from 50-70 F

II. Evaluating Seedling Ouality

Seedling quality is determined by two seedling properties: physical features and condition. Evaluating both types of properties are critical to insure quality and successful planting and seedling establishment.

A. **Physical Features**

Pine Seedlings

Many physical features are used to assess seedling quality, such as stem length, root length, root collar diameter, and lateral root abundance. Unless altered during the handling and/or planting process, these physical features remain the same from the time a seedling is lifted at the nursery until it is planted and begins to grow. For this reason, seedlings should be inspected for acceptable physical features as soon as they are received from the nursery.

Nearly all survival studies show that balanced, medium-sized seedlings (7 to 13 inch stem height) with sturdy stems and well-developed, fibrous root systems have a higher survival



rate and make better initial growth than do either larger or smaller seedlings. One of the most important physical features that is vital to seedling survival is the relationship between the size of the root mass and the size of the foliage. Seedlings with far more foliage than roots (top-heavy seedlings) have a transpiring surface out of proportion to the absorbing capacity of the roots. Simply put, the foliage of top-heavy seedlings can transpire water out of the seedling faster than the roots can bring water in to the seedling. As a result, seedlings can out and die. Table 1 lists the minimum requirements for acceptable pine seedlings. You should evaluate your seedlings before planting. If most of your seedlings meet these requirements, remember to cull inferior seedlings as they are pulled for planting. *If the majority of your seedlings do not meet these minmums, notify the nursery where you obtained your trees- don 't overlook it.*

Table 1. Minimum Quality Measures for Pine Seedlings					
Quality Measure	Loblolly Slash <u>Shortleaf</u>	Longleaf,			
Stem Length	5"	l"(needle length)			
Root Collar Diameter	I/8"	318"			
Root Length	5"	5"			
Literal Roots	Abundant	Abundant			
Winter Buds		Present			
Mycorrhizae	Present	Present :			

Hardwood Seedlings

Two primary physical features are used to assess the quality of hardwood seedlings: root collar diameter and stem length. For hardwood seedlings, the following minimum standards should be met:

Root collar diameter	-	3/8 inch
Stem length	-	18 inches

B. Seedling Condition

Unlike physical features, the condition of both pine and hardwood seedlings can change and change rapidly when seedlings are improperly stored or transported. For this reason, seedlings should be monitored and inspected continuously - from pick-up until they are successfully planted. Indicators of seedling deterioration and unacceptable condition include:

sour smell due to heat build-up and fermentation

yellow needles

seedlings are warm to the touch due to heat build-up

bark slips off easily, especially on the roots

the cambium layer is yellow to brown

seedlings are molded

The cambium is a layer of soft tissue that lies between the bark and the woody portion of the seedling. Healthy cambium is glistening white in color and is a very good indicator of seedling condition. This layer can be inspected by peeling back the bark with your fingernail or a knife blade. If the color of the cambium is yellow or brown, do not plant the seedlings.

III Seedling Care and Handling During Planting

A. On-Site Seedling Storage

When seedlings are taken to the planting site proper care and handling often takes a back seat to the actual planting task. However, this period may present the harshest conditions seedlings will face before being planted.

Take the following precautions to minimize the adverse effects of on-site storage:

Do not carry to the site more seedlings than can be planted in one day.

Stack seedlings in the shade to reduce sun exposure. If natural shade is not available use a tarpaulin to provide shade. Freezing at 20 F for 6-24 hours slightly lowers the survival of loblolly and slash pine seedlings, and completely kills longleaf pine seedlings.

Cover seedling roots in open bundles and keep them moist.

Avoidexposing seedlings to freezing temperatures. If seedlings freeze allow them to thaw gradually before separating.

Preferred weather conditions for on-site storage are:

- air temperature of 33 75 F
- relative humidity > 50 %


Plant seedlings as soon as possible.

B. Time of Planting

Planting Season

Planting season for bareroot seedlings is December through March. Containerized seedlings allow for fall planting as well as during the bareroot planting season. For best results, once bareroot seedlings have hardened off in the nursery, they should be planted as early

in the planting season as possible (December-January). If at all possible, complete planting by March 15. to avoid the traditional dry weather of late winter and early spring. Survival of seedlings planted after March 15, will depend largely on adequate rainfall. Special care, handling and storage must be used for any planting beyond March 15.



Weather Considerations During Planting

Weather conditions at the time of planting can be beneficial or adverse to young seedlings. Cool, moist conditions help minimize seedling exposure during the planting process. High temperatures, low humidity, strong winds, and dry soil conditions can harm seedlings during the planting process, especially when coupled with improper care and handling. To the maximum degree possible, avoid planting when weather conditions are critical, and take special care to protect seedlings from exposure when weather conditions are marginal.

Category	Weather Conditions		
Normal	Temperature: Relative Humidity: Wind Speed: Soil Moisture:	33 - 75 F >50% <10 mph abundant	
Marginal	Temperature: Relative Humidity: Wind Speed: Soil Moisture:	76 - 85 F 30 - 50% 10 - 15 mph marginal	
Critical	Temperature: Relative Humidity: Wind Speed: Soil Moisture:	<32F or >85 F <30% >15 mph low to absent	

If seedlings must be planted when weather conditions are marginal or critical, take precautions to protect seedlings from potential damage.

overheating	edling bundles from sun exposure and dlings from drying out.
ow humidity: protect seed	
	dlings from drying out.
	dlings from drying out.
H U R WIND.	
	the second se
	ssible, wait for soil moisture conditions to
improve, es;	pecially if dry weather patterns are expected to
persist.	

IV. Proper Planting Techniques

A. Methods of Planting

There are two broad types of tree planting techniques - hand and machine planting. Either method is acceptable and can be performed successfully using a variety of planting tools or equipment when proper techniques are employed. Common examples include:

- 1. Pine Hand Planting Dibble bars - OST, KBC, and containerized Hoedads Planting shovels
- 2. Hardwood Hand Planting Shovel Po wef auger Post hole digger Hardwood planting dibble
- 3. Machine Planting Continuous furrow planters Intermittent planters Planting hole wheel Specialized container planters

B. Equipment

Proper planting techniques begin with using adequate equipment. Pine seedlings will require a hand planting tool or machine planter that will create a planting slit at least 8 inches deep (10 inches is preferred) and 3 inches wide toaccommodate pine seedling roots and allow proper root alignment. If the equipment being used is not adequate it should be replaced. Most hardwood seedlings have a more massive root systems than pine seedlings and therefore require larger planting holes for proper planting. Hardwood planting tools and machine planters should be capable of making a slit 12-15 inches deep and 4 inches wide to accommodate proper root alignment.

C. Seedling Care and Root Protection


Seedlings should be inspected when the seedling bundles are opened. If the majority of the seedlings do not meet the minimum criteria for acceptable seedlings found in Table 1, the seedlings should not be planted - contact the nursery. Also check seedling condition; do not plant seedlings that show signs of deterioration.

Seedling roots should have been root pruned at the nursery. Tap roots should range from 5-10 inches in length and lateral roots should be abundant. Roots should not be pruned in

the field. Both tap- and lateral root alignment will be acceptable if a planting slit of proper depth and width is constructed and proper planting techniques are followed. If a few seedlings have excessively long tap roots that impair proper root alignment, cull those seedlings as they are pulled from the planting bag. If the majority of your seedlings have excessively long (greater than 10 inches)

that impair proper planting, contact the nursery where you obtained your seedlings and seek a suitable remedy.

Seedlings should be carried in seedling bags, buckets, or other suitable containers that will allow the seedlings to be kept moist and protected. Seedlings should not be carried in the tree planters hand with the roots exposed.



D. Culling Seedlings before Planting

If seedlings are not culled when they are taken from the seedling bundles, individual inferior trees should be culled as they are pulled from the planting bag. Cull seedlings which do not meet the minimum criteria listed in Table 1 as well as seedlings with forked tops, evidence of fusiform rust, or other noted undesirable traits.

E. Seedling Placement and Packing

When hand planting, planting slits should be constructed according to Exhibit A in the Appendix. After a proper planting slit is made, insert the seedling until the tap root

reaches the bottom of the planting slit, then raise the seedling to the proper planting depth. After the seedling is raised to the proper planting depth and is in a vertical it position without leaning, close the planting slit and pack the soil around the planting hole. Proper packing can be determined for pine by grasping four of the terminal needles. If the seedling moves up when the four needles are pulled, the seedling is improperly packed. If the

	Proper Planting Depth
seedling	dwood, and all pines other than longleaf, s should be planted with their root collar 1-2
beneficia	pelow ground line. Deeper planting may be I on droughty soils such as deep sands and dry
	n either situaltion, the terminal bud should be 3 inches above ground line.
depth, e	pine is more sensitive to improper planting specially shallow planting. Longleaf seedlings
collar a	be planted at a depth that will leave the root or slightly below ground level once the soil he seedling settles.

seedling remains still and the four needles release, packing is suitable.

V. Minimum Standards for Cost-Share Tree Plantings

The Alabama Forestry Commission (AFC) provides recommendations for forestry practices performed under the state and federal cost-share programs and certifies that practices are correctly implemented. The minimumstandards used when checking cost-share tree plantings serve two primary purposes: 1) To help insure that landowners receive quality tree planting which results in a planted stand with a substantial chance of long term survival and success; and 2) to insure that cost-share funds are used wisely by funding quality and successful tree planting. This section addresses the criteria that AFC personnel use to determine if tree planting practices meet minimum standards.

A. Tree Planting Inspections

After tree planting is completed, the AFC will inspect the planting for the following:

I) Is the number of trees planted per acre within +25% of the number of trees per acre recommended in the treatment plan?

2)

per acre recommended in the treatment plan?

If the answer to both questions is YES, the tree planting is approved. If the answer to either of these questions in NO, the tree planting will not pass inspection.

B. Defining a "Correctly Planted Seedling"

For AFC inspection purposes, a correctly planted seedling is a seedling that meets the minimum criteria for an acceptable seedling as noted in Table 1, and which is planted in a manner **that reflects** quality planting. The following examples are indicative of seedlings that <u>do not meet</u> "Correctly Planted" criteria.

Cull Seedling	Seedling does not meet the minimum criteria nated in Table 1, or is deformed, forked, or disease infacted.
Debris in Planting Hole	Limbs, leaves, and/or other debris are in the planting hole in amounts that create air pockets.
Improper Depth	Seedling is planted too deep (terminal bud less then 3" from ground line) or too shallow (not collar above ground line).
Excessive Lean	Seedling has top lean in excess of 45 degrees from vertical or tap root lean in excess of 30 degrees from vertical.
Improper Packing	A pine seedling that does not pass the "four needle test" or a hardwood seedling that is excessively loose.
J-rooted Seedling	Improper root alignment due to a shallow planting hole or improper placement of the seedling when planted.
L-rooted Saedling	Improper root alignment within 5" of ground line due to a shallow planting hole, improper placement of the seedling in the hole when planted, or with machine planters due to improper furrow depth and/or excessive tractor speed.
Twisted or Balled Roots	Improper root alignment due to root damage during removal from the planting bale, due to field attempts at root pruning, or due to forcing roots into planting holes that are too shallow and/or too narrow.
Improper Pruning	Improper root condition due to removal of lateral roots and/or pruning of the tap root to a length less than 6*
Needle Discoloration	Evidence of a seedling that is highly likely to die. Causes can include improper transportation, storage, and handling prior to planting, as well as many other causes.
Dead Seedling	A seedling that is no longer alive.

C. Inspection Procedures

Cost-share tree plantings will be inspected by systematically sampling I/IOOth acre plots over the planted site. Depending on tract size, the number of plots sampled will be determined by the following:

Field or Tract Size Minimum #Plots

< 10 acres	3
10-25 acres	5
> 2 5 acres	1 plot/5 acres (no more than 30)

For each plot, the total number of trees planted will be recorded and used to determine if the number of trees planted par acre is within +25% of the number of trees recommended per acre in the treatment plan. *The number of trees planted per acre* is determined using the following formula:

(# of planted trees on all plots) \div (# of plots taken) x 100

Within each plot, detailed observations will be taken on the 5 trees closest to plot center to determine if they are correctly planted. Observations will be made both above and below ground for pine. All trees which appear correctly planted based on above ground observations will be excavated for below ground observation. Trees which are incorrectly 'planted based on above ground observations will not be excavated. For hardwood, only above ground observations will be taken. The status for each of the detailed observation trees will be recorded on the <u>Tree Plantina Inspection Form</u>. The information obtained will be used to determine the percentage of the detailed observation trees which are correctly planted. The percentage of trees correctly planted and the number of trees planted per acre will be used to determine if the number of trees correctly planted per acre is 75% or more of the number of trees per acre recommended in the treatment plan. The *number of trees* correctly *planted* per *acre* is determined using the following formula:

(# of correct observations) \div (# of observations) x (# of trees planted per acre)

If this number is greater than or equal to 75% of the number of trees recommended per acre in the treatment plan, the tree planting has passed inspection.

If the tree planting does not pass inspection, the landowner has until May 1, immediately following the date of planting to correct problems noted in the inspection and avoid practice disapproval. In the event of a dispute, appeals are to be made to the appropriate AFC District office.

VI. Additional Steps to Insure Successful Planting

The risk of planting failure due to human causes can be significantly reduced by following the guidelines in this handbook. The following considerations should be employed by the landowner and vendor to insure that the chance of tree planting success is maximized.

A. Landowner Considerations

PLAN AHEAD - don't wait until the last minute to perform site preparation, acquire seedlings, and hire a tree planting vendor. You may benefit by hiring someone to handle all the work from start to finish.

If you are planting trees under a cost-share program, make sure you understand the treatment plan prepared by the local AFC office. If you don't understand the plan or would like to do things differently, contact the AFC agent who prepared the plan before proceeding to insure that you will remain eligible for cost-share.

Ask others in your area who have hired tree planting vendors in the past. Use a vendor with a good reputation and that has the ability to transport and store seedlings correctly as outlined in this handbook.

Sign a contract with the vendor that clearly states the following:

- planting equipment must be capable of creating proper planting slits
- seedlings planted must meet the minimum criteria outlined in Table 1
- seedlings must be properly transported, stored and handled at all times
- planting must be performed during suitable weather conditions
- seedlings must be planted correctly as outlined in this handbook
- the date by which planting must be completed
- how many correctly planted seedlings per acre are required to receive payment

Visit the site during the planting operation and inspect the work.

B. Vendor Considerations

Make sure you have proper transportation, storage, and handling abilities before planting season arrives

Make sure planting equipment is suitable.

Train tree planters to transport, handle and plant trees properly

Maximize tree planting during the early part of planting season


Minimize planting during adverse weather conditions, and take special precautions when planting cannot be avoided

Use a crew foreman to insure that trees are properly handled and planted

Exhibit A - Proper Hand Planting Techniques

DIBBLE PLANTING


- Fully insert dibble straight down into the soil and pull back to open the planting hole.
- Remove dibble, insert seedling roots to the bottom of the planting hole, then raise root collar to the proper depth (I-2" below ground line).
- Insert dibble halfway, several inches from the base of the seedling. Push handle forward and close top of planting hole.
- 4) Fully insert dibble several inches down behind the seedling.
- 5) Pull back handle to close bottom of planting hole. Push handle forward to close the top.
- 6) Remove dibble and close-up the opening with your heel.



HOEDAD PLANTING

- 1) Strike blade vertically to full depth in the soil. Pull handle up a few inches to open the bottom of the planting hole.
- 2) Slide hand down handle, almost to the blade. Pull handle back and down to open the top of the planting hole. Insert seedling roots to the bottom of the planting hole and raise root collar to the proper depth (I-2" below ground line).
- Hold seedling in place and remove hoedad. Loose soil should fall into the hole and hold the seedling.
- Insert blade halfway, several inches from seedling, and push soil against the seedling.
- 5) Use foot to firm the soil against the seedling.










Exhibit B - Machine Planting Notes

For Safety, tractor and tree planting machines should have protective canopies.



OPERATIONAL CONSIDERATIONS

- 1. Use equipment with sufficient horsepower to pull mechanized tree planters.
- 2. Drawbar hookup should be no more than 14" from ground line.
- 3. Hitch arms for 3-point hitch planters should be the same length to assure straight planting rows.
- 4. Tractor speed should not exceed 2-2 miles per hour.
- 5. Packing wheels should be adjusted out for sandy soils and angled in for clay soils.

Event	Elapsed Time	Type Damage	Potential Loss in % Survival	Cumulative Potential Loss in % Survival
LIFTING - loss of roots	2 minutes	Physical	9 %	9 %
mycorrhizae - Seedlings placed in tubs, exposed to sur and wind prior to pickup	20 minutes	Dessication	7 %	16%
PACKING - Unloading seedlings in packing area and separation	2 minutes	Physical and Dessication	6 %	22%
- Exposure on the grading belt	30 seconds	Dessication	2 %	24%
- Rough handling	30 seconds	Physical	5 %	29%
SHIPPING - Bags improperly stacked	2 hours	Physical Overheating	7 %	36%
STORAGE - inadequate ventilation	2 days	Overheating	9 %	45%
PLANTING - exposure during planting	20 minutes	Dessication	5 %	50%

EXHIBIT C. Four Days in the Life of a Mishandled Seedling.

This chart depicts potential losses in the survivability of a mishandled seedlings. You can minimize these adverse effects by providing proper care and handling for your seedlings throughout the entire tree planting process.

For information on cooler locations contact Hauss Nursery at 1-251-368-4854

Literature Cited

Alabama Forestry Commission. 2002. Seedling Care and Reforestation Standards. <<u>http://forestry.state.al.us/publication/PDFs/SEEDLING%20CARE%20&%20REFORESTATION%20</u> <u>STANDARDS.PDF</u>> 2 Sept 2007.

- Allen, J.A., B.D.Keeland, J.A. Stanturf, A.F. Clewell and H.E.Kennedy Jr. 2001. A Guide to Bottomland Hardwood Restoration. General Technical Report SRS-40, USDA Forest Service, Southern Research Station. <<u>http://www.treesearch.fs.fed.us/pubs/2813</u>> Accessed 10 Nov 2007.
- Barry, D. and A.J. Kroll. 1999. A Phytosociological Description of a Remnant Bottomland Hardwood Forest in Denton, Texas. Lewisville Lake Environmental Learning Area (LLELA) Research Note 5. Lewisville, Texas, USA.
- Battaglia, L.L., P.R. Minchin and D.W. Pritchett. 2002. Sixteen Years of Old-field Succession and Reestablishment of a Bottomland Hardwood Forest in the Lower Mississippi Alluvial Valley. Wetlands Vol. 22, No.1: 1-17
- Bedker, P.J., J.G. O'Brien and M.E. Mielke. 1995. How to Prune Trees. USDA Forest Service NA-FR-01-95, Northeastern Area State and Private Forestry <<u>http://na.fs.fed.us/spfo/pubs/howtos/ht_prune/prun001.htm</u> > Accessed 10 Sept 2007.
- Clatterbuck, W.K. 2006. Treatments for Improving Degraded Hardwood Stands. University of Tennessee Extension Publication SP680. <<u>http://www.utextension.utk.edu/publications/spfiles/SP680.pdf</u>> Accessed 27 Sept 2007.
- Dey, D. C, J. M. Kabrick and M. Gold. 2006. The role of large container seedlings in afforesting oaks in Bottomlands. General Technical Report SRS-92, USDA Forest Service, Southern Research Station. <<u>http://www.treesearch.fs.fed.us/pubs/23381</u>> Accessed 10 Nov 2007.
- Gardiner, E. S. 2006. Early response of interplanted nuttall oak to release from an eastern cottonwood overstory. General Technical Report SRS-92, USDA Forest Service, Southern Research Station. <<u>http://www.treesearch.fs.fed.us/pubs/23474</u> > Accessed 10 Nov 2007.
- Gardiner, E.S., J.A. Stanturf and C.J. Schweitzer. 2004. An Afforestation System for Restoring Bottomland Hardwood Forests: Biomass Accumulation of Nuttall Oak Seedlings Interplanted Beneath Eastern Cottonwood. Restoratoin Ecology Vol. 12, No. 4:525-532. <<u>http://www.treesearch.fs.fed.us/pubs/7876</u> > Accessed 10 Nov 2007.
- Guo, Y., M.G. Shelton and E. Heitzman. 2002. Effects of Flood Duration and Depth on Germination of Cherrybark, Post, Southern, White and Willow Oak Acorns. General Technical Report SRS-48. USDA Forest Service, Southern Research Station. <<u>http://www.treesearch.fs.fed.us/pubs/3143</u>> Accessed 10 Nov 2007.
- Haynes, R.J., J.A. Allen, and E.C. Pendleton. 1988. Reestablishment of Bottomland Hardwood Forests on Disturbed Sites: An Annotated Bibliography. U.S. Fish and Wildlife Service Biological Report 88(42).

- Holcomb, S.S. 2001. An Examination of the Riparian Bottomland Forest in North Central Texas Through Ecology, History, Field Study and Computer Simulation. Thesis, University of North Texas, Denton, Texas, USA. <<u>http://www.ias.unt.edu/llela/documents/holcomb.pdf</u>> Accessed 15 Nov 2007.
- Hook, D.D. 1984. Waterlogging Tolerance of Lowland Tree Species of the South. Southern Journal of Applied Forestry Vol. 8, No. 3: 136-149 (14).
- Johnson, P.S. 1994. How to Manage Oak Forest for Acorn Production. Technical Brief TB-NC-1, USDA Forest Service, North Central Research Station, Silviculture and Ecology Upland Central Hardwood Forests Research Unit. < <u>http://ncrs.fs.fed.us/pubs/tb/tb1/techbrf1.html</u>> Accessed 18 Sept 2007.
- Kays, J.S. and R. Tjaden. 2002. Woodland Management: Measuring Your Forests. Maryland Cooperative Extension Fact Sheet 629, the University of Maryland System. <<u>http://extension.umd.edu/publications/PDFs/FS629.pdf</u> > Accessed 1 Sept 2007.
- Kindscher, K. and J. Holah. 1998. An Old-Growth Definition for Western Hardwood Gallery Forests. General Technical Report SRS–22, USDA Forest Service, Southern Research Station. <<u>http://www.treesearch.fs.fed.us/pubs/458</u>> Accessed 10 Nov 2007.
- Landowner Resource Centre and the Ontario Ministry of Natural Resources. 1994. Cover Crops Help Tree Seedlings Beat Weed Competition. <<u>http://www.lrconline.com/Extension_Notes_English/pdf/cvrcrps.pdf</u>> Accessed 20 Sept 2007.
- Landowner Resource Centre and the Ontario Ministry of Natural Resources. 1999. Choosing a Silviculture System. http://www.lrconline.com/Extension Notes English/pdf/cvrcrps.pdf> Accessed 20 Sept 2007.
- Landowner Resource Centre and the Ontario Ministry of Natural Resources. 1999. Do you have a Healthy Woodlot? <<u>http://www.mnr.gov.on.ca/MNR</u> <u>/forests/extension_notes/pdf/hlthywdlt.pdf</u>> Accessed 20 Sept 2007.
- Lockhart, B.R., J.S. Meadows and J.D. Hodges. 2005. Stand development patterns in southern bottomland hardwoods: Management considerations and research needs. The state of our understanding. University of Missouri. <<u>http://www.treesearch.fs.fed.us/pubs/20850</u>> Accessed 10 Nov 2007.
- Loftis, D. L. and C.E. McGee, editors. 1993. Oak Regeneration: Serious Problems Practical Recommendations (Symposium Proceedings). General Technical Report SE-84, USDA Forest Service, Southeastern Forest Experiment Station. <<u>http://www.treesearch.fs.fed.us/pubs/939</u>> Accessed 10 Nov 2007.

Loucks, W. 1987. Flood-Tolerant Trees. Journal of Forestry, Vol. 85, No. 3: 36-40(5).

Mercker, D. 2000. Crop Tree Release in Precommercial Hardwood Stands. University of Tennessee Extension Publication SP559. <<u>http://shelby.tennessee.edu/pubs/Shelby/SP559.pdf</u>> Accessed 10 Oct 2007.

- Minnesota Department of Natural Resources, and USDA Forest Service. 1996. How to Recognize Hazardous Defects in Trees. USDA Forest Service NA-FR-01-96. <<u>http://www.na.fs.fed.us/spfo/pubs/howtos/ht_haz/ht_haz.htm</u>> Accessed 10 Sept 2007.
- Moorhead, D.J. and K.D. Coder, editors. 1994. Southern Hardwood Management. USDA Forest Service Management Bulletin R8-MB 67, Southern Region. <<u>http://www.sref.info/publications/online_pubs/hardwood_management.pdf</u>> Accessed 8 Nov 2007.
- Nebeker, T. E., T.D. Leininger, and J.S. Meadows and M.D. Warriner. 2005. Thinning southern bottomland hardwoods stands: Insect and disease considerations. Ecology and management of bottomland hardwood systems: the state of our understanding, University of Missouri-Columbia. <<u>http://www.treesearch.fs.fed.us/pubs/20854</u>> Accessed 10 Nov 2007.
- Nix, Larry E. 2004. Herbicide Release of 4 Year Old, Naturally Regenerated Bottomland Oaks -- 10 Year Results. General Technical Report SRS–71, USDA Forest Service, Southern Research Station. <<u>http://www.treesearch.fs.fed.us/pubs/6692</u>> Accessed 10 Nov 2007.
- North Carolina Division of Forest Resources. 2006. Forest Soils and Site Index. <<u>http://www.dfr.state.nc.us/starting/starting_soilsandsiteindex.htm</u>> 1 Sept 2007.
- Peairs, S.E., A.W. Ezell, K.L. Belli and J.D. Hodges. 2004. A Comparison of Oak Regeneration Conditions Following Midstory Injection and Partial Overstory Removal in a Tombigbee River Terrace. General Technical Report SRS–71, USDA Forest Service, Southern Research Station. <<u>http://www.treesearch.fs.fed.us/pubs/6746</u>> Accessed 10 Nov 2007.
- Schweitzer, C. J. and J.A. Stanturf. 1999. A Comparison of Large-Scale Reforestation Techniques Commonly Used on Abandoned Fields in the Lower Mississippi Alluvial Valley. Paper presented at the Tenth Biennial Southern Silvicultural Research Conference, Shreveport, LA, February 18-18, 1999. <<u>http://www.treesearch.fs.fed.us/pubs/2194</u>
- Scherbarth, L.L. 2002. A Survey of Allelopathic and Other Chemical Interactions of Oaks (*Quercus* sp.). Colorado State University. <<u>http://www.colostate.edu/Depts/Entomology/courses/en570/papers_2002/scherbarth.pdf</u>> Accessed 20 Nov 2007.
- Stanturf, J.A., W.H. Conner, E.S. Gardiner, C.J. Schweitzer and A.W. Ezell. 2004. Recognizing and Overcoming Difficult Site Conditions for Afforestation of Bottomland Hardwoods. Ecological Restoration Vol. 22, No. 3. <<u>http://www.srs.fs.usda.gov/pubs/7391</u>> Accessed 10 Nov 2007.
- Stanturf, J.A., E.S. Gardiner, J.P. Shepard, C.J. Schweitzer, C. J. Portwood and L. Dorris. 2004. Restoring Bottomland Hardwood Forests: A Comparison of Four Techniques. Meeting the challenge: Silvicultural Research in a Changing World, June 14-18, Montpellier, France.

Stringer, J. 2006. Oak Shelterwood: A Technique to Improve Oak Regeneration. University of Tennessee

Extension Publication SP676. <<u>http://www.utextension.utk.edu/publications/spfiles/SP676.pdf</u>> Accessed 1 Oct 2007.

- Stringer, J. 2006. Two-age System and Deferment Harvest. University of Tennessee Extension Publication SP679. <<u>http://www.utextension.utk.edu/publications/spfiles/SP679.pdf</u>> Accessed 1 Oct 2007.
- Taylor, T.S., E.F. Edward and A.H. Chappelka. 2004. Improving Species Composition in Mismanaged Bottomland Hardwood Stands in Western Alabama. General Technical Report SRS-71, USDA Forest Service, Southern Research Station. <<u>http://www.srs.fs.usda.gov/pubs/6705</u>> Accessed 10 Nov 2007.
- USDA Forest Service and Centers for Urban and Interface Forestry. 1998. When the Forest Becomes a Community: A Forester's Handbook for the Wildland/Urban Interface. <<u>http://www.interfacesouth.usda.gov/resources/pubs/sgsfhandbook.pdf</u>> Accessed 8 Nov 2007.
- USDA Texas Forest Service. 2007. Reforestation Practices- Estimated Costs. <<u>http://txforestservice.tamu.edu/main/article.aspx?id=1802</u>> Accessed 1 Nov 2007.

USDA Texas Forest Service. 2007. Site Preparation. <<u>http://txforestservice.tamu.edu/main/article.aspx?id=1802</u>> Accessed 8 Nov 2007.